

ATAL-BELGOPROCESS JOINT VENTURE

Contract No. EP/SP/40/02

Low Level Radioactive Waste Storage Facility at Siu A Chau

**Seventeenth Environmental Monitoring and Audit Report
(Operation Phase)**

September 2013

Certified By



(Environmental Team Leader)

REMARKS:

The information supplied and contained within this report is, to the best of our knowledge, correct at the time of printing.

The Environmental Team Leader accepts no responsibility for changes made to this report by third parties.

Dr. John K.C. Leung
Department of Physics, The University of Hong Kong
Pokfulam Road, Hong Kong.
Tel: +852 2859 2858 Fax: +852 2471 8888
E-mail: jkcleung@hku.hk

TABLE OF CONTENTS

EXECUTIVE SUMMARY	1
BACKGROUND.....	2
PURPOSE OF THE REPORT	2
2. MONITORING RESULTS	3
AMBIENT γ DOSE RATES	5
SAND	8
GRASS	9
SEA WATER	10
MARINE ORGANISMS.....	11
AIRBORNE PARTICULATES.....	13
3. REPORT ON ELEVATED ENVIRONMENTAL RADIATION BACKGROUND	14
DOSE FOR RADIATION WORKERS	15
DOSE RATES AT UN-CONTROLLED AREAS	15
LIQUID EFFLUENT DISCHARGE.....	15
AIRBORNE EFFLUENT DISCHARGE.....	15
5. RESULT OF ENVIRONMENTAL COMPLIANCE AUDITS	16
APPENDIX 1.....	17
LIMIT LEVEL AND ACTION LEVEL	17
INVESTIGATION LEVEL.....	17

LIST OF TABLES

Table 2.1(a)	Ambient γ Dose Rates at 1 m above Ground
Table 2.1(b)	Comparison of Ambient γ Dose Rates with Previous Results
Table 2.2(a)	Activity Concentration of Some Major Radionuclides in Soil Samples
Table 2.2(b)	Comparison of Activities in Soil with Previous Results
Table 2.3(a)	Activity Concentration of Some Major Radionuclides in Sand Samples
Table 2.3(b)	Comparison of Activities in Sand with Previous Results
Table 2.4(a)	Activity Concentration of Gross α and β Emitters in Grass Samples
Table 2.4(b)	Comparison of α/β Activities in Grass Samples with Previous Results
Table 2.5(a)	Activity Concentration of Gross α/β Emitters in Sea Water Samples
Table 2.5(b)	Comparison of α/β Activities in Sea Water Samples with Previous Results
Table 2.6(a)	Activity Concentration of Gross α/β Emitters in Fish Samples
Table 2.6(b)	Comparison of α/β Activities in Fish Samples with Previous Results
Table 2.7(a)	Activity Concentration of Gross α/β Emitters in Sea Snail Samples
Table 2.7(b)	Comparison of α/β Activities in Sea Snails with Previous Results
Table 2.8(a)	Net Gross α/β Counts in Airborne Particulate Samples
Table 2.8(b)	Comparison of α/β in Airborne Particulate Samples with Previous Samples
Table A1.1	Limit Levels for Non-compliance and Action Levels
Table A1.2	Investigation Levels for Environmental Samples

LIST OF FIGURES

Figure 2.1	Locations of the Sampling Sites
------------	---------------------------------

EXECUTIVE SUMMARY

This is the seventh yearly report after the Facility has been operated for 8 years and it presents the results of the radiological monitoring work performed between August 23, 2012 and the date of sampling, which is August 27, 2013.

The Facility was operating smoothly over the years and there is no indication that the operation of the Facility has risen or will raise the environmental radiation level around the Facility. There is also no indication that the environment has been affected by the nuclear accident that occurred in Japan on March 2011.

All measurement results are normal and there is no sign of elevated radiation level in the environment.

This is the first time that one sea snail sample exhibits a β activity above the Investigation Level. But the exceedance is small and there is no indication that the water is contaminated as their α , β and γ activities are all normal.

Air-borne particulate samples were not measured in the past 2 years because the cloth for collecting the particulates were either severely damaged by weathering or lost completely. To rectify the situation, the operation team has carried out a routine check during the bi-weekly visits since Oct 2012. If any damaged cloth is found, it will be replaced. The damaged cloth will be stored for airborne particulate detection to ensure that all collected airborne particulates can be measured.

1. INTRODUCTION

Background

- 1.1 Various industrial, educational and medical facilities in Hong Kong have, for a number of years, used radioactive materials and generated radioactive waste. Most of the existing waste arisings were stored in disused air raid tunnels close to Queen's Road East in Wan Chai. Other arisings were stored temporarily (although in some cases for several years) at the point of use in educational institutions or hospitals.
- 1.2 A consultancy study in 1995 concluded that Siu A Chau was a suitable location for a purpose-built storage facility to which all waste will be transported, placed in stainless steel drums and stored.
- 1.3 In July 2003 ATAL-Belgoprocess Joint Venture Limited (ABJV) was awarded a contract to design, construct, and operate the LRWF at Siu A Chau for 10 years. Thereafter, the ABJV will transfer the waste management skills for this Facility to Hong Kong.
- 1.4 The LRWF was designed to have a storage vault that can initially store 260 drums of waste, each drum of 275 litres net capacity. The building also contains facilities for waste reception and repackaging waste, and administering the process. A jetty was built to provide marine access to the Facility.
- 1.5 The Facility is equipped with various radiation monitors inside the building specially installed for detecting all possible leakage of effluents from the building.
- 1.6 However, it is possible that minute activities may escape from detection and enter the biosphere, or an unexpected incidence would have resulted in a significant release of radionuclide from the Facility. It is one of the objectives of this environmental monitoring scheme to monitor whether in the long-term, the operation of the Facility will cause deterioration to the environment.

Purpose of the Report

- 1.7 This is the seventeenth EM&A (Operation Phase) report, which is also the seventh annual report on measurement results of environmental samples taken after the commencement of operation of the LRWF on July 28, 2005. This report covers the monitoring period from August 23, 2012 to the date of sampling which was August 27, 2013.
- 1.8 The requirements of the operation phase monitoring and audit; monitoring scheme and monitoring equipment and procedures have been fully described in the First EM&A (Operation Phase) Report. Please refer to that report for reference.
- 1.9 This report also covers the monitoring of personnel doses, the non-active areas of the Facility and the liquid and gaseous effluents.

2. MONITORING RESULTS

- 2.1 The sampling scheme remained unchanged. 15 in-situ ambient γ dose rates were measured. 3 soil samples; 3 sand samples; 3 grass samples; 8 seawater samples from 4 locations at two depths; 1 kg of sea snails; a few fish and 3 airborne particulate samples were collected and analysed as in previous monitoring. **Figure 2.1** shows the locations for taking various samples.
- 2.2 Ambient γ dose rates were taken at exactly the same locations and would give a true picture of the variation of the radiation environment if there were any.
- 2.3 Soil and grass samples were collected at more or less the same place as for the baseline. Since we need fresh surface soils that would have stored information of fallout since the commencement of the operation, the sampling sites shifted a little bit every time.
- 2.4 The uncertainties of the measurement results are given as standard deviation (SD) or standard uncertainty (SU). SD is given for individual sample and is calculated according to the number of counts recorded and assuming a normal distribution for the counts. SU is reported for each group of samples and it takes into account of the variance between samples. Please refer to the First EM&A Report (Operation Phase) for details.




Fig. 2.1 Locations of the Sampling Sites

(γ : Ambient gamma dose rate; S: Soil or Sand; W: Water; P: Air particulates)
(Grass sampling sites are the same as soil sampling sites)

Ambient γ Dose Rates

2.5 The measurement results are given in the last column in **Table 2.1(a)**. The 1st year results are average of all previous results measured within the 1st year. **Table 2.1(b)** also shows the results of the previous measurements for comparison. It is noted that the overall average value has not changed during the monitoring period.

Table 2.1(a) Ambient γ Dose Rates at 1 m above Ground

Location	γ Dose Rate ± 1 SD ($\mu\text{Sv h}^{-1}$)								
	Baseline (2005)	2006	2007	2008	2009	2010	2011	2012	2013
Boat	0.07	0.06	0.06	0.07	0.06	0.07	0.07	0.06	0.06 \pm 0.004
A	0.21	0.22	0.21	0.20	0.19	0.21	0.23	0.20	0.21 \pm 0.01
B	0.22	0.24	0.25	0.24	0.23	0.25	0.24	0.24	0.23 \pm 0.01
D	0.23	0.26	0.26	0.25	0.25	0.26	0.27	0.25	0.24 \pm 0.01
E	0.25	0.23	0.22	0.20	0.20	0.22	0.22	0.20	0.22 \pm 0.01
F	0.24	0.27	0.29	0.26	0.25	0.23	0.26	0.25	0.27 \pm 0.01
G	0.23	0.26	0.26	0.25	0.25	0.27	0.25	0.27	0.25 \pm 0.01
H	0.27	0.29	0.30	0.29	0.30	0.25	0.32	0.29	0.28 \pm 0.01
I	--	0.29	0.29	0.27	0.27	0.26	0.24	0.26	0.28 \pm 0.01
J	0.21	0.23	0.24	0.22	0.24	0.25	0.25	0.24	0.23 \pm 0.01
K	0.28	0.27	0.27	0.26	0.27	0.25	0.28	0.26	0.25 \pm 0.01
L	0.22	0.26	0.27	0.25	0.26	0.25	0.28	0.25	0.24 \pm 0.01
M	0.27	0.28	0.27	0.30	0.26	0.31	0.25	0.31	0.28 \pm 0.01
N	0.25	0.25	0.27	0.25	0.24	0.28	0.23	0.25	0.27 \pm 0.01
O	--	0.22	0.24	0.20	0.22	0.22	0.24	0.22	0.23 \pm 0.01
P	--	0.25	0.27	0.27	0.25	0.29	0.28	0.27	0.24 \pm 0.01

-- Not measured

2.6 No exceedance of Investigation Level was observed.

Table 2.1(b) Comparison of Ambient γ Dose Rates with Previous Results

EM&A Report No.	Mean γ Dose Rate ($\mu\text{Sv h}^{-1}$)	SU
1 (Baseline)	0.24	0.026
2	0.26	0.028
3	0.25	0.022
4	0.25	0.029
5	0.26	0.027
6	0.25	0.024
7	0.26	0.022
8	0.25	0.031
9	0.24	0.031
10	0.25	0.029
11 (2007)	0.26	0.025
12 (2008)	0.25	0.031
13 (2009)	0.25	0.028
14 (2010)	0.25	0.027
15 (2011)	0.26	0.026
16 (2012)	0.25	0.030
17 (2013)	0.25	0.023

2.7 The overall mean ambient γ dose-rate for this year is the same as those in previous years.

Soil

2.8 Soil samples were collected at 3 locations only, all from the undisturbed areas. These locations correspond to the passive air sampler locations which aim to detect dispersion of effluent leakages, if any, in the prevailing wind directions. The measurement results are given in **Table 2.2(a) & (b)**.

Table 2.2(a) Activity Concentration of Some Major Radionuclides in Soil Samples

Location	Collection Date	Activity Concentration (Bq kg ⁻¹)							
		²²⁶ Ra	SD	²²⁸ Th	SD	⁴⁰ K	SD	¹³⁷ Cs	SD
A	27 Aug 13	56.3	0.6	57.7	1.0	659	5.3	0.23	0.13
B	27 Aug 13	55.4	0.6	73.0	1.0	729	5.9	*	*
C	27 Aug 13	46.9	0.6	72.9	1.0	626	5.1	*	*

* Not detected

Table 2.2(b) Comparison of Activities in Soil Samples with Previous Results

EM&A Report No.	Mean Activity Concentration (Bq kg ⁻¹)							
	²²⁶ Ra	SU	²²⁸ Th	SU	⁴⁰ K	SU	¹³⁷ Cs	SU
1 (Baseline)	50.0	13.9	80.2	16.1	606	297	0.25	0.37
2	41.7	17.0	63.7	20.5	387	219	*	*
3	41.8	15.4	75.6	20.1	423	237	*	*
4	45.3	7.1	104.5	11.4	574	319	0.25	0.43
5	57.8	17.7	95.8	4.2	535	294	0.41	0.42
6	59.9	19.0	103.9	14.3	479	277	0.25	0.23
7	60.8	22.4	102.9	16.2	464	258	0.36	0.33
8	51.9	17.6	95.0	14.8	449	263	0.19	0.17
9	52.5	18.6	98.4	16.3	523	307	0.07	0.12
10	50.7	16.1	97.7	9.5	498	282	0.18	0.17
11 (2007)	52.8	15.7	106.8	16.9	483	253	0.27	0.01
12 (2008)	64.3	34.5	99.1	25.3	506	218	0.18	0.04
13 (2009)	59.3	12.0	116.0	12.0	474	199	*	*
14 (2010)	58.7	9.7	105	23.2	626	290	*	*
15 (2011)	48.9	31.5	52.5	27.1	504	249	0.1	0.03
16 (2012)	44.9	33.2	63.6	36.9	460	308	*	*
17 (2013)	52.9	5.2	67.9	8.8	671	52.6	0.07	0.13

* Not detected

2.9 No exceedance of Investigation Level is observed.

Sand

2.10 The measurement results are shown in Table 2.3(a) & (b).

Table 2.3(a) Activity Concentration of Some Major Radionuclides in Sand Samples

Location	Collection Date	Activity Concentration (Bq kg ⁻¹)					
		²²⁶ Ra	SD	²²⁸ Th	SD	⁴⁰ K	SD
A	27 Aug 13	19.7	0.4	14.5	0.6	353	3.8
B	27 Aug 13	21.8	0.4	14.0	0.6	257	3.3
C	27 Aug 13	21.3	0.4	13.5	0.6	274	3.4

Table 2.3(b) Comparison of Activities in Sand Samples with Previous Results

EM&A Report No.	Mean Activity Concentration (Bq kg ⁻¹)					
	²²⁶ Ra	SU	²²⁸ Th	SU	⁴⁰ K	SU
1 (Baseline)	18.8	4.4	21.6	5.5	576	106
2	11.1	3.8	12.8	5.0	357	100
3	11.4	3.2	13.2	4.4	382	141
4	28.3	22.8	24.5	17.4	360	165
5	23.3	12.7	25.6	17.9	323	117
6	20.8	8.0	25.8	18.0	329	95.7
7	30.2	24.8	24.3	17.0	320	173
8	15.4	4.6	15.4	4.1	246	30.5
9	14.5	1.2	17.3	5.8	380	99.1
10	18.4	1.7	18.5	2.4	377	124
11 (2007)	17.0	2.4	18.6	4.4	397	71.3
12 (2008)	18.0	4.7	16.5	1.7	382	20.6
13 (2009)	19.1	2.2	17.3	3.4	313	115
14 (2010)	14.3	2.2	12.5	0.3	301	71.5
15 (2011)	16.9	1.8	10.9	1.4	331	80
16 (2012)	19.2	2.5	13.9	3.0	273	49
17 (2013)	20.9	1.1	14.0	0.5	295	51.2

2.11 No exceedance of Investigation Level is observed.

Grass

2.12 Grass samples were collected in locations near to the soil samples. The measurement results are given in Table 2.4(a) & (b). The γ -spectra are identical to the background of the γ spectrometer and do not reveal the presence of any significant γ -emitting radionuclides, hence they are not reported here.

Table 2.4(a) Activity Concentration of Gross α and β Emitters in Grass Samples

Location	Collection Date	α Activity* (Bq g ⁻¹)	SD (Bq g ⁻¹)	β Activity* (Bq g ⁻¹)	SD (Bq g ⁻¹)
A	27 Aug 13	0.02	0.003	0.268	0.005
B	27 Aug 13	0.015	0.003	0.212	0.005
C	27 Aug 13	0.043	0.004	0.339	0.006

* Bq g⁻¹ refers to dry mass of grass

Table 2.4(b) Comparison of α/β Activities in Grass with Previous Results

EM&A Report No.	Mean α Activity (Bq g ⁻¹)	SU (Bq g ⁻¹)	Mean β Activity (Bq g ⁻¹)	SU (Bq g ⁻¹)
1 (Baseline)	0.083	0.044	0.33	0.03
2	0.037	0.012	0.25	0.01
3	0.081	0.017	0.30	0.10
4	0.093	0.009	0.26	0.03
5	0.084	0.020	0.23	0.04
6	0.081	0.056	0.22	0.09
7	0.077	0.046	0.25	0.08
8	0.068	0.047	0.28	0.05
9	0.050	0.023	0.29	0.02
10	0.051	0.008	0.40	0.07
11 (2007)	0.030	0.022	0.27	0.06
12 (2008)	0.012	0.020	0.17	0.04
13 (2009)	0.014	0.016	0.10	0.03
14 (2010)	0.038	0.027	0.21	0.04
15 (2011)	0.021	0.019	0.15	0.03
16 (2012)	0.022	0.022	0.10	0.03
17 (2013)	0.026	0.015	0.27	0.06

2.13 No exceedance of Investigation Level is observed.

Sea Water

- 2.14 The same 4 locations were chosen to collect the water samples at 2 depths. The measurement results are given in **Table 2.5(a) & (b)**.
- 2.15 Similar to grass samples, the γ spectra are not reported. There is no sign of presence of γ emitters.
- 2.16 No exceedance of Investigation Level is observed.

Table 2-5(a) Activity Concentration of Gross α/β Emitters in Sea Water Samples

Location	Collection Date	Water Depth (m)	α Activity (Bq L ⁻¹)	SD (Bq L ⁻¹)	β Activity (Bq L ⁻¹)	SD (Bq L ⁻¹)
A	27 Aug 13	1	0.00 [#]	0.00	2.92	0.72
		3.5	0.00 [#]	0.00	1.32	0.71
B	27 Aug 13	1	0.00 [#]	0.00	1.48	1.25
		6.5	0.00 [#]	0.00	2.92	0.72
C	27 Aug 13	1	0.00 [#]	0.00	3.58	0.69
		7.5	0.00 [#]	0.00	4.76	0.69
D	27 Aug 13	1	0.00 [#]	0.00	3.04	0.69
		5	0.00 [#]	0.00	2.02	0.71

[#] Below minimum detectable activity of 1.5 Bq L⁻¹ for α and 1.0 Bq L⁻¹ for β .

Table 2.5(b) Comparison of α/β Activities in Sea Water with Previous Results

EM&A Report No.	Mean α Activity (Bq L ⁻¹)	SU (Bq L ⁻¹)	Mean β Activity (Bq L ⁻¹)	SU (Bq L ⁻¹)
1 (Baseline)	0.77	0.25	7.20	0.70
2	0.49	0.47	6.10	0.46
3	0.57	0.21	7.43	0.80
4	0.71	0.50	7.00	0.81
5	0.92	0.44	6.15	0.64
6	0.63	0.28	6.99	0.37
7	0.25	0.28	6.30	0.45
8	0.19	0.23	5.84	1.34
9	0.32	0.29	5.21	0.38
10	0.70	0.35	8.35	2.19
11 (2007)	0.00	0.00	2.35	0.21
12 (2008)	0.00	0.00	4.08	0.42
13 (2009)	0.32	0.29	5.44	1.27
14 (2010)	0.00	0.00	4.80	0.41
15 (2011)	0.14	0.21	2.88	1.39
16 (2012)	0.03	0.07	3.74	0.96
17 (2013)	0.00	0.00	2.76	1.14

Marine Organisms

- 2.17 Fishes were caught along the jetty and sea snails were collected randomly along the shores.
- 2.18 The measurement results are given in Table 2.6(a) & (b) and Table 2.7(a) & (b) for the gross α/β activities in fish and sea snails respectively.

Table 2.6(a) Activity Concentration of Gross α/β Emitters in Fish Samples

Sample	Collection Date	α Activity* (Bq g ⁻¹)	SD (Bq g ⁻¹)	β Activity* (Bq g ⁻¹)	SD (Bq g ⁻¹)
1	27 Aug 13	0.000 [#]	0.000	0.04	0.001
2	27 Aug 13	0.000 [#]	0.000	0.04	0.001
3	27 Aug 13	0.000 [#]	0.000	0.04	0.001

* Bq g⁻¹ refers to wet mass of fish flesh.

[#] Below minimum detectable α activity of 0.008 Bq g⁻¹.

Table 2.6(b) Comparison of α/β Activities in Fish Samples with Previous Results

EM&A Report No.	Mean α Activity (Bq g ⁻¹)	SU (Bq g ⁻¹)	Mean β Activity (Bq g ⁻¹)	SU (Bq g ⁻¹)
1 (Baseline)	0.0093	0.004	0.068	0.003
2	0.0068	0.004	0.16	0.15
3	0.0116	0.005	0.026	0.006
4	0.0066	0.004	0.065	0.005
5	0.0040	0.004	0.056	0.010
6	0.0069	0.002	0.063	0.002
7	0.0120	0.021	0.047	0.035
8	0.0037	0.002	0.074	0.006
9	0.0100	0.004	0.062	0.050
10	0.0060	0.005	0.078	0.007
11 (2007)	0.0003	0.001	0.055	0.012
12 (2008)	0.0000	0.000	0.067	0.003
13 (2009)	0.0075	0.002	0.079	0.000
14 (2010)	0.0030	0.003	0.111	0.023
15 (2011)	0.0032	0.001	0.040	0.001
16 (2012)	0.0000	0.000	0.027	0.004
17 (2013)	0.0000	0.000	0.040	0.000

- 2.19 No exceedance is observed.

Table 2.7(a) Activity Concentration of Gross α/β Emitters in Sea Snail Samples

Sample	Collection Date	α Activity* (Bq g ⁻¹)	SD (Bq g ⁻¹)	β Activity* (Bq g ⁻¹)	SD (Bq g ⁻¹)
1	27 Aug 13	0.001 [#]	0.001	0.065	0.001
2	27 Aug 13	0.000 [#]	0.000	0.099	0.001

* Bq g⁻¹ refers to wet mass of sea snail flesh.[#] Below minimum detectable α activity of 0.008 Bq g⁻¹**Table 2.7(b) Comparison of α/β Activities in Sea Snails with Previous Results**

EM&A Report No.	Mean α Activity (Bq g ⁻¹)	SU (Bq g ⁻¹)	Mean β Activity (Bq g ⁻¹)	SU (Bq g ⁻¹)
1 (Baseline)	0.029	0.006	0.064	0.004
2	0.010	0.008	0.034	0.007
3	0.009	0.002	0.032	0.002
4	0.032	0.011	0.050	0.002
5	0.004	0.005	0.045	0.007
6	0.007	0.005	0.042	0.006
7	0.014	0.006	0.063	0.008
8	0.005	0.001	0.040	0.004
9	0.000	0.000	0.023	0.002
10	0.010	0.009	0.045	0.005
11 (2007)	0.000	0.001	0.043	0.002
12 (2008)	0.000	0.000	0.024	0.002
13 (2009)	0.003	0.003	0.035	0.004
14 (2010)	0.005	0.000	0.034	0.002
15 (2011)	0.002	0.001	0.048	0.001
16 (2012)	0.000	0.000	0.014	0.001
17 (2013)	0.001	0.000	0.082	0.024

2.20 One snail sample shows a slight exceedance of Investigation Level.

Airborne Particulates

2.21 The measurement results are given in Table 2.8(a) & 2.8(b).

Table 2.8(a) Net Gross α/β Counts in Airborne Particulate Samples

Location	α Count-rate (cpm)	SD	β Count-rate (cpm)	SD
Blank	0.00 [#]	0.00	0.00 [#]	0.00
A1	0.00 [#]	0.00	4.62	1.03
A2	0.00 [#]	0.00	19.3	1.30
B1	0.00 [#]	0.00	5.85	1.06
B2	0.00 [#]	0.00	13.8	1.30
C1	0.00 [#]	0.00	7.19	1.20
C2	3.86	0.47	42.9	1.93

[#] Below minimum detectable limit of 0.66 cpm for α and 1.6 cpm for β

Table 2.8(b) Comparison of α/β in Airborne Particulate Samples with Previous Results

EM&A Report No.	A		B		C	
	α (cpm)	β (cpm)	α (cpm)	β (cpm)	α (cpm)	β (cpm)
1 (Baseline)	0.00	0.00	0.00	0.00	0.00	1.17
2	0.09	1.38	0.00	0.39	0.00	0.00
3	0.04	0.45	0.00	1.18	0.13	0.86
4	0.12	1.75	0.65	2.18	0.00	0.28
5	0.35	0.94	0.24	0.66	0.07	0.83
6	0.18	0.33	0.00	0.02	0.00	0.00
7	0.16	0.75	0.16	0.09	0.20	0.00
8	0.84	4.87	0.24	1.64	0.09	1.84
9	0.91	3.03	0.29	1.36	0.09	0.42
10	0.32	1.97	0.11	1.05	0.00	0.03
11 (2007)	0.35	1.00	0.11	0.04	0.02	1.75
12 (2008)	0.26	6.18	0.42	5.51	0.19	2.83
13 (2009)	0.97	3.62	0.25	3.09	0.79	1.78
14 (2010)	0.93	16.3	0.00	6.27	0.00	3.55
15 (2011)	--	--	--	--	--	--
16 (2012)	--	--	--	--	--	--
17 (2013)	0.00	12.0	0.00	9.83	1.93	25.1

2.22 A small amount of α and β were detected, particularly in sample C, indicating that sample C is possibly contaminated.

3. REPORT ON ELEVATED ENVIRONMENTAL RADIATION BACKGROUND

- 3.1 The Investigation Levels for environmental samples have been established and they are given in Appendix 1. The relevant action plan is given in the First EM&A Report (Operation Phase).
- 3.2 One of the two snail samples shows a slight exceedance of Investigation Level, which was not observed previously.
- 3.3 All the air particulate samples show rather high β activities when compared to measurements in previous years. Investigations are recommended.
- 3.4 All other samples are normal.

4. REPORT ON NON-COMPLIANCE

- 4.1 The Action Level and Limit Level (A/L Levels) for non-compliance have been established and they are given in Appendix 1 for easy reference. The relevant Event and Action Plan have been developed. Please refer to the First EM&A Report (Operation Phase) for details.

Dose for Radiation Workers

- 4.2 There was no record of exceeding the A/L Levels as recorded by TLDs.

Dose Rates at Un-controlled Areas

- 4.3 No exceedance of the A/L Levels was observed.

Liquid Effluent Discharge

- 4.4 There was no liquid effluent discharged during the monitoring period.

Airborne Effluent Discharge

- 4.5 The average total radon released during the monitoring period was estimated to be 4.08×10^8 Bq/month, which is below the A/L Levels.
- 4.6 The discharged α and β activities were also below the A/L Levels.
- 4.7 The total airborne effluent discharge was below the A/L Levels.

5. RESULT OF ENVIRONMENTAL COMPLIANCE AUDITS

5.1 No compliant was received during the period.

APPENDIX 1

Limit Level and Action Level

The Limit Levels for non-compliance with the Environmental Performance Requirements during the Operation are shown in **Table A1-1**.

Table A1-1 Limit Levels for Non-compliance and Action Levels

Environmental Performance Requirements	Limit Levels	Action Levels (3/10th of Limit Levels)
Dose for radiation workers	1.67 mSv per month	0.5 mSv per month
Dose rate at un-controlled areas	1 µSv per hour	0.3 µSv per hour
Liquid effluent discharge	10 ALI per month	3 ALI per month
Airborne effluent discharge	10 ALI per month	3 ALI per month

Investigation Level

With the help of all the internal monitoring, it is unlikely that the effluents will cause any observable increase in the radiation levels in the vicinity of the Facility under normal operation. It is also not anticipated that any significant quantity of the radioactive wastes would be released to the environment under even the most severe natural disasters. Nevertheless when the environmental samples are found to have radioactivities higher than the normal fluctuation of the established baseline levels, some investigation has to be initiated. The levels that trigger the investigation are called investigation levels and they are given in **Table A1.2**.

Table A1.2 Investigation Levels for Environmental Samples

Environmental Samples		Investigation Levels	
Ambient γ dose rate ($\mu\text{Sv h}^{-1}$)	A	0.23	3 × SD of individual baseline dose rate
	B	0.25	
	D	0.27	
	E	0.29	
	F	0.28	
	G	0.27	
	H	0.31	
	I	0.32	
	J	0.24	
	K	0.32	
	L	0.30	
	M	0.31	
	N	0.29	
	O	0.24	
P	0.29		
Soil (Bq kg^{-1})	²²⁶ Ra	91.7	3 × SU of baseline samples
	²²⁸ Th	128.5	
	⁴⁰ K	1497	

	¹³⁷ Cs	1.36	
	Other γ emitters		Occurrence in any quantities
Sand (Bq kg ⁻¹)	²²⁶ Ra	32.0	3 × SU of baseline samples
	²²⁸ Th	38.1	
	⁴⁰ K	894	
	Other γ emitters		Occurrence in any quantities
Grass (Bq g ⁻¹)	Gross α	0.22	3 × SU of baseline samples
	Gross β	0.43	
	γ emitters not found in baseline		Occurrence in any quantities
Sea water (Bq L ⁻¹)	Gross α	1.52	3 × SU of baseline samples
	Gross β	9.3	
	γ emitters not found in baseline		Occurrence in any quantities
Fish (Bq g ⁻¹)	Gross α	0.021	3 × SU of baseline samples
	Gross β	0.076	
Sea snails (Bq g ⁻¹)	Gross α	0.048	3 × SU of baseline samples
	Gross β	0.076	
Airborne particulates (cpm)	Gross α		Occurrence in any quantities
	Gross β		

- SD is the standard deviation of a single sample.
- SU is standard uncertainty of the sample group.