



**Agreement No. DP 04/2012
Post-Construction Ecological Monitoring
of Drainage Improvement Works in Southern Lantau
Implemented under 4128CD in Contract DC/2006/11**

Monthly EM&A Report – April 2013

May 2013

	Name	Signature
Prepared & Checked:	Chiu Ming Ho (Ecologist)	
Reviewed & Approved:	Sharne McMillan (Environmental Team Leader)	
Version:	2	Date: May 2013


AECOM Asia Co. Ltd.

8/F, Grand Central Plaza, Tower 2, 138 Shatin Rural Committee Road, Shatin, NT, Hong Kong

Tel: (852) 3922 9000 Fax: (852) 3922 9797 www.aecom.com

Pursuant to Condition 4.3 of Environmental Permit No. EP-237/2005/B, this monthly EM&A Report for post-construction ecological monitoring and ecological water monitoring during April 2013 has been certified by the Environmental Team Leader (ETL) and verified by the Independent Environmental Checker (IEC)

Certified by:

Signature:  _____

Date: 16/05/2013

Ms. Sharne McMillan
Environmental Team Leader (ETL)
AECOM Asia Co. Ltd

Verified by:

Signature:  _____

Date: 16/05/2013

Mr. Roger Leung
Independent Environmental Checker (IEC)
ENVIRON Hong Kong Limited

Table of Contents

	Page
EXECUTIVE SUMMARY	1
1. INTRODUCTION.....	2
1.1. Background	2
1.2. Project Description	2
1.3. Report Objectives.....	2
2. ECOLOGICAL MONITORING PARAMETERS	2
2.1. Ecological Surveys.....	2
2.2. Ecological Water Quality Monitoring	5
2.3. Limitations	6
3. MONITORING RESULTS	8
3.1. Ecological Survey Findings	8
3.2. Ecological Water Quality Monitoring (EWQM)	20
4. ECOLOGICAL MONITORING SCHEDULE.....	21
5. DISCUSSION AND RECOMMENDATIONS	21
6. REFERENCES.....	25

List of Figures

- Figure 1 Ecological Monitoring Locations at Pak Ngan Heung River, Luk Tei Tong River, and Luk Tei Tong Bypass Channel and the Reference Site
- Figure 2 Ecological Water Quality Monitoring Locations at Pak Ngan Heung River, Luk Tei Tong River, Luk Tei Tong Bypass Channel and the Reference Site

List of Tables

- Table 2.1 Limit of Reporting for Water Quality Parameters
- Table 3.1 Number of Avifauna Recorded at Pak Ngan Heung River (PNH)
- Table 3.2 Number of Dragonfly Recorded at Pak Ngan Heung River (PNH)
- Table 3.3 Relative Abundance of Aquatic Macroinvertebrate and Fish Recorded at Pak Ngan Heung River (PNH)
- Table 3.4 Number of Avifauna Recorded at Luk Tei Tong River (LTT)
- Table 3.5 Relative Abundance of Aquatic Macroinvertebrate and Fish Recorded at Luk Tei Tong River (LTT)
- Table 3.6 Vegetation Coverage at Luk Tei Tong Bypass Channel (LBC) and Reference Site (RS)
- Table 3.7 Number of Avifauna Recorded at Luk Tei Tong Bypass Channel (LBC)
- Table 3.8 Number of Avifauna Recorded at Reference Site (RS)
- Table 3.9 Relative Abundance of Aquatic Macroinvertebrate and Fish Recorded at Luk Tei Tong Bypass Channel (LBC)
- Table 3.10 Summarized Ecological Water Quality Monitoring Results (April 2013)
- Table 3.11 Baseline Results of Ecological Water Quality Monitoring Results (September 2007)
- Table 5.1 Key Observations/Comments and Recommendations Arising from the April 2013 Monitoring Period

List of Appendices

Appendix 1	Calibration Certificate of the Instruments (pH Meter and Multi-meter)
Appendix 2a	Plant Species Recorded in Pak Ngan Heung River and Luk Tei Tong River
Appendix 2b	Plant Species Recorded in Luk Tei Tong Bypass Channel and the Reference Site
Appendix 3	Ecological Water Quality Monitoring – Raw Data

EXECUTIVE SUMMARY

This is the fourth bi-monthly post-construction ecological monitoring and audit exercise for “Drainage Improvement in Southern Lantau” conducted by AECOM. This report concludes the post-construction phase ecological monitoring and audit requirement for the activities undertaken during the period of 1 April 2013 to 30 April 2013.

Ecological monitoring and ecological water quality monitoring were performed on 11 April 2013 and 23 April 2013, respectively. Results obtained are presented in this report.

The Environmental Team (ET) will continue to implement the environmental monitoring & audit (EM&A) programme in accordance with the EM&A Manual and Environmental Permit requirement. The report is available for public inspection and will be uploaded to the dedicated project website (<http://www.envproject.com/sldiwema.htm>).

1. INTRODUCTION

1.1. Background

1.1.1. The Drainage Services Department (DSD) has implemented Contract No. DC/2006/11 “Drainage Improvement in Southern Lantau and Construction of Mui Wo Village Sewerage Phase 1”. The monitoring requirements of the drainage improvement works are subject to the conditions specified in Environmental Permit (EP) No. EP-237/2005/B issued by the Environmental Protection Department (25 January 2006). In compliance with the EP, an Environmental Monitoring and Audit (EM&A) programme was established during the construction and post-construction phases of the project. The operation of the project is subject to the conditions in EP No. EP-434/2012.

1.1.2. The Post-Construction Ecological Monitoring and Audit of Drainage Improvement Works in Southern Lantau under Agreement No. DP 04/2012, commenced in January 2012. AECOM Asia Co. Ltd. was appointed by DSD as the Environmental Team (ET) to conduct the above captioned monitoring project from October 2012 onwards. This is the fourth bi-monthly post-construction ecological monitoring and audit report under that appointment.

1.2. Project Description

1.2.1. Under Contract No. DC/2006/11, the improvement works were undertaken at Pak Ngan Heung River (PNH), Luk Tei Tong River (LTT) and Tai Tei Tong River (TTT) in Southern Lantau, west of Mui Wo. The works for which the post-construction ecological monitoring required by EP No. EP-237/2005/B covered included:

- the drainage channel and a three-cell box culvert at PNH;
- the drainage channel at LTT; and
- the bypass channel at LTT.

1.2.2. No ecological monitoring and ecological water monitoring was required following the drainage improvement works at TTT and village sewerage works in Mui Wo.

1.2.3. Both PNH and LTT are part of the Mui Wo River (also named as Silver River) in Lantau Island. These two tributaries of Mui Wo River, together with Tai Tei Tong River, then joined and connected to Silver Mine Bay next to Mui Wo.

1.3. Report Objectives


This report presents the findings of the ecological monitoring and the ecological water monitoring conducted in April 2013.

2. ECOLOGICAL MONITORING PARAMETERS

2.1. Ecological Surveys

2.1.1. Details of the monitoring parameters and survey methodology are described below. According to the Final EM&A Manual, a specific ecological monitoring programme of the improved section of PNH, LTT, LTT Bypass Channel (LBC) and its Reference Site (RS) is recommended.

Figure 1 Ecological Monitoring Locations at Pak Ngan Heung River, Luk Tei Tong River, Luk Tei Tong Bypass Channel and the Reference Site



Pak Ngan Heung River and Luk Tei Tong River

- 2.1.2. The ecological survey for these two rivers was divided into nine 50 m sections and comprised the following:
- Two sections for downstream of PNH (PNH1 and 2), two sections for upstream of PNH (PNH3 and 4);
 - Five sections for LTT (LTT1 to 5).
- 2.1.3. The location plan is shown in **Figure 1** for reference.
- 2.1.4. The monitoring parameters and survey methodology for each section are described below:
- (a) Bird species in each 50 m section were surveyed quantitatively using transect count method. Birds within the river channel and on the riverbank were identified to species and their abundance was recorded. Birds that flew over/across the river channel without landing were not considered to be utilising the area and thus excluded from the records. This does not apply to species which rarely land and is known to associate with the habtiat such as Barn Swallow.
 - (b) Surveys on aquatic macroinvertebrate focused on determination of the diversity and abundance. Sampling methods included active searching, direct observation, hand netting and kick sampling. In each section, the macroinvertebrate species composition was identified and their relative abundance was recorded.
 - (c) Surveys on fish focused on determination of the diversity and abundance of fish communities. Sampling methods included active searching, direct observation, and hand netting, and were determined in accordance with site conditions. In each section, the fish species composition was identified and their relative abundance was recorded.
 - (d) Adult odonate community in each 50 m section were surveyed quantitatively by transect count method. Adult odonate within the river channel and on the riverbank were identified to species and their abundance was recorded. Species requiring close examination were netted.
 - (e) Aquatic, emergent and riparian vegetation community was recorded by walk-through survey. Plant species composition and their relative abundance were recorded.

Luk Tei Tong Bypass Channel

- 2.1.5. The ecological survey for the Luk Tei Tong Bypass Channel (LBC) and its Reference Sites (RS) were carried out in every 50 m section and comprised the following:
- Five sections for LBC (LBC1 to 5);
 - Five sections for RS (RS1 to 5).
- 2.1.6. The location plan is shown in **Figure 1** for reference.
- 2.1.7. The monitoring parameters and survey methodology are described below:
- (a) Bird species in each 50 m section were surveyed quantitatively using transect count method. Birds within the river channel and on the riverbank were identified to species and their abundance was recorded. Birds that flew over/across the river channel without landing were not considered to be utilising the area and thus excluded from the records. This does not apply to species which rarely land and is known to associate with the habtiat such as Barn Swallow.
 - (b) Where/when water was present, surveys of aquatic macroinvertebrate focused on determination of their diversity and abundance of stream aquatic communities. Sampling

methods included active searching, direct observation, hand netting and kick sampling. In each section, macroinvertebrate species composition was identified and their relative abundance was recorded.

- (c) Where/when water was present, surveys of fish focused on determination of their diversity and abundance. Sampling methods included active searching, direct observation, and hand netting, were determined in accordance with site conditions. In each section, fish species composition was identified and their relative abundance was recorded.
 - (d) Adult odonate community in each 50 m section were surveyed quantitatively by transect count method. Adult dragonflies within the river channel and on the riverbank were identified to species and their abundance was recorded. Species requiring close examination were netted.
 - (e) Line-intercept method was adopted to determine the relative plant cover of aquatic, emergent and riparian vegetation. One line transect of 10 m was set perpendicular to the stream channel at each section, and five 1 m x 1 m quadrats were placed along the transect. Relative coverage and plant species intercepting the transect line was recorded. Percentage cover of each species within the quadrat was recorded to the nearest 10% (except "1" = present but insignificant cover, normally 1 to 2 individuals, and 5% = up to 5%). The conditions of vegetation were described.
 - (f) Herpetofauna community within LBC and RS were surveyed by active searching in potential habitats. Reptiles were identified and their abundance was recorded. Amphibians were identified by their calls and the number of calling males in each section was recorded.
- 2.1.8. For all surveys, identification of plant species and distribution status in Hong Kong were made with reference to Corlett *et al.* (2000), Hu *et al.* (2003), Hong Kong Herbarium (2012), and Hong Kong Herbarium and South China Botanical Gardens (2007; 2008; 2009; 2011).
- 2.1.9. In terms of assessing geographical distribution, published references specializing in the distribution of specific faunal groups in Hong Kong have been utilized. For general status, these have included Fellowes *et al.* (2002) and the Hong Kong Biodiversity Database (AFCD, 2013), and for specific faunal groups, these have included: Avifauna – Carey *et al.* (2001), Viney *et al.* (2006); Dragonflies – Tam *et al.* (2011); Butterflies – Lo (2005); and Chan *et al.* (2011); Amphibians – Chan *et al.* (2005); Reptiles – Chan *et al.* (2006), Chan *et al.* (2009), and Karsen *et al.* (1998); Terrestrial Mammals – Shek (2006); Freshwater Fish – Lee *et al.* (2004); and Freshwater Community – Dudgeon (2003). The status and rarity of vascular plants has been based on Hu *et al.* (2003) and Corlett *et al.* (2000).

2.2. Ecological Water Quality Monitoring

- 2.2.1. Ecological water quality monitoring along PNH, LTT, LBC, and RS was carried out. Ten locations were selected and comprised the following:
- Three locations for existing PNH (WE1 to 3);
 - Three locations for existing LTT (WE4 to 6);
 - Two locations for RS (WE7 to 8);
 - Two locations for existing LBC (WE9 to 10).
- 2.2.2. The location plan for ecological water quality monitoring is shown in **Figure 2** for reference.
- 2.2.3. Water Quality Monitoring along PNH, LTT, LBC and RS included the monitoring parameters shown below:

- Biochemical Oxygen Demand (BOD₅)
- Nitrate
- Ammonia
- Reactive Phosphorus
- Total Suspended Solids (SS)
- Temperature
- Dissolved Oxygen (DO)
- Water Depth* and Water Flow Rate
- Conductivity
- pH
- Salinity
- Sediment Characteristics

Note:

*As referred to in the Final EM&A Manual, Water Depth is required only for LBC.

2.2.4. The DO, water depth and water flow rate, conductivity, pH, temperature, salinity and sediment characteristics were measured in-situ while the other water samples were analyzed in a HOKLAS accredited laboratory and the analyses followed the standard methods according to APHA Standard Methods for the Examination of Water and Wastewater, 20th Edition, or equivalent. The limit of reporting for the laboratory analysis is summarized in **Table 2.1**.

Table 2.1 Limit of Reporting for Water Quality Parameters

Parameters	Limit of Reporting (mg/L)
Total Suspended Solids	2
Biochemical Oxygen Demand (BOD ₅)	2
Nitrate	0.01
Ammonia	0.01
Reactive Phosphorus	0.01

2.2.5. The instrument for in-situ measurement of temperature, DO, salinity and conductivity is a portable and weather proof multi-meter complete with cable and uses a DC power source (YSI 85), whereas Orion 230A+ is used as for pH measurement. Calibration certificates are attached in **Appendix 1**. The instruments are capable of measuring:

- pH in the range of 0 to 14
- Temperature of -5 to +65°C
- DO in the range of 0 to 20 mg/L and 0 to 200% saturation
- Salinity in the range of 0-80ppt
- Conductivity in the range of 0 to 4999 µS/cm


2.2.6. According to the requirement of the Final EM&A Manual, two consecutive measurements for parameters of DO concentration, and DO saturation are required to be taken at each monitoring location. When the difference in value between the first and second reading of DO is more than 25%, the reading was discarded and a further reading taken.

2.3. Limitations

2.3.1. No water was present at LBC2 to LBC5 at the time of survey, therefore aquatic fauna surveys were not undertaken in these locations.

2.3.2. No water was present at WE7 - WE10 at the time of survey, therefore water quality monitoring was not undertaken at these locations.

Figure 2 Ecological Water Quality Monitoring Locations at Pak Ngan Heung River, Luk Tei Tong River, Luk Tei Tong Bypass Channel and the Reference Site



3. MONITORING RESULTS

3.1. Ecological Survey Findings

Pak Ngan Heung River (PNH)

- 3.1.1. The lower stream of PNH (PNH1 and PNH2) is subject to tidal influence from Silver Mine Bay. Vertical concrete retaining wall formed the banks of the river channel. The two sections were located at the mouth of the PNH. PNH1 and PNH2 were adjacent to each other. The bridge formed the southern boundary of PNH1 whereas the box-culvert formed the northern boundary of PNH2. Small boulders and sandy substrate formed the main component of the streambed.
- 3.1.2. Rock-filled gabion formed the eastern bank and the gabion and a vertical concrete retaining wall formed the western bank of the upper stream (PNH3 and PNH4). PNH3 and PNH4 are adjacent to each other. PNH4 comprised a man-made cascade, including a fish ladder, while PNH3 comprised a pool below the cascade and was bounded by a bridge at its downstream end. Small boulders and sandy substrate were the main component in the middle streambed which allowed water flow and pool formation, whereas big boulders were scattered on both sides of the streambed and had an absence of water. The width of the fish ladder at PNH4 is approximately 7 m.
- 3.1.3. The cascade/fish ladder at PNH4 was open and free of vegetation, exposing the feature and allowing free water flow and pool formation.

Vegetation

- 3.1.4. At PNH3 and PNH4, a total of 16 plant species were recorded. The vegetation was dominated by exotic species such as Mile-a-minute (*Mikania micrantha*) and *Bidens alba* at both PNH3 and PNH4. The vegetation predominantly grew on the banks of PNH3 pool and outer edges of the PNH4 cascade. Species such as *Polygonum* sp. and *Commelina* sp. were scattered within the flowing water of the PNH4 cascade.
- 3.1.5. At PNH1 and PNH2, no plant species were recorded within the river channel. The vegetation recorded on the vertical wall has not changed significantly since the last monitoring period, which includes a record of *Bidens alba* and seedlings of Opposite-leaved Fig (*Ficus hispida*).
- 3.1.6. The list of plant species is presented in **Appendix 2a**.

Terrestrial Fauna

- 3.1.7. Eleven avifauna species were recorded at PNH, all of which are common in Hong Kong (**Table 3.1**). Six avifauna species were recorded at lower PNH (PNH1 and PNH2). Little Egret (*Egretta garzetta*) was the only species of conservation importance recorded, which occurred at PNH2. One individual was recorded moving at PNH2. During the monitoring, the water level at lower PNH was approximately 30 cm.
- 3.1.8. Eight avifauna species were recorded at PNH3 whilst no avifauna species were recorded at PNH4. The birds at upper PNH3 were mostly observed along the banks of the river channel, including the wetland species: Common Sandpiper (*Actitis hypoleucos*) and White-breasted Waterhen (*Amaurornis phoenicurus*). Other recorded species were generalist such as Japanese White-eye (*Zosterops japonicus*) and Crested Myna (*Acridotheres cristatellus*). All are of them are common and abundant in Hong Kong (AFCD, 2013).
- 3.1.9. One individual of dragonfly, Yellow Featherlegs (*Copera marginipes*), was recorded at PNH4 (**Table 3.2**). This species is abundant in Hong Kong (AFCD, 2013).
- 3.1.10. No herpetofauna was recorded at PNH.

Aquatic Macroinvertebrate and Fish

- 3.1.11. One fish species, one crab species and six species of other aquatic invertebrates were recorded at PNH (PNH1 to PNH4) (**Table 3.3**). A low abundance and species richness of common species were recorded at PNH.

- 3.1.12. At lower PNH (PNH1 and PNH2), one fish, one crab and six other aquatic invertebrate species (such as the snail, *Nerita albicilla*; Lymnaeidae; and Amphipoda) were recorded. Goldfish (*Carassius auratus*) was found foraging in the waterbody at PNH2 and PNH4.
- 3.1.13. At upper PNH (PNH3 and PNH4), one fish species and one other aquatic macroinvertebrate species were recorded. Goldfish was also recorded foraging at PNH4. The insect, *Baetidae*, was found under boulders in the waterbody at both PNH3 and PNH4.

Table 3.1 Number of Avifauna Recorded at Pak Ngan Heung River (PNH)

Common Name ⁽¹⁾	Scientific Name	Distribution in Hong Kong ⁽²⁾	Principal Status ⁽³⁾	Level of Concern ⁽⁴⁾	Protection Status in China ⁽⁵⁾	China Red Data Book ⁽⁶⁾	IUCN Red List ⁽⁷⁾	PNH1	PNH2	PNH3	PNH4
Little Egret ⁽⁸⁾	<i>Egretta garzetta</i>	Common	P	PRC (RC)	-	-	-		1		
White-breasted Waterhen ⁽⁸⁾	<i>Amaurornis phoenicurus</i>	Common	R	-	-	-	-			1	
Common Sandpiper ⁽⁸⁾	<i>Actitis hypoleucos</i>	Common	M,W	-	-	-	-			1	
Common Kingfisher ⁽⁸⁾	<i>Alcedo atthis</i>	Common	AM,P	-	-	-	-	1			
Red-whiskered Bulbul	<i>Pycnonotus jocosus</i>	Abundant	R	-	-	-	-	2		1	
Barn Swallow	<i>Hirundo rustica</i>	Abundant	SpM, Su	-	-	-	-	1		3	
Yellow-bellied Prinia	<i>Prinia flaviventris</i>	Common	R	-	-	-	-			4	
Japanese White-eye	<i>Zosterops japonicus</i>	Abundant	R,?W	-	-	-	-	5		1	
Crested Myna	<i>Acridotheres cristatellus</i>	Common	R	-	-	-	-			1	
Oriental Magpie Robin	<i>Copsychus saularis</i>	Abundant	R	-	-	-	-			1	
White Wagtail	<i>Motacilla alba</i>	Common	W,R	-	-	-	-	1			

Note:

(1) All wild birds are protected under Wild Animal Protection Ordinance (Cap. 170).

(2) AFCD (2013) Hong Kong Biodiversity Database.

(3) R=resident; Sp=spring; Su=summer; A=autumn; W=winter visitor; M=migrant; P=present all year, exact composition unknown; ?W=the extent of immigration in winter is unclear.

(4) Fellowes *et al.* (2002);RC=Regional Concern; PRC=Potential Regional Concern. Letters in parentheses indicate that the assessment is on the basis of restrictedness in breeding and/or roosting sites rather than in general occurrence.

(5) List of Wild Animals Under State Protection (promulgated by State Forestry Administration and Ministry of Agriculture on 14 January, 1989).

(6) Zheng and Wang (1998).

(7) IUCN (2012). IUCN Red List of Threatened Species. Version 2012.2.

(8) Wetland-dependent species (including wetland-dependent species and waterbirds).

Table 3.2 Number of Dragonfly Recorded at Pak Ngan Heung River (PNH)

Common Name	Scientific Name	Distribution in Hong Kong ⁽¹⁾	Level of Concern ⁽²⁾	Protection Status in China ⁽³⁾	China Red Data Book ⁽⁴⁾	IUCN Red List ⁽⁵⁾	PNH 1	PNH 2	PNH 3	PNH 4
Yellow Featherlegs	<i>Coperla marginipes</i>	Abundant	-	-	-	-				1

Note:

- (1) AFCD (2013). Hong Kong Biodiversity Database.
- (2) Fellowes *et al.* (2002).
- (3) List of Wild Animals Under State Protection (promulgated by State Forestry Administration and Ministry of Agriculture on 14 January, 1989).
- (4) Zheng and Wang (1998).
- (5) IUCN (2012). IUCN Red List of Threatened Species. Version 2012.2.

Table 3.3 Relative Abundance of Aquatic Macroinvertebrate and Fish Recorded at Pak Ngan Heung River (PNH)

Fauna Group	Scientific Name	Common Name	Distribution in Hong Kong ⁽¹⁾	Level of Concern ⁽²⁾	Protection Status in China ⁽³⁾	China Red Data Book ⁽⁴⁾	IUCN Red List ⁽⁵⁾	PNH1	PNH2	PNH3	PNH4
Fish	<i>Carassius auratus</i>	Goldfish	-	-	-	-	-		+		+
Crab	-	Unidentified crab species	-	-	-	-	-	+			
Polychaete	<i>Polychaete</i>	-	-	-	-	-	-	+			
Oligochaeta	<i>Oligochaeta</i>	-	-	-	-	-	-	+			
Snail (Nerites)	<i>Nerita albicilla</i>	-	-	-	-	-	-	+++	+++		
Snail (Lymnaeidae)	Lymnaeidae	-	-	-	-	-	-	++	++		
Snail (Amphipoda)	Amphipoda	-	-	-	-	-	-	++	++		
Insect	<i>Baetidae</i>	-	-	-	-	-	-	+	++	+	++

Note:

- (1) AFCD (2013). Hong Kong Biodiversity Database.
- (2) Fellowes *et al.* (2002).
- (3) List of Wild Animals Under State Protection (promulgated by State Forestry Administration and Ministry of Agriculture on 14 January, 1989).
- (4) Zheng and Wang (1998).
- (5) IUCN (2012). IUCN Red List of Threatened Species. Version 2012.2
- (6) Relative abundance: + = occasional, less than 5 individuals were found; ++ = common, 5-20 individuals were found; +++ = abundant, more than 20 individuals were found.

Luk Tei Tong River (LTT)

- 3.1.14. The LTT is subject to tidal influence from Silver Mine Bay and is estuarine in nature. It is a north-south running river. A vertical concrete retaining wall formed the river bank of the LTT1 whereas rock-filled gabion formed the river bank of LTT2 to LTT5. LTT1 was located at the confluence with Pak Ngan Heung River, Tai Tei Tong River and Luk Tei Tong River. Since it is subject to tidal flow, water flowed from south to north during the survey when the tide was going out. LTT1 and LTT2 had sandy substrate whilst LTT3 to LTT5 had muddy substrate. Clusters of boulders occurred at both sides of the river channel. The width of the river channel was approximately 8-10 m.
- 3.1.15. No evidence of maintenance works (including those relevant to Conditions 2.1 to 2.4 of EP No. EP-434/2012) was observed during the monitoring period.

Vegetation

- 3.1.16. A total of 10 plant species were recorded in LTT. More than half of the recorded species were exotic. The majority were herbs or climbers scattered along the gabion such as Smooth Crotalaria (*Crotalaria pallid*) and Guinea Grass (*Panicum maximum*). In addition to the mangrove species (Spiny Bears Breech, *Acanthus ilicifolius*; and *Kandelia obovata*) that colonized inside the river channel at LTT2 and LTT3, several seedlings of *Kandelia obovata* have naturally regenerated in LTT1, and LTT5.
- 3.1.17. The list of plant species is presented in **Appendix 2a**.

Terrestrial Fauna

- 3.1.18. A total of ten avifauna species were recorded at LTT, all of them are common or abundant in Hong Kong (AFCD, 2013) (**Table 3.4**). Waterbirds species, including Grey Heron (*Ardea cinerea*), Little Egret and Common Sandpiper (*Actis hypoleucos*), and other lowland species such as White Wagtail (*Motacilla alba*) were recorded feeding in the main river channel. Grey Heron and Little Egret are common species of conservation importance. Individuals of Barn Swallow (*Hirundo rustica*) were observed flying over the river channel across LTT.
- 3.1.19. No dragonflies or herpetofauna were recorded at the LTT during the monitoring.

Aquatic Macroinvertebrate and Fish

- 3.1.20. A total of three fish species, two crab species and twelve species of other aquatic invertebrates were recorded from the LTT (**Table 3.5**). All fish species recorded mainly occur in river mouth or estuarine environments in Hong Kong (AFCD, 2013). Grey Mullet (*Mugil cephalus*) were recorded through LTT2 to LTT3 while White-spotted Rabbit Fish (*Siganus canaliculatus*) was recorded at LTT2. A Goby sp. was recorded at LTT5.

Table 3.4 Number of Avifauna Recorded at Luk Tei Tong River (LTT)

Common Name ⁽¹⁾	Scientific Name	Distribution in Hong Kong ⁽²⁾	Principal Status ⁽³⁾	Level of Concern ⁽⁴⁾	Protection Status in China ⁽⁵⁾	China Red Data Book ⁽⁶⁾	IUCN Red List ⁽⁷⁾	LTT1	LTT2	LTT3	LTT4	LTT5
Grey Heron ⁽⁸⁾	<i>Ardea cinerea</i>	Common	W	PRC	-	-	-	2				
Little Egret ⁽⁸⁾	<i>Egretta garzetta</i>	Common	P	PRC (RC)	-	-	-	1	1			
Common Sandpiper ⁽⁸⁾	<i>Actitis hypoleucos</i>	Common	M, W	-	-	-	-	1				
Spotted Dove	<i>Streptopelia chinensis</i>	Abundant	R	-	-	-	-		2	1		
Barn Swallow	<i>Hirundo rustica</i>	Abundant	Sp, M, Su	-	-	-	-	1	1	2	1	1
Dusky Warbler	<i>Phylloscopus fuscatus</i>	Common	W	-	-	-	-					1
Plain Prinia	<i>Prinia inornata</i>	Common	R	-	-	-	-			1		
Crested Myna	<i>Acridotheres cristatellus</i>	Common	R	-	-	-	-				1	
Oriental Magpie Robin	<i>Copsychus saularis</i>	Abundant	R							1	1	1
White Wagtail	<i>Motacilla alba</i>	Common	W, R								1	

Note:

(1) All wild birds are protected under Wild Animal Protection Ordinance (Cap. 170).

(2) AFCD (2013). Hong Kong Biodiversity Database

(3) R=resident; Sp=spring; Su=summer; W=winter visitor; M=migrant; P=present all year, exact composition unknown.

(4) Fellowes *et al.* (2002); RC=Regional Concern; PRC=Potential Regional Concern. Letters in parentheses indicate that the assessment is on the basis of restrictedness in breeding and/or roosting sites rather than in general occurrence.

(5) List of Wild Animals Under State Protection (promulgated by State Forestry Administration and Ministry of Agriculture on 14 January, 1989).

(6) Zheng and Wang (1998).

(7) IUCN (2012). IUCN Red List of Threatened Species. Version 2012.2.

(8) Wetland-dependent species (including wetland-dependent species and waterbirds).

Table 3.5 Relative Abundance of Aquatic Macroinvertebrate and Fish Recorded at Luk Tei Tong River (LTT)

Fauna Groups	Scientific Name	Common Name	Distribution in Hong Kong ⁽¹⁾⁽²⁾⁽³⁾	Level of Concern ⁽⁴⁾	Protection Status in China ⁽⁵⁾	China Red Data Book ⁽⁶⁾	IUCN Red List ⁽⁷⁾	LTT1	LTT2	LTT3	LTT4	LTT5
Fish	<i>Mugil cephalus</i>	Grey Mullet	Common	-	-	-	-		+	++		
Fish	<i>Siganus canaliculatus</i>	White-spotted Rabbit Fish	Common	-	-	-	-		+			
Fish	Goby sp.	-	-	-	-	-	-					+
Crabs	-	Unidentified crab species	-	-	-	-	-					+
Crabs	<i>Perisesarma bidens</i>	-	-	-	-	-	-		+			
Polychaete	<i>Polychaete</i>	-	-	-	-	-	-		+		+	++
Oligochaeta	<i>Oligochaeta</i>	-	-	-	-	-	-	+				+
Amphipod	Amphipoda	-	-	-	-	-	-	+	++	++	+++	
Tube-worms	<i>Hydroides</i> spp.	-	Very common	-	-	-	-	+				
Snail	<i>Clithon</i> sp.	-	-	-	-	-	-	++	++			
Snail (Nerites)	<i>Nerita albicilla</i>	-	Common	-	-	-	-	++			+	
Snail	<i>Cerithidea cingulata</i>	-	Very common	-	-	-	-			+		
Bivalves	Lymnaeidae	-	-	-	-	-	-			+	++	+
Bivalves	<i>Grafrarium pectinatum</i>	-	Common	-	-	-	-		++			
Bivalves	<i>Saccostrea cucullata</i>	Rock oyster	Very common	-	-	-	-	++				
Barnacles	<i>Balanus amphitrite</i>	-	Very common	-	-	-	-	+	+++			
Insect	<i>Baetidae</i>	-	-	-	-	-	-		+	+		+

Note:

(1) AFCD (2013). Hong Kong Biodiversity Database.

(2) Williams, G (2003). Hong Kong Field Guides – Rocky Shores

(3) Chan *et al.* (2003). Hong Kong Field Guides – Sandy Shores

(4) Fellowes *et al.* (2002).

(5) List of Wild Animals Under State Protection (promulgated by State Forestry Administration and Ministry of Agriculture on 14 January, 1989).

(6) Zheng and Wang (1998).

(7) IUCN (2012). IUCN Red List of Threatened Species. Version 2012.2

(8) Relative abundance: + = occasional, less than 5 individuals were found; ++ = common, 5-20 individuals were found; +++ = abundant, more than 20 individuals were found.

Luk Tei Tong Bypass Channel (LBC) and Reference Site (RS)

- 3.1.21. The LBC is linked to the end of LTT5 and runs east to west but the connection with LTT5 is blocked by a layer of gabion wall approximately 1 m in height which allows water flow between LBC and LTT when water level is higher than the height of the gabion. It is located in the Luk Tei Tong Marsh to the west of the original LTT. Gabion walls formed both sides of the channel bank. Generally, all sections were heavily vegetated except in LBC1 where a small pool of approximately 60 m² in size was located at the western end of LBC1. The pool was separated from the LTT by a weir constructed from a single layer of rock-filled gabion. The substrate comprised soil which was translocated from the marsh area prior to construction of the bypass. The width of the bypass channel was approximately 15 m.
- 3.1.22. The RS was located parallel to the northern side of the LBC. Next to the RS was village housing. The site was vegetated and did not have any free-standing water at the time of survey.

Vegetation

- 3.1.23. A total of 39 plant species were recorded in LBC, of which 19 species were found in the quadrats sampled. The list of plant species is presented in **Appendix 2b**. More than half of the recorded species were exotic. During the survey, only LBC1 included a small patch of open shallow water. Other sections were dry.
- 3.1.24. The habitat at LBC1 differed from the remaining sections in terms of vegetation type. It may be subject to tidal influence during high tide because of its location immediately next to LTT. The sedge, Ferruginous-scale Fimbristylis (*Fimbristylis sieboldii*), dominated LBC1 with a pool of open water forming the western part of the section next to LTT.
- 3.1.25. The plant species recorded in LBC2 to LBC5 were dominated by the exotic species *Wedelia trilobata*. Herbaceous species commonly encountered along the transects were Mile-a-Minute (*Mikania micrantha*) and Ciliate Microstegium (*Microstegium ciliatum*). Other species only formed a small proportion of the vegetation.
- 3.1.26. A total of 41 plant species were recorded in the RS, of which 17 species were found in the quadrats (**Table 3.6**). Seventeen of the 41 species were exotic and common. All sections were dry and were located next to the village housing. The dominant species was exotic *Wedelia trilobata*. Exotic Mile-a-minute and *Aster subulatus* were commonly recorded across the RS sections. The majority of vegetation recorded at the RS could typically be found in disturbed land. Marsh species (e.g. Ginger Lily, *Hedychium coronarium*) was recorded at RS2, RS3 and RS5.
- 3.1.27. The list of plant species is presented in **Appendix 2b**.

Table 3.6 Vegetation Coverage at Luk Tei Tong Bypass Channel (LBC) and Reference Site (RS)

	LBC	RS
No. of species recorded in quadrats	19	17
Total No. of species	39	41
Total No. of exotic species	17	17
Average vegetation coverage	89%	77%
Bare ground coverage	11%	23%

Note:

(1) The transect was not laid along any open water, thus open water coverage was not provided in this table.

Terrestrial Fauna

- 3.1.28. Five species of avifauna were recorded at LBC (**Table 3.7**) whereas eight species of avifauna were recorded at the RS (**Table 3.8**). All recorded species are common or abundant in Hong Kong (AFCD, 2013). Only Greater Coucal (*Centropus sinensis*) is regarded as species of

conservation importance, which is classified as Class II Protected Animal in China and is assessed as vulnerable by China Red Data Book. The species was recorded moving into the vegetation next to the bank at LBC5. All avifauna species are lowland bird species such as Crested Myna (*Acridotheres cristatellus*), Barn Swallow and Oriental Magpie Robin (*Copsychus saularis*) and are generalist adapted to disturbed environment.

3.1.29. No dragonflies or herpetofauna were recorded at LBC and RS during the monitoring.

Aquatic Macroinvertebrate and Fish

3.1.30. Three species of fish were recorded at LBC1 which included Grey Mullet, Jarbua Terapon (*Terapon jarbua*), and White-spotted Rabbit Fish (**Table 3.9**). No species of conservation importance were recorded. All fish species were found foraging in the waterbody.

3.1.31. No aquatic fauna was recorded at the RS or the remaining sections of the LBC as they were dry during the monitoring.

Table 3.7 Number of Avifauna Recorded at Luk Tei Tong Bypass Channel (LBC)

Common Name ⁽¹⁾	Scientific Name	Distribution in Hong Kong ⁽²⁾	Principal Status ⁽³⁾	Level of Concern ⁽⁴⁾	Protection Status in China ⁽⁵⁾	China Red Data Book ⁽⁶⁾	IUCN Red List ⁽⁷⁾	LBC1	LBC2	LBC3	LBC4	LBC5
Greater Coucal	<i>Centropus sinensis</i>	Common	R	-	Class II	Vulnerable	-					1
Barn Swallow	<i>Hirundo rustica</i>	Abundant	SpM, Su	-	-	-	-			5	1	1
Yellow-bellied Prinia	<i>Prinia flaviventris</i>	Common	R	-	-	-	-					1
Crested Myna	<i>Acridotheres cristatellus</i>	Common	R	-	-	-	-			1	3	
Oriental Magpie Robin	<i>Copsychus saularis</i>	Abundant	R	-	-	-	-			2	1	

Note:

- (1) All wild birds are protected under Wild Animal Protection Ordinance (Cap. 170).
- (2) AFCD (2013). Hong Kong Biodiversity Database.
- (3) R=resident; Sp=spring; Su=summer; M=migrant.
- (4) Fellowes *et al.* (2002).
- (5) List of Wild Animals Under State Protection (promulgated by State Forestry Administration and Ministry of Agriculture on 14 January, 1989).
- (6) Zheng and Wang (1998).
- (7) IUCN (2012). IUCN Red List of Threatened Species. Version 2012.2.

Table 3.8 Number of Avifauna Recorded at Reference Site (RS)

Common Name ⁽¹⁾	Scientific Name	Distribution in Hong Kong ⁽²⁾	Principal Status ⁽³⁾	Level of Concern ⁽⁴⁾	Protection Status in China ⁽⁵⁾	China Red Data Book ⁽⁶⁾	IUCN Red List ⁽⁷⁾	RS1	RS2	RS3	RS4	RS5
Spotted Dove	<i>Streptopelia chinensis</i>	Abundant	R	-	-	-	-				2	
Greater Coucal	<i>Centropus sinensis</i>	Common	R	-	Class II	Vulnerable	-					1
Red-whiskered Bulbul	<i>Pycnonotus jocosus</i>	Abundant	R	-	-	-	-				2	
Barn Swallow	<i>Hirundo rustica</i>	Abundant	SpM, Su	-	-	-	-			3		2
Yellow-bellied Prinia	<i>Prinia flaviventris</i>	Common	R	-	-	-	-					1
Masked Laughingthrush	<i>Garrulax perspicillatus</i>	Abundant	R	-	-	-	-					2
Crested Myna	<i>Acridotheres cristatellus</i>	Common	R	-	-	-	-			1	3	
Oriental Magpie Robin	<i>Copsychus saularis</i>	Abundant	R	-	-	-	-			2	1	

Note:

- (1) All wild birds are protected under Wild Animal Protection Ordinance (Cap. 170).
- (2) AFCD (2013). Hong Kong Biodiversity Database.
- (3) R=resident; W=winter visitor.
- (4) Fellowes *et al.* (2002).
- (5) List of Wild Animals Under State Protection (promulgated by State Forestry Administration and Ministry of Agriculture on 14 January, 1989).
- (6) Zheng and Wang (1998).
- (7) IUCN (2012). IUCN Red List of Threatened Species. Version 2012.2.

Table 3.9 Relative Abundance of Aquatic Macroinvertebrate and Fish Recorded at Luk Tei Tong Bypass Channel (LBC)

Fauna Groups	Scientific Name	Common Name	Distribution in Hong Kong ⁽¹⁾	Level of Concern ⁽²⁾	Protection Status in China ⁽³⁾	China Red Data Book ⁽⁴⁾	IUCN Red List ⁽⁵⁾	LBC1	LBC2	LBC3	LBC4	LBC5
Fish	<i>Mugil cephalus</i>	Grey Mullet	Common	-	-	-	-	+				
Fish	<i>Terapon jarbua</i>	Jarbua Terapon	Common	-	-	-	-	++				
Fish	<i>Siganus canaliculatus</i>	White-spotted Rabbit Fish	Common	-	-	-	-	+				

Note:

- (1) AFCD (2013). Hong Kong Biodiversity Database.
- (2) Fellowes *et al.* (2002)
- (3) List of Wild Animals Under State Protection (promulgated by State Forestry Administration and Ministry of Agriculture on 14 January, 1989).
- (4) Zheng and Wang (1998).
- (5) IUCN (2012). IUCN Red List of Threatened Species. Version 2012.2.
- (6) Relative abundance: + = occasional, less than 5 individuals were found; ++ = common, 5-20 individuals were found; +++ = abundant, more than 20 individuals were found.

3.2. Ecological Water Quality Monitoring (EWQM)

- 3.2.1. The post-construction phase EWQM was conducted on 23 April 2013. The monitoring results are presented in **Appendix 3** and summarised in **Table 3.10**, which includes reference to the key Water Quality Objectives (WQOs). Baseline surveys were conducted in 2007 prior to the start of the drainage improvement works. The baseline survey results are presented in **Table 3.11**.
- 3.2.2. Comparison of the data in **Table 3.10** demonstrates that the April 2013 monitoring results meet the key WQOs for the Mui Wo River (EPD, 2011), indicating a reasonable water quality of the subject watercourses. The exception to this is a slight decrease in the pH at WE1 and WE6 which were slightly lower than the lower limit of WQOs. WE1 is the water sampling station at PNH4, whereas WE6 is the water sampling station at the original LTT river near the end of LBC. This phenomenon will be monitored during the future monitoring periods.
- 3.2.3. The water quality monitoring results are discussed in **Section 5**.

Table 3.10 Summarized Ecological Water Quality Monitoring Results (April 2013)

Parameters	Key Water Quality Objectives ⁽¹⁾	WE1	WE2	WE3	WE4	WE5	WE6
Suspended Solids (mg/L)	<20	3.0	3.0	3.0	3.0	3.0	4.0
Nitrogen (Ammonia) (mg/L)	-	0.02	0.18	0.16	0.57	1.95	0.04
Nitrogen (Nitrate) (mg/L)	-	0.17	0.26	0.25	0.56	0.35	0.40
Reactive Phosphorous (mg/L)	-	0.03	0.04	0.05	0.06	0.11	0.01
5-day Biochemical Oxygen Demand (BOD ₅) (mg/L)	<5	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
Dissolved Oxygen (mg/L)	>4	7.52	7.55	7.43	6.83	5.54	7.60
Temperature (°C)	-	20.50	20.80	20.80	22.30	21.80	20.90
pH	6.5 – 8.5	6.29	6.83	6.85	7.40	7.03	6.35
Salinity (ppt)	-	0.02	0.04	0.03	10.3	2.47	0.03
Conductivity (µs/cm)	-	66.2	122.3	124.9	17,411	46.6	75.4
Water Flow (m/s)	-	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1
Water Depth (cm)	-	53	12	18	22	18	33

Note:

(1) The available key Water Quality Objectives (WQOs) for River Monitoring Stations at Mui Wo River on Lantau Island (EPD, 2011).

Table 3.11 Baseline Results of Ecological Water Quality Monitoring Results (September 2007)

Parameters	WE1	WE2	WE3	WE4	WE5	WE6
Suspended Solids (mg/L)	1.0	2.0	3.0	3.0	<1.0	<1.0
Nitrogen (Ammonia) (mg/L)	0.07	0.12	0.11	0.23	0.03	0.02
Nitrogen (Nitrate) (mg/L)	0.12	0.13	0.13	0.31	0.04	0.05
Reactive Phosphorous (mg/L)	0.04	0.06	0.06	0.09	0.06	0.05
5-day Biochemical Oxygen Demand (BOD ₅) (mg/L)	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
Dissolved Oxygen (mg/L)	6.58	6.82	6.37	7.61	6.87	5.70
pH	6.4	7.1	7.0	6.8	6.6	6.1
Salinity (ppt)	<0.1	0.1	0.3	7.6	0.1	<0.1

4. ECOLOGICAL MONITORING SCHEDULE

- 4.1. The next ecological surveys monitoring and ecological water quality is tentatively scheduled for mid-June 2013.

5. DISCUSSION AND RECOMMENDATIONS

- 5.1. The aim of the monitoring programme is to provide data on the re-establishment of aquatic/riparian communities in the PNH and LTT, and allow an assessment of the relative success of the mitigation measures to be made. In addition, monitoring of the LBC will assess whether the proposed channel design has provided suitable compensation for the impacts to the Luk Tei Tong Marsh.
- 5.2. Key observations made during the April 2013 monitoring period in relation to the implemented mitigation measures are presented in **Table 5.1**. Where applicable, recommendations for improving the functionality of the mitigation measures have been made.
- 5.3. Re-establishment of cleared vegetation was observed in PNH and LBC. No further recommendations were made on vegetation re-establishment at PNH since it was not overgrown and blocking water flow; however, this will be monitored.
- 5.4. The cleared bypass channel is re-establishing with exotic species (mainly *Wedelia trilobata*). Whilst a low abundance marsh species were also observed, the re-establishment of weedy species may outcompete the preferred marsh species. One of the potential underlying issues may lie in the fact that water retention in LBC is not sufficient to support growth of marsh species during dry season. However, more observations have to be made and more data has to be collected during wet season.
- 5.5. Seedlings of the mangrove species (*Kandalia obovata*) were seen at other sections of LTT in addition to the existing mangrove stand at the junction at LTT2 and LTT3. Several mangrove seedlings were observed at LTT1, which is the confluence of PNH, TTT and LTT, as well as LTT5, which may imply natural re-colonization of mangrove. The natural regeneration of the mangroves will be monitored and re-planting recommended later, if required.
- 5.6. Whilst some differences between the original 2007 baseline surveys and the April 2013 monitoring surveys are evident, findings from water monitoring could be attributed to a range of factors including seasonal variations, and climatic conditions and/or the influence of tidal status at the time of survey. Taking this into account, the key Water Quality Objectives

(WQOs) for River Monitoring Stations at Mui Wo River (EPD, 2011) have been included to provide a comparison with standard water quality goals applicable to the area (refer to **Table 3.10**).

- 5.7. The Environmental Protection Department (EPD) analyses and presents data from its annual water monitoring programme to express the level of compliance with the statutory WQOs including pH, Suspended Solids (SS), 5-day Biochemical Oxygen Demand (BOD₅), and Dissolved Oxygen (DO). These WQOs specify the long-term water quality goals that the Government is to achieve and maintain for individual rivers in Hong Kong, including the Mui Wo River. As part of the programme five locations are sampled from the Mui Wo River, three of which are associated with the monitoring area for the drainage improvement works (MW1, MW2 and MW4). The objectives related to these sampling locations, have been used in this report. Water quality of the subject watercourses has met the WQOs during the survey with the exception of pH level at WE1 and WE6 slightly lower than the WQO. The cause is not known. The phenomenon will be monitored during the future monitoring periods.
- 5.8. Results of some parameters, such as Ammonia and Nitrate, demonstrated an increase from the baseline survey; however, the reason for this and implications for the re-establishment of the aquatic/riparian communities is not currently known.
- 5.9. Concentration of Nitrate, Ammonia and Reactive Phosphorous was in general lower than the last sampling period, returning to a more reasonable level compared to other sampling periods. Nitrate concentration at WE6, however, demonstrated a 40-fold increase compared to the last sampling period, which was also higher than other sampling periods. Suspended solid concentration demonstrated a slight decrease at all monitoring stations except WE2 and WE6 compared to the last sampling record. This phenomenon will be monitored during the future monitoring periods.
- 5.10. No observable evidence of environmental changes such as odour, or discharge within the surveyed area, were recorded. Frequent precipitation events may contribute to some of the discrepancy of results in the current sampling period because of dilution effect but the precise effect of rainfall is not known.
- 5.11. The April 2013 monitoring period occurs early in the post-construction monitoring programme and provides only a snap-shot of the water quality conditions. Further monitoring is required to draw conclusions regarding the overall success of the mitigation measures implemented into the project. The assessment will be on-going over the course of the monitoring programme and will be presented in subsequent reports as additional information becomes available.

Table 5.1 Key Observations/Comments and Recommendations Arising from the April 2013 Monitoring Period

Location	Mitigation Measure	Observations/Comments	Recommendations
PNH and LTT	Construction of a small fish ladder at the upstream end of the PNH	Vegetation has re-established in PNH4; however, the fish ladder is not currently overgrown and blocking water flow.	<p>The retention of native species, particularly at the edges of the river channel, during any future maintenance activities is recommended, to maintain existing habitat and minimize the re-colonization of exotic species.</p> <p>Some pits have been incorporated into the gabion banks, but do not appear to have been planted up. Planting of riparian vegetation, preferably with native species is suggested.</p> <p>On-going weed</p>

Location	Mitigation Measure	Observations/Comments	Recommendations
			management is recommended, as required, to maintain the open nature of the fish ladder.
		The fish ladder does not meet the lip of the weir at the up-stream end of PNH4 due to a drop of approximately 30 cm. This could limit the overall function of the fish ladder for fish passage/movement up and downstream.	Connection of PNH4 with the upper stream should be maintained in order to assist the movement of the fish.
	Re-establishment of aquatic / riparian communities	Only one fish species (Goldfish, <i>Carassius auratus</i>), which is not of conservation concern, was recorded at PNH4 during this monitoring. The two fish species of conservation importance, Flagtail (<i>Kuhlia marginata</i>) and Predaceous Chub (<i>Parazacco spilurus</i>) were not recorded. Flagtail was recorded in the 2003-2004 EIA surveys, and Predaceous Chub was recorded in the December 2012 survey of post-construction monitoring.	The presence of species of conservation importance in both PNH3 and PNH4 including relative abundance will continue to be monitored.
LBC	Provision of suitable habitat compensation	Vegetation has re-established, which is dominated by exotic plant species (<i>Wedelia trilobata</i>). Only limited marsh species of very low abundance were recorded.	Removal and control of vegetation is recommended to maintain the sub-climatic vegetation community found in marsh habitats. It includes retaining native species (particularly marsh species) within the LBC during any future maintenance activities is recommended, to maintain existing habitat and minimize the re-colonization of exotic species. The regeneration of marsh species in the LBC is to be monitored. To avoid recolonisation of unwanted species (e.g. <i>Wedelia trilobata</i>), replanting of marsh species would be recommended upon confirmation of the water level/availability to support marsh habitat.
		The limited occurrence of typical marsh plant species (although this was also limiting in the RS) suggests that the water levels/availability	On-going monitoring of water levels and species composition within the channel are required.

Location	Mitigation Measure	Observations/Comments	Recommendations
		within the channel may not be adequate to sustain a marsh habitat.	Further assessment should take into account the timing of the surveys (wet/dry season).

6. REFERENCES

AFCD (2013). Hong Kong Biodiversity Database. Available at <http://www.afcd.gov.hk/english/conservation/hkbiodiversity/database/resultlist.asp?lang=en> Accessed on 28 April 2013.

Carey, G. J., Chalmers, M. L., Diskin, D. A., Kennerley, P. R., Leader, P. J., Leven, M. R., Lewthwaite, R. W., Melville, D. S., Turnbull, M., and Young, L. (2001). *The Avifauna of Hong Kong*. Hong Kong Bird Watching Society, Hong Kong.

Chan, A., Cheung, J., Sze, P., Wong, A., Wong, E. and Yau, E. (2011). A Review of the Local Restrictedness of Hong Kong Butterflies. *Hong Kong Biodiversity Newsletter* 21: 1-6. Agriculture, Fisheries and Conservation Department, Hong Kong Special Administrative Region.

Chan, S. K.-F., Cheung, K.-S., Ho, C.-Y., Lam, F.-N., Tang, W.-S., Lau, M. W.-N., Bogadek, A. (2005). *A Field Guide to the Amphibians of Hong Kong*. Agriculture, Fisheries and Conservation Department, Friends of the Country Parks and Cosmos Books Ltd.

Chan, S. K.-F., Cheung, K. S., Ho, C. Y., Lam, F. N., Tang, W. S., Tse, M. L. (2006). *A Field Guide to the Venomous Land Snakes of Hong Kong*. Agriculture, Fisheries and Conservation Department, Friends of the Country Parks and Cosmos Books Ltd.

Chan, S.K.F., Chan, A.S.W., Cheung, K.S., Ho, C.Y., Ng, C.K.Y., Tang, W.S. (2009). *The Skinks of Hong Kong*. *Hong Kong Biodiversity Newsletter: Issue 17*

Corlett, R., Xing, W. F., Ng, C. S., Chau, K. C. L., & Wong, M. Y. L. (2000). *Hong Kong Vascular Plants: Distribution and Status*. *Memoirs of the Hong Kong Natural History Society*, 23, 1-157.

Dudgeon (2003). *Hong Kong Field Guides – Hillstreams*. The Department of Ecology and Biodiversity, The University of Hong Kong.

Environmental Protection Department (2011). *River Water Quality in Hong Kong in 2011*. The Government of the Hong Kong Special Administrative Region.

Fellows, J. R., Lau, M. W., Dudgeon, D., Reels, G. T., Ades, G. W., & Carey, G. J. (2002). *Wild Animals to Watch: Terrestrial and Freshwater Fauna of Conservation Concern in Hong Kong*. *Memoirs of the Hong Kong Natural History Society*, 25, 123-159.

Hong Kong Herbarium and South China Botanical Garden (2007). *Flora of Hong Kong*. Volume 1. Agriculture, Fisheries and Conservation Department, Government of Hong Kong Special Administrative Region.

Hong Kong Herbarium and South China Botanical Garden (2008). *Flora of Hong Kong*. Volume 2. Agriculture, Fisheries and Conservation Department, Government of Hong Kong Special Administrative Region.

Hong Kong Herbarium and South China Botanical Garden (2009). *Flora of Hong Kong*. Volume 3. Agriculture, Fisheries and Conservation Department, Government of Hong Kong Special Administrative Region.

Hong Kong Herbarium and South China Botanical Garden (2011). Flora of Hong Kong. Volume 4. Agriculture, Fisheries and Conservation Department, Government of Hong Kong Special Administrative Region.

Hong Kong Herbarium (2012). Check List of Hong Kong Plants 2012. Agriculture, Fisheries and Conservation Department, HKSAR Government.

Hu, Q.M, Wu, T.L., Xia, N.H., Xing F.W., Patrick C.C.L., Yip, K.W. (2003). Rare and Precious Plants of Hong Kong. Agriculture, Fisheries and Conservation Department, Hong Kong Special Administrative Government.

IUCN (2012). IUCN Red List of Threatened Species. Version 2012.1. Available at www.iucnredlist.org. Accessed on 28 April 2013.

Lee, V. L. F., Lam, S. K. S., Ng, F. K. Y., Chan, T. K. T. and Young, M. L. C. (2004). Field Guide to the Freshwater Fish of Hong Kong. Agriculture, Fisheries and Conservation Department, Friends of the Country Parks and Cosmos Books Ltd. Hong Kong.

Lo, P.Y.F. (2005). Hong Kong Butterflies, 2nd edition. Agriculture, Fisheries and Conservation Department.

Karsen, S. J., Lau, M. W. N. and Bogadek, A. (1998). Hong Kong Amphibians and Reptiles. Urban Council, Hong Kong.

Shek, C.T. (2006). A Field Guide to the Terrestrial Mammals of Hong Kong. Agriculture, Fisheries and Conservation Department, Hong Kong.

Tam, T.W., Leung, K.K., Kwan, B.S.P., Wu, K.K.Y., Tang, S.S.H., So, I.W.Y., Cheng, J.C.Y., Yuen, E.F.M., Tsang, Y.M., and Hui, W.L. (2011). The Hong Kong Dragonflies. AFCD, Friends of Country Park and Cosmos Books Ltd. Hong Kong.

Territory Development Department (1999). Planning and Development Study of Potential Housing Site in Area 54, Tuen Mun: EIA – Final Assessment Report. Prepared by ERM for Territory Development Department, The Government of the Hong Kong Special Administrative Region.

Wan, P. H. (2009). The role of Masked Palm Civet (*Paguma larvata*) and Small Indian Civet (*Viverricula indica*) in seed dispersal in Hong Kong, China. Mphil Thesis, HKU.

William, G.A. (2003). Hong Kong Field Guides: Rocky Shore. The Department of Ecology and Biodiversity, The Hong Kong University of Hong Kong, Hong Kong.

Wong, L. C., Lam, V. W. Y., and Ades, G. W. J. (2009). Ecology of the Birds of Hong Kong. Kadoorie Farm and Botanic Garden, Hong Kong Special Administrative Region.

Wong, Y.H., Li, P.K., Sze, W.C. and Wong, K.C. (2005). Butterfly Garden in the Shing Mun Country Park. In: Hong Kong Biodiversity, Issue 10.

Zheng, G. M. and Wang, Q. S. (1998). China Red Data Book of Endangered Animals: Aves. Science Press, Beijing.

Appendix 1. Calibration certificate of the instruments (pH meter and multi-meter)



ALS Technichem (HK) Pty Ltd

REPORT OF EQUIPMENT PERFORMANCE CHECK/CALIBRATION

CONTACT: MR MIKE SHEK
CLIENT: AECOM ASIA COMPANY LIMITED
ADDRESS: 11/F, TOWER 2, GRAND CENTRAL PLAZA,
138 SHATIN RURAL COMMITTEE ROAD,
SHATIN, N.T.,
HONG KONG.
PROJECT: --

WORK ORDER: HK1301561
LABORATORY: HONG KONG
DATE RECEIVED: 17/01/2013
DATE OF ISSUE: 24/01/2013

COMMENTS

It is certified that the item under calibration/checking has been calibrated/checked by corresponding calibrated equipment in the laboratory.
Maximum Tolerance and calibration frequency stated in the report, **unless** otherwise stated, the internal acceptance criteria of ALS will be followed.

Scope of Test: pH
Description: pH Meter
Brand Name: ORION
Model No.: ORION 230A+
Serial No.: 020365
Equipment No.: W.039.04
Date of Calibration: 24 January, 2013

NOTES


This is the Final Report and supersedes any preliminary report with this batch number.
Results apply to sample(s) as submitted. All pages of this report have been checked and approved for release.

ISSUING LABORATORY: HONG KONG

Address

ALS Technichem (HK) Pty Ltd
11/F Chung Shun Knitting Centre
1-3 Wing Yip Street
Kwai Chung
HONG KONG

Phone: 852-2610 1044
Fax: 852-2610 2021
Email: hongkong@alsglobal.com


Mr. Ewan Kwok Eai Godfrey
Laboratory Manager - Hong Kong

This report may not be reproduced except with prior written approval from ALS Technichem (HK) Pty Ltd.

Page 1 of 2

ADDRESS 11/F, Chung Shun Knitting Centre, 1-3 Wing Yip Street, Kwai Chung, N.T., Hong Kong · PHONE +852 2610 1044 · FAX +852 2610 2021
ALS TECHNICHEM (HK) PTY LTD · An ALS Limited Company

Life Sciences

www.alsglobal.com

RIGHT SOLUTIONS RIGHT PARTNER

REPORT OF EQUIPMENT PERFORMANCE CHECK/CALIBRATION

Work Order: HK1301561
Date of Issue: 24/01/2013
Client: AECOM ASIA COMPANY LIMITED



Description: pH Meter
Brand Name: ORION
Model No.: ORION 230A+
Serial No.: 020365
Equipment No.: W.039.04
Date of Calibration: 24 January, 2013

Date of next Calibration: 24 April, 2013

Parameters:

pH Value

Method Ref: APHA 21st Ed. 4500H:B

Expected Reading (pH Unit)	Displayed Reading (pH Unit)	Tolerance (pH unit)
4.0	3.98	-0.02
7.0	7.02	0.02
10.0	9.93	-0.07
Tolerance Limit (\pm pH unit)		0.20

Temperature

Method Ref: Section 6 of International Accreditation New Zealand Technical Guide No. 3 Second edition March 2008: Working Thermometer Calibration Procedure.

Reading of Ref. thermometer ($^{\circ}$ C)	Displayed Reading ($^{\circ}$ C)	Tolerance ($^{\circ}$ C)
11.0	11.8	0.8
21.0	20.3	-0.7
45.0	45.3	0.3
Tolerance Limit (\pm $^{\circ}$ C)		2.0

Remark: "Displayed Reading" presents the figures shown on item under calibration / checking regardless of equipment precision or significant figures.



ALS Technichem (HK) Pty Ltd

REPORT OF EQUIPMENT PERFORMANCE CHECK/CALIBRATION

CONTACT: MR MIKE SHEK
CLIENT: AECOM ASIA COMPANY LIMITED
ADDRESS: 11/F, TOWER 2, GRAND CENTRAL PLAZA,
138 SHATIN RURAL COMMITTEE ROAD,
SHATIN, N.T.,
HONG KONG.

WORK ORDER: HK1302694
LABORATORY: HONG KONG
DATE RECEIVED: 30/01/2013
DATE OF ISSUE: 06/02/2013

PROJECT: -

COMMENTS

It is certified that the item under calibration/checking has been calibrated/checked by corresponding calibrated equipment in the laboratory.

Maximum Tolerance and calibration frequency stated in the report, unless otherwise stated, the internal acceptance criteria of ALS will be followed.

Scope of Test: Conductivity, Dissolved Oxygen, Salinity and Temperature
Description: YSI PROFESSIONAL PLUS
Brand Name: YSI
Model No.: -
Serial No.: 12M100515
Equipment No.: W.040.01
Date of Calibration: 05 February, 2013

NOTES


This is the Final Report and supersedes any preliminary report with this batch number.
Results apply to sample(s) as submitted. All pages of this report have been checked and approved for release.

ISSUING LABORATORY: HONG KONG

Address

ALS Technichem (HK) Pty Ltd
11/F Chung Shun Knitting Centre
1-3 Wing Yip Street
Kwai Chung
HONG KONG

Phone: 852-2610 1044
Fax: 852-2610 2021
Email: hongkong@alsglobal.com


Mr. Chan Kwok Fai, Godfrey
Laboratory Manager - Hong Kong

This report may not be reproduced except with prior written approval from ALS Technichem (HK) Pty Ltd.

Page 1 of 2

ADDRESS 11/F, Chung Shun Knitting Centre, 1-3 Wing Yip Street, Kwai Chung, N.T., Hong Kong | PHONE +852 2610 1044 | FAX +852 2610 2021
ALS TECHNICHEM (HK) PTY LTD An ALS Limited Company

Life Sciences

www.alsglobal.com

RIGHT SOLUTIONS RIGHT PARTNER

REPORT OF EQUIPMENT PERFORMANCE CHECK/CALIBRATION



Work Order: HK1302694
Date of Issue: 06/02/2013
Client: AECOM ASIA COMPANY LIMITED

Description: YSI PROFESSIONAL PLUS
Brand Name: YSI
Model No.: --
Serial No.: 12M100515
Equipment No.: W.040.01
Date of Calibration: 05 February, 2013 **Date of next Calibration:** 05 May, 2013

Parameters:

Conductivity

Method Ref: APHA (21st edition), 2510B

Expected Reading (uS/cm)	Displayed Reading (uS/cm)	Tolerance (%)
146.9	141.9	-3.4
6667	6522	-2.2
12890	11749	-8.9
58670	55430	-5.5
Tolerance Limit (±%)		10.0

Dissolved Oxygen

Method Ref: APHA (21st edition), 4500O: G

Expected Reading (mg/L)	Displayed Reading (mg/L)	Tolerance (mg/L)
2.20	2.19	-0.01
3.91	3.97	0.06
8.21	8.31	0.10
Tolerance Limit (±mg/L)		0.20

Salinity

Method Ref: APHA (21st edition), 2520B

Expected Reading (NTU)	Displayed Reading (NTU)	Tolerance (%)
0	0	-
10	9.44	-5.6
20	19.13	-4.4
30	27.78	-7.4
Tolerance Limit (±%)		10.0

Temperature

Method Ref: Section 6 of International Accreditation New Zealand Technical Guide No. 3 Second edition March 2006: Working Thermometer Calibration Procedure.

Expected Reading (°C)	Displayed Reading (°C)	Tolerance (°C)
12.0	11.7	-0.3
21.5	21.5	0.0
37.0	36.7	-0.3
Tolerance Limit (±°C)		2.0

Remark: "Displayed Reading" presents the figures shown on item under calibration / checking regardless of equipment precision or significant figures.

Mr. Chan Kwok Fai, Godfrey
 Laboratory Manager - Hong Kong

Appendix 2a: Plant Species Recorded in Pak Ngan Heung River and Luk Tei Tong River

Scientific Name	Growth Form	Native / Exotic to Hong Kong	Distribution in Hong Kong	PNH1	PNH2	PNH3	PNH4	LTT1	LTT2	LTT3	LTT4	LTT5
<i>Acanthus ilicifolius</i>	shrub	native	common						+	+		
<i>Ageratum houstonianum</i>	herb	exotic	common				+					
<i>Bidens alba</i>	herb	exotic	very common	+		+	++			+		
<i>Bidens pilosa</i>	herb	exotic	very common				+			+		
<i>Celtis sinensis</i>	tree	native	common				+					
<i>Colocasia esculenta</i>	herb	native	N/A				+					
<i>Commelina diffusa</i>	herb	native	common			+	++					
<i>Crotalaria pallida</i>	herb	exotic	common							+	+	+
<i>Ficus hispida</i>	tree	native	very common	+								
<i>Ipomoea cairica</i>	climber	exotic	very common									+
<i>Kandelia obovata</i>	shrub or small tree	native	common					+	+	+		+
<i>Ludwigia octovalvis</i>	perennial herb	native	common			+						
<i>Mallotus paniculatus</i>	tree	native	very common				+					
<i>Microstegium ciliatum</i>	perennial procumbent herb	native	very common			++	+					
<i>Mikania micrantha</i>	climber	exotic	very common			++	+++					+
<i>Neyraudia reynaudiana</i>	herb	native	very common							+	+	
<i>Oxalis corniculata</i>	perennial herb	native	very common				+					
<i>Panicum maximum</i>	herb	exotic	very common				+					+
<i>Polygonum chinense</i>	herb	native	very common	+								
<i>Polygonum</i> spp.	herb	N/A	N/A				+					
<i>Praxelis clematidea</i>	perennial herb	exotic	very common	+								
<i>Pueraria</i> spp.	climber	N/A	N/A							+	+	
<i>Rhus succedanea</i>	tree	native	common				+					
<i>Urena lobata</i>	shrub	native	common				+					
<i>Wedelia trilobata</i>	perennial herb	exotic	common				+					

Note:

Code for Abundance: +++=abundant; ++=frequent; +=occasional

Appendix 2b: Plant Species Recorded in Luk Tei Tong Bypass Channel and Reference Site

LLT Bypass Channel (LBC)

Scientific Name	Growth Form	Native / Exotic to Hong Kong	Distribution in Hong Kong	LBC1	LBC2	LBC3	LBC4	LBC5	Average
Species recorded in the quadrats along the transects				Average Percentage Cover					
<i>Apluda mutica</i>	herb	native	very common	0.00	0.00	0.00	0.00	0.00	0.00
<i>Aster subulatus</i>	herb	exotic	N/A	0.06	0.00	0.00	0.00	0.00	0.01
<i>Bidens alba</i>	herb	exotic	very common	0.00	0.00	0.00	0.00	0.00	0.00
<i>Bidens pilosa</i>	herb	exotic	very common	0.00	0.00	0.00	0.00	0.00	0.00
<i>Celosia argentea</i>	herb	native	very common	0.00	0.00	0.00	0.00	0.00	0.00
<i>Colocasia esculenta</i>	herb	native	N/A	0.00	0.00	0.00	0.00	0.00	0.00
<i>Crotalaria pallida</i>	herb	exotic	common	0.00	0.00	0.00	0.02	0.01	0.01
<i>Cyclosorus interruptus</i>	herb	native	common	0.00	0.00	0.03	0.00	0.00	0.01
<i>Fimbristylis sieboldii</i>	herb	native	common	0.81	0.00	0.00	0.00	0.00	0.16
<i>Ipomoea cairica</i>	climber	exotic	very common	0.01	0.00	0.00	0.00	0.00	0.00
<i>Microstegium ciliatum</i>	perennial procumbent herb	native	very common	0.01	0.14	0.03	0.05	0.15	0.08
<i>Mikania micrantha</i>	climber	exotic	very common	0.00	0.00	0.12	0.02	0.05	0.04
<i>Mimosa diplotricha</i>	herb	exotic	rare	0.00	0.00	0.00	0.00	0.01	0.00
<i>Paspalum conjugatum</i>	perennial herb	exotic	common	0.00	0.00	0.00	0.07	0.00	0.01
<i>Praxelis clematidea</i>	perennial herb	exotic	very common	0.00	0.00	0.07	0.00	0.00	0.01
<i>Ruellia coerulea</i>	herb	exotic	N/A	0.12	0.00	0.00	0.00	0.00	0.02
<i>Sapium sebiferum</i>	tree	native	common	0.00	0.00	0.00	0.00	0.00	0.00
<i>Solanum americanum</i>	herb	exotic	very common	0.00	0.00	0.04	0.00	0.01	0.01
<i>Wedelia trilobata</i>	perennial herb	exotic	common	0.00	0.53	0.70	0.74	0.65	0.52
Other species recorded during the walk-through survey				Occurrence of the Species					
<i>Apluda mutica</i>	herb	native	very common	+					
<i>Artemisia japonica Thunb.</i>	herb	native	common				+	+	
<i>Aster subulatus</i>	herb	exotic	N/A	+					
<i>Bidens alba</i>	herb	exotic	very common			+	+	+	
<i>Bidens pilosa</i>	herb	exotic	very common				+	+	
<i>Cassytha filiformis</i>	climber	native	very common					+	
<i>Celosia argentea</i>	herb	native	very common				+	+	
<i>Celtis sinensis</i>	tree	native	common				+		
<i>Colocasia esculenta</i>	herb	native	N/A		+	+	+	+	
<i>Commelina diffusa</i>	herb	native	common			+			
<i>Conyza canadensis</i>	herb	exotic	very common	+					
<i>Crotalaria pallida</i>	herb	exotic	common			+	+	+	
<i>Cyclosorus interruptus</i>	herb	native	common		+	+	+		
<i>Cynodon dactylon</i>	perennial herb	native	very common		+				

Note:

Code: +=occurrence of the species

Appendix 2b: Plant Species Recorded in Luk Tei Tong Bypass Channel and Reference Site

LLT Bypass Channel (LBC)

Scientific Name	Growth Form	Native / Exotic to Hong Kong	Distribution in Hong Kong	LBC1	LBC2	LBC3	LBC4	LBC5
Other species recorded during the walk-through survey				Occurrence of the Species				
<i>Erechtites hieracifolia</i>	herb	exotic	common			+	+	+
<i>Fimbristylis sieboldii</i>	herb	native	common	+				
<i>Hedychium coronarium</i>	shrub	exotic	N/A		+	+	+	
<i>Ipomoea cairica</i>	climber	exotic	very common	+		+	+	+
<i>Ipomoea pes-caprae</i>	perennial herb	native	common					+
<i>Kandelia obovata</i>	shrub or small tree	native	common	+				
<i>Lophatherum gracile</i>	herb	native	common				+	
<i>Ludwigia octovalvis</i>	perennial herb	native	common			+		
<i>Macaranga tanarius</i>	tree	native	common		+			
<i>Microstegium ciliatum</i>	perennial procumbent herb	native	very common	+	+	+	+	+
<i>Mikania micrantha</i>	climber	exotic	very common		+	+	+	+
<i>Mimosa diplotricha</i>	herb	exotic	rare			+	+	+
<i>Nerium oleander</i>	shrub	exotic	common		+			
<i>Neyraudia reynaudiana</i>	herb	native	very common	+				
<i>Panicum maximum</i>	herb	exotic	very common				+	
<i>Paspalum conjugatum</i>	perennial herb	exotic	common				+	
<i>Plantago major</i>	herb	native	very common		+			
<i>Polygonum chinense</i>	herb	native	very common		+		+	
<i>Polygonum spp.</i>	herb	N/A	N/A			+	+	
<i>Praxelis clematidea</i>	perennial herb	exotic	very common		+	+	+	+
<i>Ruellia coerulea</i>	herb	exotic	N/A	+				
<i>Sapium sebiferum</i>	tree	native	common		+	+		
<i>Solanum americanum</i>	herb	exotic	very common		+	+	+	+
<i>Urena lobata</i>	shrub	native	common				+	
<i>Wedelia trilobata</i>	perennial herb	exotic	common		+		+	

Note:

Code: +=occurrence of the species

Appendix 2b: Plant Species Recorded in Luk Tei Tong Bypass Channel and Reference Site

Reference Site (RS)

Scientific Name	Growth Form	Native / Exotic to Hong Kong	Distribution in Hong Kong	RS1	RS2	RS3	RS4	RS5	Average
Species recorded in the quadrats along the transects				Average Percentage Cover					
<i>Aster subulatus</i>	herb	exotic	N/A	0.00	0.00	0.00	0.00	0.04	0.01
<i>Bambusa sp.</i>	bamboo	N/A	common	0.00	0.00	0.00	0.00	0.00	0.00
<i>Bidens alba</i>	herb	exotic	very common	0.00	0.03	0.00	0.03	0.02	0.02
<i>Colocasia esculenta</i>	herb	native	N/A	0.00	0.00	0.00	0.00	0.00	0.00
<i>Conyza canadensis</i>	herb	exotic	very common	0.00	0.00	0.00	0.00	0.00	0.00
<i>Cyclosorus interruptus</i>	herb	native	common	0.00	0.00	0.00	0.00	0.00	0.00
<i>Erechtites hieracifolia</i>	herb	exotic	common	0.00	0.00	0.00	0.00	0.00	0.00
<i>Praxelis clematidea</i>	perennial herb	exotic	very common	0.00	0.00	0.00	0.00	0.00	0.00
<i>Hedychium coronarium</i>	shrub	exotic	N/A	0.00	0.00	0.00	0.00	0.00	0.00
<i>Ipomoea cairica</i>	climber	exotic	very common	0.00	0.00	0.00	0.00	0.02	0.00
<i>Microstegium ciliatum</i>	perennial procumbent herb	native	very common	0.00	0.05	0.01	0.00	0.18	0.05
<i>Mikania micrantha</i>	climber	exotic	very common	0.00	0.00	0.01	0.00	0.25	0.05
<i>Mimosa diplotricha</i>	herb	exotic	rare	0.04	0.02	0.00	0.01	0.00	0.01
<i>Paspalum conjugatum</i>	perennial herb	exotic	common	0.00	0.00	0.00	0.00	0.00	0.00
<i>Pueraria phaseoloides</i>	climber	native	very common	0.00	0.00	0.00	0.00	0.00	0.00
<i>Urena lobata</i>	shrub	native	common	0.00	0.02	0.00	0.00	0.04	0.01
<i>Wedelia trilobata</i>	perennial herb	exotic	common	0.64	0.65	0.82	0.63	0.28	0.60
Other species recorded during the walk-through survey				Occurrence of the Species					
<i>Acacia confusa</i>	tree	exotic	N/A		+		+		
<i>Acrostichum aureum</i>	herb	native	restricted	+					
<i>Amaranthus viridis</i>	herb	native	very common		+				
<i>Aster subulatus</i>	herb	exotic	N/A	+	+	+	+	+	
<i>Bambusa sp.</i>	bamboo	N/A	common	+				+	
<i>Bauhinia blakeana</i>	tree	native	common	+					
<i>Bidens alba</i>	herb	exotic	very common	+	+	+	+	+	
<i>Calliandra haematocephala</i>	shrub	exotic	common					+	
<i>Cassytha filiformis</i>	climber	native	very common					+	
<i>Celosia argentea</i>	herb	native	very common		+		+		
<i>Celtis sinensis</i>	tree	native	common			+			
<i>Colocasia esculenta</i>	herb	native	N/A			+		+	
<i>Conyza canadensis</i>	herb	exotic	very common	+		+	+	+	
<i>Crotalaria pallida</i>	herb	exotic	common				+		
<i>Cyclosorus interruptus</i>	herb	native	common					+	
<i>Cynodon dactylon</i>	perennial herb	native	very common	+					
<i>Emilia sonchifolia</i>	herb	native	very common	+					
<i>Erechtites hieracifolia</i>	herb	exotic	common	+	+				
<i>Ficus hispida</i>	tree	native	very common			+	+		

Note:

Code: +=occurrence of the species

Appendix 2b: Plant Species Recorded in Luk Tei Tong Bypass Channel and Reference Site

Reference Site (RS)

Scientific Name	Growth Form	Native / Exotic to Hong Kong	Distribution in Hong Kong	RS1	RS2	RS3	RS4	RS5
Other species recorded during the walk-through survey				Occurrence of the Species				
<i>Hedychium coronarium</i>	shrub	exotic	N/A		+	+		+
<i>Ipomoea cairica</i>	climber	exotic	very common	+	+		+	+
<i>Ipomoea pes-caprae</i>	perennial herb	native	common			+		+
<i>Lantana camara</i>	shrub	exotic	very common	+			+	+
<i>Leucaena leucocephala</i>	tree	exotic	common				+	
<i>Mallotus paniculatus</i>	tree	native	very common				+	+
<i>Melastoma malabathricum</i> L.	herb	native	common			+		
<i>Microstegium ciliatum</i>	perennial procumbent herb	native	very common		+	+	+	+
<i>Mikania micrantha</i>	climber	exotic	very common	+		+	+	+
<i>Mimosa diplotricha</i>	herb	exotic	rare	+	+	+	+	+
<i>Miscanthus floridulus</i>	perennial herb	native	common	+	+			+
<i>Oxalis corymbosa</i>	perennial herb	exotic	common					+
<i>Paspalum conjugatum</i>	perennial herb	exotic	common				+	
<i>Plantago major</i>	herb	native	very common		+			
<i>Polygonum chinense</i>	herb	native	very common			+		
<i>Praxelis clematidea</i>	perennial herb	exotic	very common					+
<i>Pueraria phaseoloides</i>	climber	native	very common		+	+		+
<i>Saccharum arundinaceum</i>	herb	native	N/A				+	+
<i>Sageretia thea</i>	shrub	native	very common	+	+		+	
<i>Sapium sebiferum</i>	tree	native	common	+	+			+
<i>Urena lobata</i>	shrub	native	common		+	+	+	+
<i>Wedelia trilobata</i>	perennial herb	exotic	common	+	+	+	+	+

Note:

Code: +=occurrence of the species

Appendix 3: Ecological Water Quality Monitoring Raw Data

(April 2013)

Date of Monitoring: 23 April 2013

Weather : Sunny

Monitoring Location	Suspended Solids (mg/L)	Nitrogen (Ammonia) (mg/L)	Nitrogen (Nitrate) (mg/L)	Reactive Phosphorous (mg/L)	5-day Biochemical Oxygen Demand (BOD5) (mg/L)	Dissolved Oxygen (mg/L)	
						M1	M2
WE1	3.0	0.02	0.17	0.03	<2.0	7.58	7.46
WE2	3.0	0.18	0.26	0.04	<2.0	7.56	7.54
WE3	3.0	0.16	0.25	0.05	<2.0	7.46	7.40
WE4	3.0	0.57	0.56	0.06	<2.0	6.89	6.77
WE5	3.0	1.95	0.35	0.11	<2.0	5.55	5.53
WE6	4.0	0.04	0.40	0.01	<2.0	7.61	7.59
WE7	No water - Not sampled						
WE8	No water - Not sampled						
WE9	No water - Not sampled						
WE10	No water - Not sampled						

Monitoring Location	Temperature (°C)		pH		Salinity (ppt)		Conductivity (µs/cm)	Water Flow (m/s)	Water Depth (cm)
	M1	M2	M1	M2	M1	M2			
WE1	20.5	20.5	6.28	6.29	0.02	0.02	66.2	<0.1	53
WE2	20.7	20.8	6.83	6.83	0.04	0.04	122.3	<0.1	12
WE3	20.8	20.8	6.85	6.85	0.03	0.03	124.9	<0.1	18
WE4	22.3	22.3	7.39	7.41	9.98	10.56	17411	<0.1	22
WE5	21.8	21.8	7.03	7.03	2.50	2.43	46.6	<0.1	18
WE6	20.9	20.9	6.34	6.36	0.03	0.03	75.4	<0.1	33
WE7	No water - Not sampled								
WE8	No water - Not sampled								
WE9	No water - Not sampled								
WE10	No water - Not sampled								

Note:

Where more than one measurement was taken, the data is represented by Measurement M1 and M2.