



**Agreement No. DP 04/2012**  
**Post-Construction Ecological Monitoring**  
**of Drainage Improvement Works in Southern Lantau**  
**Implemented under 4128CD in Contract DC/2006/11**

**Monthly EM&A Report – October 2013**

November 2013

	Name	Signature
Prepared & Checked:	Chiu Ming Ho (Ecologist)	
Reviewed & Approved:	Sharne McMillan (Environmental Team Leader)	
Version:	2	Date: November 2013

AECOM Asia Co. Ltd.

8/F, Grand Central Plaza, Tower 2, 138 Shatin Rural Committee Road, Shatin, NT, Hong Kong

Tel: (852) 3922 9000 Fax: (852) 3922 9797 www.aecom.com

**Pursuant to Condition 4.3 of Environmental Permit No. EP-237/2005/B, this monthly EM&A Report for post-construction ecological monitoring and ecological water monitoring during October 2013 has been certified by the Environmental Team Leader (ETL) and verified by the Independent Environmental Checker (IEC)**

Certified by:

Signature: 

Ms. Shame McMillan  
Environmental Team Leader (ETL)  
AECOM Asia Co. Ltd

Date: 26/11/2013

Verified by:

Signature: 

Mr. Roger Leung  
Independent Environmental Checker (IEC)  
ENVIRON Hong Kong Limited

Date: 26/11/2013

## Table of Contents

	Page
<b>EXECUTIVE SUMMARY .....</b>	<b>1</b>
<b>1. INTRODUCTION .....</b>	<b>2</b>
1.1. Background .....	2
1.2. Project Description .....	2
1.3. Report Objectives.....	2
<b>2. ECOLOGICAL MONITORING PARAMETERS .....</b>	<b>2</b>
2.1. Ecological Surveys .....	2
2.2. Ecological Water Quality Monitoring.....	5
2.3. Limitations.....	6
<b>3. MONITORING RESULTS .....</b>	<b>8</b>
3.1. Ecological Survey Findings.....	8
3.2. Ecological Water Quality Monitoring (EWQM) .....	20
<b>4. ECOLOGICAL MONITORING SCHEDULE .....</b>	<b>21</b>
<b>5. DISCUSSION AND RECOMMENDATIONS.....</b>	<b>21</b>
<b>6. REFERENCES .....</b>	<b>24</b>

### List of Figures

- Figure 1 Ecological Monitoring Locations at Pak Ngan Heung River, Luk Tei Tong River, and Luk Tei Tong Bypass Channel and the Reference Site
- Figure 2 Ecological Water Quality Monitoring Locations at Pak Ngan Heung River, Luk Tei Tong River, Luk Tei Tong Bypass Channel and the Reference Site

### List of Tables

- Table 2.1 Limit of Reporting for Water Quality Parameters
- Table 3.1 Number of Avifauna Recorded at Pak Ngan Heung River (PNH)
- Table 3.2 Number of Dragonfly Recorded at Pak Ngan Heung River (PNH)
- Table 3.3 Relative Abundance of Aquatic Macroinvertebrate and Fish Recorded at Pak Ngan Heung River (PNH)
- Table 3.4 Number of Avifauna Recorded at Luk Tei Tong River (LTT)
- Table 3.5 Relative Abundance of Aquatic Macroinvertebrate and Fish Recorded at Luk Tei Tong River (LTT)
- Table 3.6 Vegetation Coverage at Luk Tei Tong Bypass Channel (LBC) and Reference Site (RS)
- Table 3.7 Number of Avifauna Recorded at Luk Tei Tong Bypass Channel (LBC)
- Table 3.8 Number of Avifauna Recorded at Reference Site (RS)
- Table 3.9 Number of Dragonfly Recorded at Luk Tei Tong Bypass Channel (LBC)
- Table 3.10 Relative Abundance of Aquatic Macroinvertebrate and Fish Recorded at Luk Tei Tong Bypass Channel (LBC)
- Table 3.11 Summarized Ecological Water Quality Monitoring Results (October 2013)
- Table 3.12 Baseline Results of Ecological Water Quality Monitoring Results (September 2007)
- Table 5.1 Key Observations/Comments and Recommendations Arising from the October 2013 Monitoring Period

**List of Appendices**

- Appendix 1 Calibration Certificate of the Instruments (pH Meter and Multi-meter)
- Appendix 2a Plant Species Recorded in Pak Ngan Heung River and Luk Tei Tong River
- Appendix 2b Plant Species Recorded in Luk Tei Tong Bypass Channel and the Reference Site
- Appendix 3 Ecological Water Quality Monitoring – Raw Data
- Appendix 4 Representative Photographs taken during the Monitoring

## **EXECUTIVE SUMMARY**

This is the seventh bi-monthly post-construction ecological monitoring and audit exercise for "Drainage Improvement in Southern Lantau" conducted by AECOM. This report concludes the post-construction phase ecological monitoring and audit requirement for the activities undertaken during the period of 1 October 2013 to 31 October 2013.

Ecological monitoring and ecological water quality monitoring were performed on 4 October 2013 and 25 October 2013, respectively. Results obtained are presented in this report.

The Environmental Team (ET) will continue to implement the environmental monitoring & audit (EM&A) programme in accordance with the EM&A Manual and Environmental Permit requirement. The report is available for public inspection and will be uploaded to the dedicated project website (<http://www.envproject.com/sldiwema.htm>).

## **1. INTRODUCTION**

### **1.1. Background**

1.1.1. The Drainage Services Department (DSD) has implemented Contract No. DC/2006/11 “Drainage Improvement in Southern Lantau and Construction of Mui Wo Village Sewerage Phase 1”. The monitoring requirements of the drainage improvement works are subject to the conditions specified in Environmental Permit (EP) No. EP-237/2005/B issued by the Environmental Protection Department (25 January 2006). In compliance with the EP, an Environmental Monitoring and Audit (EM&A) programme was established during the construction and post-construction phases of the project. The operation of the project is subject to the conditions in EP No. EP-434/2012.

1.1.2. The Post-Construction Ecological Monitoring and Audit of Drainage Improvement Works in Southern Lantau under Agreement No. DP 04/2012, commenced in January 2012. AECOM Asia Co. Ltd. was appointed by DSD as the Environmental Team (ET) to conduct the above captioned monitoring project from October 2012 onwards. This is the seventh bi-monthly post-construction ecological monitoring and audit report under that appointment.

### **1.2. Project Description**

1.2.1. Under Contract No. DC/2006/11, the improvement works were undertaken at Pak Ngan Heung River (PNH), Luk Tei Tong River (LTT) and Tai Tei Tong River (TTT) in Southern Lantau, west of Mui Wo. The works for which the post-construction ecological monitoring required by EP No. EP-237/2005/B covered included:

- the drainage channel and a three-cell box culvert at PNH;
- the drainage channel at LTT; and
- the bypass channel at LTT.

1.2.2. No ecological monitoring and ecological water monitoring was required following the drainage improvement works at TTT and village sewerage works in Mui Wo.

1.2.3. Both PNH and LTT are part of the Mui Wo River (also named as Silver River) in Lantau Island. These two tributaries of Mui Wo River, together with Tai Tei Tong River, then joined and connected to Silver Mine Bay next to Mui Wo.

### **1.3. Report Objectives**


1.3.1. This report presents the findings of the ecological monitoring and the ecological water monitoring conducted in October 2013.

## **2. ECOLOGICAL MONITORING PARAMETERS**

### **2.1. Ecological Surveys**

2.1.1. Details of the monitoring parameters and survey methodology are described below. According to the Final EM&A Manual, a specific ecological monitoring programme of the improved section of PNH, LTT, LTT Bypass Channel (LBC) and its Reference Site (RS) is recommended.

Figure 1 Ecological Monitoring Locations at Pak Ngan Heung River, Luk Tei Tong River, Luk Tei Tong Bypass Channel and the Reference Site



*Pak Ngan Heung River and Luk Tei Tong River*

- 2.1.2. The ecological survey for these two rivers was divided into nine 50 m sections and comprised the following:
- Two sections for downstream of PNH (PNH1 and 2), two sections for upstream of PNH (PNH3 and 4);
  - Five sections for LTT (LTT1 to 5).
- 2.1.3. The location plan is shown in **Figure 1** for reference.
- 2.1.4. The monitoring parameters and survey methodology for each section are described below:
- (a) Bird species in each 50 m section were surveyed quantitatively using transect count method. Birds within the river channel and on the riverbank were identified to species and their abundance was recorded. Birds that flew over/across the river channel without landing were not considered to be utilising the area and thus excluded from the records. This does not apply to species that rarely land and are associated with specific habitats (e.g. Barn Swallow).
  - (b) Surveys on aquatic macroinvertebrate focused on determination of the diversity and abundance. Sampling methods included active searching, direct observation, hand netting and kick sampling. In each section, the macroinvertebrate species composition was identified and their relative abundance was recorded.
  - (c) Surveys on fish focused on determination of the diversity and abundance of fish communities. Sampling methods included active searching, direct observation, and hand netting, and were determined in accordance with site conditions. In each section, the fish species composition was identified and their relative abundance was recorded.
  - (d) Adult odonate community in each 50 m section were surveyed quantitatively by transect count method. Adult odonates within the river channel and on the riverbank were identified to species and their abundance was recorded. Species requiring close examination were netted.
  - (e) Aquatic, emergent and riparian vegetation community was recorded by walk-through survey. Plant species composition and their relative abundance were recorded.

*Luk Tei Tong Bypass Channel*

- 2.1.5. The ecological survey for the Luk Tei Tong Bypass Channel (LBC) and its Reference Sites (RS) were carried out in every 50 m section and comprised the following:
- Five sections for LBC (LBC1 to 5);
  - Five sections for RS (RS1 to 5).
- 2.1.6. The location plan is shown in **Figure 1** for reference.
- 2.1.7. The monitoring parameters and survey methodology are described below:
- (a) Bird species in each 50 m section were surveyed quantitatively using transect count method. Birds within the river channel and on the riverbank were identified to species and their abundance was recorded. Birds that flew over/across the river channel without landing were not considered to be utilising the area and thus excluded from the records. This does not apply to species that rarely land and are associated with specific habitats (e.g. Barn Swallow).
  - (b) Where/when water was present, surveys of aquatic macroinvertebrate focused on determination of their diversity and abundance of stream aquatic communities. Sampling



methods included active searching, direct observation, hand netting and kick sampling. In each section, macroinvertebrate species composition was identified and their relative abundance was recorded.

- (c) Where/when water was present, surveys of fish focused on determination of their diversity and abundance. Sampling methods included active searching, direct observation, and hand netting, were determined in accordance with site conditions. In each section, fish species composition was identified and their relative abundance was recorded.
  - (d) Adult odonate community in each 50 m section were surveyed quantitatively by transect count method. Adult dragonflies within the river channel and on the riverbank were identified to species and their abundance was recorded. Species requiring close examination were netted.
  - (e) Line-intercept method was adopted to determine the relative plant cover of aquatic, emergent and riparian vegetation. One line transect of 10 m was set perpendicular to the stream channel at each section, and five 1 m x 1 m quadrats were placed along the transect. Relative coverage and plant species intercepting the transect line was recorded. Percentage cover of each species within the quadrat was recorded to the nearest 10% (except "1" = present but insignificant cover, normally 1 to 2 individuals, and 5% = up to 5%). The conditions of vegetation were described.
  - (f) Herpetofauna community within LBC and RS were surveyed by active searching in potential habitats. Reptiles were identified and their abundance was recorded. Amphibians were identified by their calls and the number of calling males in each section was recorded.
- 2.1.8. For all surveys, identification of plant species and distribution status in Hong Kong were made with reference to Corlett *et al.* (2000), Hu *et al.* (2003), Hong Kong Herbarium (2012), and Hong Kong Herbarium and South China Botanical Gardens (2007; 2008; 2009; 2011).
- 2.1.9. In terms of assessing geographical distribution, published references specializing in the distribution of specific faunal groups in Hong Kong have been utilized. For general status, these have included Fellowes *et al.* (2002) and the Hong Kong Biodiversity Database (AFCD, 2013), and for specific faunal groups, these have included: Avifauna – Carey *et al.* (2001), Viney *et al.* (2006); Dragonflies – Tam *et al.* (2011); Butterflies – Lo (2005); and Chan *et al.* (2011); Amphibians – Chan *et al.* (2005); Reptiles – Chan *et al.* (2006), Chan *et al.* (2009), and Karsen *et al.* (1998); Terrestrial Mammals – Shek (2006); Freshwater Fish – Lee *et al.* (2004); and Freshwater Community – Dudgeon (2003). The status and rarity of vascular plants has been based on Hu *et al.* (2003) and Corlett *et al.* (2000).

## 2.2. Ecological Water Quality Monitoring

- 2.2.1. Ecological water quality monitoring along PNH, LTT, LBC, and RS was carried out. Ten locations were selected and comprised the following:
- Three locations for existing PNH (WE1 to 3);
  - Three locations for existing LTT (WE4 to 6);
  - Two locations for RS (WE7 to 8);
  - Two locations for existing LBC (WE9 to 10).
- 2.2.2. The location plan for ecological water quality monitoring is shown in **Figure 2**.
- 2.2.3. Water Quality Monitoring along PNH, LTT, LBC and RS included the monitoring parameters shown below:

- Biochemical Oxygen Demand (BOD<sub>5</sub>)
- Nitrate
- Ammonia
- Reactive Phosphorus
- Total Suspended Solids (SS)
- Temperature
- Dissolved Oxygen (DO)
- Water Depth\* and Water Flow Rate
- Conductivity
- pH
- Salinity
- Sediment Characteristics

Note:

\*As referred to in the Final EM&A Manual, Water Depth is required only for LBC.

2.2.4. The DO, water depth and water flow rate, conductivity, pH, temperature, salinity and sediment characteristics were measured in-situ while the other water samples were analyzed in a HOKLAS accredited laboratory and the analyses followed the standard methods according to APHA Standard Methods for the Examination of Water and Wastewater, 20th Edition, or equivalent. The limit of reporting for the laboratory analysis is summarized in **Table 2.1**.

**Table 2.1 Limit of Reporting for Water Quality Parameters**

Parameters	Limit of Reporting (mg/L)
Total Suspended Solids	2
Biochemical Oxygen Demand (BOD <sub>5</sub> )	2
Nitrate	0.01
Ammonia	0.01
Reactive Phosphorus	0.01

2.2.5. The instrument for in-situ measurement of temperature, DO, salinity and conductivity is a portable and weather proof multi-meter complete with cable and uses a DC power source (YSI 85), whereas Orion 230A+ is used as for pH measurement. Calibration certificates are attached in **Appendix 1**. The instruments are capable of measuring:


- pH in the range of 0 to 14
- Temperature of -5 to +65<sup>0</sup>C
- DO in the range of 0 to 20 mg/L and 0 to 200% saturation
- Salinity in the range of 0-80ppt
- Conductivity in the range of 0 to 4999 μS/cm

2.2.6. According to the requirement of the Final EM&A Manual, two consecutive measurements for parameters of DO concentration, and DO saturation are required to be taken at each monitoring location. When the difference in value between the first and second reading of DO is more than 25%, the reading was discarded and a further reading taken.

### 2.3. Limitations

- 2.3.1. No water was present at LBC2 to LBC5 at the time of survey, therefore aquatic fauna surveys were not undertaken in these locations.
- 2.3.2. No water was present at WE7 - WE10 at the time of survey, therefore water quality monitoring was not undertaken at these locations.

**Figure 2 Ecological Water Quality Monitoring Locations at Pak Ngan Heung River, Luk Tei Tong River, Luk Tei Tong Bypass Channel and the Reference Site**



### 3. MONITORING RESULTS

#### 3.1. Ecological Survey Findings

##### *Pak Ngan Heung River (PNH)*

- 3.1.1. The lower stream of PNH (PNH1 and PNH2) is subject to tidal influence from Silver Mine Bay. Vertical concrete retaining wall formed the banks of the river channel. The two sections were located at the mouth of the PNH. PNH1 and PNH2 were adjacent to each other. The bridge formed the southern boundary of PNH1 whereas the box-culvert formed the northern boundary of PNH2. Small boulders and sandy substrate formed the main component of the streambed.
- 3.1.2. Rock-filled gabion formed the eastern bank and the gabion and a vertical concrete retaining wall formed the western bank of the upper stream (PNH3 and PNH4). PNH3 and PNH4 are adjacent to each other. PNH4 comprised a man-made cascade, including a fish ladder, while PNH3 comprised a pool below the cascade and was bounded by a bridge at its downstream end. Small boulders and sandy substrate were the main component in the middle streambed which allowed water flow and pool formation, whereas big boulders were scattered on both sides of the streambed and had an absence of water. The width of the fish ladder at PNH4 is approximately 7 m.
- 3.1.3. The cascade/fish ladder at PNH4 was open and free of vegetation, exposing the feature and allowing free water flow and pool formation.

##### Vegetation

- 3.1.4. At PNH3 and PNH4, a total of 15 plant species were recorded. The vegetation was dominated by exotic species, Mile-a-minute (*Mikania micrantha*), at both PNH3 and PNH4. The vegetation predominantly grew on the banks of PNH3 pool and outer edges of the PNH4 cascade. Species such as *Polygonum barbatum* and *Ludwigia octovalvis* were scattered along the edge of the PNH4 cascade.
- 3.1.5. At PNH1 and PNH2, no plant species were recorded within the river channel. The vegetation recorded on the vertical wall has not changed significantly since the last monitoring period, which includes a record of *Wedelia trilobata* as well as seedlings of *Kandelia obovata* and Opposite-leaved Fig (*Ficus hispida*).
- 3.1.6. The list of plant species is presented in **Appendix 2a**.

##### Terrestrial Fauna

- 3.1.7. Twelve avifauna species were recorded at PNH, all of which are common and abundant in Hong Kong (**Table 3.1**). Four avifauna species were recorded at lower PNH (PNH1 and PNH2). Two species of conservation importance, Little Egret (*Egretta garzetta*) and Common Redshank (*Tringa tetanus*), were recorded roosting within the river channel. They are waterbirds, which are considered as Potential Regional Concern and Regional Concern respectively by Fellowes *et al.* (2002). Other recorded species were common resident species. During the monitoring, the water level at lower PNH was approximately 30 cm.
- 3.1.8. Eight avifauna species were recorded at upper PNH. The birds at PNH3 and PNH4 were mostly observed along the banks of the river channel, and were dominated by resident species. A group of five Large-billed Crow (*Corvus macrohynchos*) were recorded perching near the bank of PNH3. Other recorded species included Asian Koel (*Eudynamis scolopacea*) and Blue Whistling Thrush (*Myophonus caeruleus*). All recorded species are common and abundant in Hong Kong (AFCD, 2013).
- 3.1.9. A total of four odonate species were recorded at PNH. One species, Crimson Dropwing (*Trithemis aurora*), was recorded at PNH1 in low abundance. Four species were recorded at PNH3. All of the recorded species are abundant in Hong Kong (**Table 3.2**).
- 3.1.10. No herpetofauna was recorded at PNH.

##### Aquatic Macroinvertebrate and Fish

- 3.1.11. At lower PNH (PNH1 and PNH2), four fish species, one crab species and four other aquatic macroinvertebrate species (such as the Sea Slater, *Ligia exotica*) were recorded (**Table 3.3**). One species of conservation importance, Predaceous Chub (*Parazacco spilurus*), was recorded in low numbers at PNH2. Predaceous Chub although is one of the most common freshwater fishes in local streams, it is considered as “Vulnerable” in China Red Data Book (Lee *et al.*, 2004). Other recorded fish species are common in Hong Kong, such as Jarbua Terapon (*Terapon jarbua*) (AFCD, 2013).
- 3.1.12. At upper PNH (PNH3 and PNH4), no fish but four aquatic macroinvertebrate species were recorded (**Table 3.3**). Baetidae was dominated at both PNH3 and PNH4.

**Table 3.1 Number of Avifauna Recorded at Pak Ngan Heung River (PNH)**

Common Name <sup>(1)</sup>	Scientific Name	Distribution in Hong Kong <sup>(2)</sup>	Principal Status <sup>(3)</sup>	Level of Concern <sup>(4)</sup>	Protection Status in China <sup>(5)</sup>	China Red Data Book <sup>(6)</sup>	IUCN Red List <sup>(7)</sup>	PNH1	PNH2	PNH3	PNH4
<b>Little Egret</b> <sup>(8)</sup>	<i>Egretta garzetta</i>	Common	P	PRC (RC)	-	-	-		1		
<b>Common Redshank</b> <sup>(8)</sup>	<i>Tringa totanus</i>	Common	W	RC	-	-	-	1			
Spotted Dove	<i>Streptopelia chinensis</i>	Abundant	R	-	-	-	-	3			
Asian Koel	<i>Eudynamys scolopacea</i>	Common	Su,R	-	-	-	-			1	
Large-billed Crow	<i>Corvus macrorhynchos</i>	Common	R	-	-	-	-			5	
Great Tit	<i>Parus major</i>	Common	R	-	-	-	-			2	
Red-whiskered Bulbul	<i>Pycnonotus jocosus</i>	Abundant	R	-	-	-	-			2	
Yellow-bellied Prinia	<i>Prinia flaviventris</i>	Common	R	-	-	-	-				1
Japanese White-eye	<i>Zosterops japonicus</i>	Abundant	R,?W	-	-	-	-			3	
Blue Whistling Thrush	<i>Myophonus caeruleus</i>	Common	R	-	-	-	-			1	
Oriental Magpie Robin	<i>Copsychus saularis</i>	Abundant	R	-	-	-	-			1	
White Wagtail	<i>Motacilla alba</i>	Common	W,R	-	-	-	-	1			

Note:

(1) All wild birds are protected under Wild Animal Protection Ordinance (Cap. 170).

(2) AFCD (2013) Hong Kong Biodiversity Database.

(3) R=resident; Su=summer; W=winter; P=present all year, exact composition unknown; ?W=the extent of immigration in winter is unclear.

(4) Fellowes *et al.* (2002); RC=Regional Concern; PRC =Potential Regional Concern; Letters in parentheses indicate that the assessment is on the basis of restrictedness in breeding and/or roosting sites rather than in general occurrence.

(5) List of Wild Animals Under State Protection (promulgated by State Forestry Administration and Ministry of Agriculture on 14 January, 1989).

(6) Zheng and Wang (1998).

(7) IUCN (2013). IUCN Red List of Threatened Species. Version 2013.1.

(8) Wetland-dependent species (including wetland-dependent species and waterbirds).

Species of conservation importance is noted in bold type face

**Table 3.2 Number of Dragonfly Recorded at Pak Ngan Heung River (PNH)**

Common Name	Scientific Name	Distribution in Hong Kong <sup>(1)</sup>	Level of Concern <sup>(2)</sup>	Protection Status in China <sup>(3)</sup>	China Red Data Book <sup>(4)</sup>	IUCN Red List <sup>(5)</sup>	PNH 1	PNH 2	PNH 3	PNH 4
Black Threadtail	<i>Prodasineura autumnalis</i>	Abundant	-	-	-	-			3	
Common Red Skimmer	<i>Orthetrum prunosum neglectum</i>	Abundant	-	-	-	-			2	
Crimson Dropwing	<i>Trithemis aurora</i>	Abundant	-	-	-	-	1		2	
Indigo Dropwing	<i>Trithemis festiva</i>	Abundant	-	-	-	-			2	

Note:

- (1) AFCD (2013). Hong Kong Biodiversity Database.
- (2) Fellowes *et al.* (2002).
- (3) List of Wild Animals Under State Protection (promulgated by State Forestry Administration and Ministry of Agriculture on 14 January, 1989).
- (4) Zheng and Wang (1998).
- (5) IUCN (2013). IUCN Red List of Threatened Species. Version 2013.1.

**Table 3.3 Relative Abundance of Aquatic Macroinvertebrate and Fish Recorded at Pak Ngan Heung River (PNH)**

Fauna Group	Scientific Name	Common Name	Distribution in Hong Kong <sup>(1)(2)(3)</sup>	Level of Concern <sup>(4)</sup>	Protection Status in China <sup>(5)</sup>	China Red Data Book <sup>(6)</sup>	IUCN Red List <sup>(7)</sup>	PNH1	PNH2	PNH3	PNH4
Fish	<b><i>Parazacco spilurus</i></b>	<b>Predaceous Chub</b>	Common	-	-	Vulnerable	-		+		
Fish	<i>Goby sp.</i>	-	-	-	-	-	-		+		
Fish	<i>Terapon jarbua</i>	Jarbua Terapon	Common	-	-	-	-	++			
Fish	<i>Cyprinidae sp.</i>	-	-	-	-	-	-	++			
Crabs	<i>Sesarmops sinensis</i>	-	-	-	-	-	-		++		
Oligochaeta	Oligochaeta	-	-	-	-	-	-			+++	++
Worms	<i>Capitella capitata</i>	-	-	-	-	-	-			+	+
Snail	<i>Clithon sp.</i>	-	-	-	-	-	-	++++			
Snails	Lymnaeidae	-	-	-	-	-	-	++			
Amphipod	Amphipoda	-	-	-	-	-	-	++	++	++	
Insects	<i>Ligia exotica</i>	Sea Slater	Common	-	-	-	-	+			
Insects	Baetidae	-	-	-	-	-	-			+++	+++

Note:

(1) AFCD (2013). Hong Kong Biodiversity Database.

(2) Williams, G. (2003). Hong Kong Field Guides – Rocky Shores.

(3) Chan *et al.* (2003). Hong Kong Field Guides – Sandy Shores.

(4) Fellowes *et al.* (2002).

(5) List of Wild Animals Under State Protection (promulgated by State Forestry Administration and Ministry of Agriculture on 14 January, 1989).

(6) Zheng and Wang (1998).

(7) IUCN (2013). IUCN Red List of Threatened Species. Version 2013.1.

Relative abundance: + = occasional, less than 5 individuals were found; ++ = common, 5-20 individuals were found; +++ = abundant, more than 20 individuals were found.

Species of conservation importance is noted in bold type face



### *Luk Tei Tong River (LTT)*

- 3.1.13. The LTT is subject to tidal influence from Silver Mine Bay and is estuarine in nature. It is a north-south running river. A vertical concrete retaining wall formed the riverbank of the LTT1 whereas rock-filled gabion formed the riverbank of LTT2 to LTT5. LTT1 was located at the confluence with Pak Ngan Heung River, Tai Tei Tong River and Luk Tei Tong River. Since it is subject to tidal flow, water flowed from south to north during the survey when the tide was going out. LTT1 and LTT2 had sandy substrate whilst LTT3 to LTT5 had muddy substrate. Clusters of boulders occurred at both sides of the river channel. The width of the river channel was approximately 8-10 m.
- 3.1.14. No evidence of maintenance works (including those relevant to Conditions 2.1 to 2.4 of EP No. EP-434/2012) was observed during the monitoring period.

#### Vegetation

- 3.1.15. A total of 12 plant species were recorded in LTT. Five out of 12 of the recorded species were exotic. The majority were herbs or climbers scattered along the gabion such as *Bidens alba*. In addition to the mangrove stand (Spiny Bears Breech, *Acanthus ilicifolius* and *Kandelia obovata*) that colonized inside the river channel at LTT2 and LTT3, several seedlings of *Kandelia obovata* have naturally regenerated in LTT1, LTT2, LTT3 and LTT5. Herbaceous species such as Mile-a-minute (*Mikania micrantha*) and Sea Sword Bean (*Canavalia maritima*) were recorded on the gabion at LTT2, LTT3, LTT4 and LTT5.
- 3.1.16. The list of plant species is presented in **Appendix 2a**.

#### Terrestrial Fauna

- 3.1.17. A total of 16 avifauna species were recorded at LTT, all of them are common and abundant in Hong Kong (AFCD, 2013) (**Table 3.4**). Waterbirds and wetland dependent species, including ardeids (Black-crowned Night Heron *Nycticorax nycticorax*, Grey Heron *Ardea cinerea* and Little Egret), Common Sandpiper (*Actitis hypoleucos*) and Common Kingfisher (*Alcedo atthis*) were recorded along the main river channel. Other recorded species included generalist species (such as Crested Myna *Acridotheres cristatellus*). Among the recorded species, Grey Heron and Little Egret are of conservation importance for they are considered as Potential Regional Concern by Fellowes *et al.* (2002).
- 3.1.18. No dragonfly and herpetofauna species were recorded at the LTT during the monitoring.

#### Aquatic Macroinvertebrate and Fish

- 3.1.19. A total of six fish species, one crab species and seven species of other aquatic macroinvertebrate were recorded from the LTT (**Table 3.5**).
- 3.1.20. All fish species recorded mainly occur in river mouth or estuarine environments in Hong Kong (AFCD, 2013). Among the recorded species, Predaceous Chub, which is a species of conservation importance, was recorded. It is considered as "Vulnerable" in China Red Data Book (Lee, *et al.*, 2004). Whilst one fish in the family of Cyprinidae was recorded in all sections of LTT.

**Table 3.4 Number of Avifauna Recorded at Luk Tei Tong River (LTT)**

Common Name <sup>(1)</sup>	Scientific Name	Distribution in Hong Kong <sup>(2)</sup>	Principal Status <sup>(3)</sup>	Level of Concern <sup>(4)</sup>	Protection Status in China <sup>(5)</sup>	China Red Data Book <sup>(6)</sup>	IUCN Red List <sup>(7)</sup>	LTT1	LTT2	LTT3	LTT4	LTT5
Black-crowned Night Heron <sup>(8)</sup>	<i>Nycticorax nycticorax</i>	Common	P	(LC)	-	-	-		1			
<b>Grey Heron</b> <sup>(8)</sup>	<i>Ardea cinerea</i>	Common	W	PRC	-	-	-	1				1
<b>Little Egret</b> <sup>(8)</sup>	<i>Egretta garzetta</i>	Common	P	PRC (RC)	-	-	-	1				
Common Sandpiper <sup>(8)</sup>	<i>Actitis hypoleucos</i>	Common	M,W	-	-	-	-	1				
Spotted Dove	<i>Streptopelia chinensis</i>	Abundant	R	-	-	-	-	2				
House Swift	<i>Apus nipalensis</i>	Common	R,SpM	-	-	-	-					1
Common Kingfisher <sup>(8)</sup>	<i>Alcedo atthis</i>	Common	AM,P	-	-	-	-					1
Black Drongo	<i>Dicrurus macrocercus</i>	Common	M,Su	-	-	-	-					1
Red-whiskered Bulbul	<i>Pycnonotus jocosus</i>	Abundant	R	-	-	-	-	2				
Yellow-bellied Prinia	<i>Prinia flaviventris</i>	Common	R	-							1	
Plain Prinia	<i>Prinia inornata</i>	Common	R	-							1	
Common Tailorbird	<i>Orthotomus sutorius</i>	Common	R	-								1
Japanese White-eye	<i>Zosterops japonicus</i>	Abundant	R,?W	-							5	
Crested Myna	<i>Acridotheres cristatellus</i>	Common	R	-								30
Grey Wagtail	<i>Motacilla cinerea</i>	Common	W	-				1				
White Wagtail	<i>Motacilla alba</i>	Common	W,R	-				1				

Note:

(1) All wild birds are protected under Wild Animal Protection Ordinance (Cap. 170).

(2) AFCD (2013). Hong Kong Biodiversity Database.

(3) R=resident; A=Autumn; Su=summer; W=winter visitor; Sp=spring; M=migrant; P=present all year, exact composition unknown; ?W=the extent of immigration in winter is unclear.

(4) Fellowes *et al.* (2002); LC=Local Concern; RC=Regional Concern; PRC=Potential Regional Concern. Letters in parentheses indicate that the assessment is on the basis of restrictedness in breeding and/or roosting sites rather than in general occurrence.

(5) List of Wild Animals Under State Protection (promulgated by State Forestry Administration and Ministry of Agriculture on 14 January, 1989).

(6) Zheng and Wang (1998).

(7) IUCN (2013). IUCN Red List of Threatened Species. Version 2013.1.

(8) Wetland-dependent species (including wetland-dependent species and waterbirds).  
 Species of conservation importance is noted in bold type face.

**Table 3.5 Relative Abundance of Aquatic Macroinvertebrate and Fish Recorded at Luk Tei Tong River (LTT)**

Fauna Groups	Scientific Name	Common Name	Distribution in Hong Kong (1)(2)(3)	Level of Concern (4)	Protection Status in China (5)	China Red Data Book (6)	IUCN Red List (7)	LTT1	LTT2	LTT3	LTT4	LTT5
Fish	<i>Mugil cephalus</i>	Grey Mullet	-	-	-	-	-				++++	++++
Fish	<b><i>Parazacco spilurus</i></b>	<b>Predaceous Chub</b>	Common	-	-	Vulnerable	-	+++				++
Fish	<i>Periophthalmus cantonensis</i>	-	Very common	-	-	-	-		+			+
Fish	<i>Goby sp.</i>	-	-	-	-	-	-					+
Fish	<i>Terapon jarbua</i>	Jarbua Terapon	Common	-	-	-	-	++++	++			
Fish	Cyprinidae	-	-	-	-	-	-	+++++	+++	+++	+++	++++
Crabs	<i>Sesarmops sinensis</i>	-	-	-	-	-	-			+		
Snail	<i>Clithon sp.</i>	-	-	-	-	-	-	++++				
Snail	<i>Nerita sp.</i>	-	-	-	-	-	-	++++				
Snails	Lymnaeidae	-	-	-	-	-	-	+				
Bivalves	<i>Saccostrea cucullata</i>	Rock oyster	Very common	-	-	-	-	+++	++	+		
Barnacles	<i>Balanus amphitrite</i>	-	Very common	-	-	-	-	++	+	++		
Amphipod	Amphipoda	-	-	-	-	-	-	+	+			
Insects	<i>Ligia exotica</i>	Sea Slater	Common	-	-	-	-	+		+	+	++

Note:

- (1) AFCD (2013). Hong Kong Biodiversity Database.
- (2) Williams, G (2003). Hong Kong Field Guides – Rocky Shores.
- (3) Chan *et al.* (2003). Hong Kong Field Guides – Sandy Shores.
- (4) Fellowes *et al.* (2002).
- (5) List of Wild Animals Under State Protection (promulgated by State Forestry Administration and Ministry of Agriculture on 14 January, 1989).
- (6) Zheng and Wang (1998).
- (7) IUCN (2013). IUCN Red List of Threatened Species. Version 2013.1.
- (8) Relative abundance: + = occasional, less than 5 individuals were found; ++ = common, 5-20 individuals were found; +++ = abundant, more than 20 individuals were found.  
 Species of conservation importance is noted in bold type face

*Luk Tei Tong Bypass Channel (LBC) and Reference Site (RS)*

- 3.1.21. The LBC is linked to the end of LTT5 and runs east to west but the connection with LTT5 is blocked by a layer of gabion wall approximately 1 m in height, which allows water flow between LBC and LTT when water level is higher than the height of the gabion. It is located in the Luk Tei Tong Marsh to the west of the original LTT. Gabion walls formed both sides of the channel bank. Generally, all sections were heavily vegetated except in LBC1 where a small pool of approximately 60 m<sup>2</sup> in size was located at the western end of LBC1. The pool was separated from the LTT by a weir constructed from a single layer of rock-filled gabion. The substrate comprised soil, which was translocated from the marsh area prior to construction of the bypass. The width of the bypass channel was approximately 15 m.
- 3.1.22. The RS was located parallel to the northern side of the LBC. Next to the RS was village housing. The site was vegetated and did not have any free-standing water at the time of survey.

Vegetation

- 3.1.23. A total of 37 plant species were recorded in LBC, of which 17 species were found in the quadrats sampled. The list of plant species is presented in **Appendix 2b**. Among the recorded species, about 38% were exotic. During the survey, half of the LBC1 section has included a patch of open water. Other sections were dry.
- 3.1.24. The habitat at LBC1 differed from the remaining sections in terms of vegetation type. It adjoined LTT5 and had a pool of open water at the western tip. LBC1 may be subject to tidal influence during high tide because of its location immediately next to LTT. The sedge, Ferruginous-scale *Fimbristylis* (*Fimbristylis sieboldii*), dominated LBC1.
- 3.1.25. The newly established *Polygonum perfoliatum* that was observed in the last monitoring at LBC3 to LBC5 two months ago was not recorded during this monitoring. *Polygonum perfoliatum* is a fringe-of-mangrove species that requires some moisture for establishment. The absence of this species may imply a drier environment compared to the last monitoring two months ago. The exotic species, *Wedelia trilobata*, continues to be the dominant species.
- 3.1.26. Other herbaceous species commonly encountered along the transects were exotic Mile-a-Minute (*Mikania micrantha*) and *Crotalaria pallida*. Other species only formed a small proportion of the vegetation. Records of wetland species such as Taro (*Colocasia esculenta*) and Ginger Lily (*Hedychium coronarium*) were occasional.
- 3.1.27. A total of 36 plant species were recorded in the RS, of which 14 species were found in the quadrats (**Table 3.6**). Half of the 36 species were exotic. All sections were dry and were located next to the village housing. The dominant species was exotic *Wedelia trilobata*. Exotic *Mimosa diplotricha*, *Bidens alba* and *Paspalum conjugatum* were commonly recorded across the RS sections. The majority of vegetation recorded at the RS could typically be found in disturbed land. Records of wetland species such as Taro (*Colocasia esculenta*) and Ginger Lily (*Hedychium coronarium*) were occasional.
- 3.1.28. The list of plant species is presented in **Appendix 2b**.

**Table 3.6 Vegetation Coverage at Luk Tei Tong Bypass Channel (LBC) and Reference Site (RS)**

	LBC	RS
No. of species recorded in quadrats	17	14
Total No. of species	37	36
Total No. of exotic species	14	18
Average vegetation coverage	100%	89%
Bare ground coverage	0%	11%

Note:

(1) The transect was not laid along any open water, thus open water coverage was not provided in this table.

#### Terrestrial Fauna

- 3.1.29. Five species of avifauna were recorded at LBC (**Table 3.7**) whereas three species were recorded at RS (**Table 3.8**). All recorded species are common or abundant in Hong Kong (AFCD, 2013). All recorded avifauna were generalists that have adapted to disturbed environments such as Crested Myna (*Acridotheres cristatellus*) and Black-collared Starling (*Gracupica nigricollis*).
- 3.1.30. Three dragonfly species were recorded at LBC (**Table 3.9**), while no dragonfly species were recorded at RS. All of the recorded species are abundant in Hong Kong (AFCD, 2013). Ten individuals of Wandering Glider (*Pantala flavescens*) were observed flying over LBC1.
- 3.1.31. No herpetofauna were recorded at LBC and RS during the monitoring.

#### Aquatic Macroinvertebrate and Fish

- 3.1.32. Two species of fish were recorded at LBC1 which included Jarbua Terapon, which is a common species in Hong Kong and widespread in estuaries and coastal water, and *Goby* sp. (**Table 3.10**) (AFCD, 2013). Other recorded macroinvertebrate included snails (*Nerita* sp. and Lymnaeidae) and sea slaters. No species of conservation importance were recorded.
- 3.1.33. No aquatic fauna was recorded at the RS or the remaining sections of the LBC as they were dry during the monitoring.

**Table 3.7 Number of Avifauna Recorded at Luk Tei Tong Bypass Channel (LBC)**

Common Name <sup>(1)</sup>	Scientific Name	Distribution in Hong Kong <sup>(2)</sup>	Principal Status <sup>(3)</sup>	Level of Concern <sup>(4)</sup>	Protection Status in China <sup>(5)</sup>	China Red Data Book <sup>(6)</sup>	IUCN Red List <sup>(7)</sup>	LBC1	LBC2	LBC3	LBC4	LBC5
Spotted Dove	<i>Streptopelia chinensis</i>	Abundant	R	-	-	-	-	1		1		
Black Drongo	<i>Dicrurus macrocercus</i>	Common	M,Su	-	-	-	-	1				
Crested Myna	<i>Acridotheres cristatellus</i>	Common	R	-	-	-	-	10		3	3	
Black-collared Starling	<i>Gracupica nigricollis</i>	Common	R	-	-	-	-	1			2	3
White Wagtail	<i>Motacilla alba</i>	Common	W,R	-	-	-	-			1		

Note:

- (1) All wild birds are protected under Wild Animal Protection Ordinance (Cap. 170).
- (2) AFCD (2013). Hong Kong Biodiversity Database.
- (3) R=resident; Su=summer; W=winter; M=migrant.
- (4) Fellowes *et al.* (2002).
- (5) List of Wild Animals Under State Protection (promulgated by State Forestry Administration and Ministry of Agriculture on 14 January, 1989).
- (6) Zheng and Wang (1998).
- (7) IUCN (2013). IUCN Red List of Threatened Species. Version 2013.1.

**Table 3.8 Number of Avifauna Recorded at Reference Site (RS)**

Common Name <sup>(1)</sup>	Scientific Name	Distribution in Hong Kong <sup>(2)</sup>	Principal Status <sup>(3)</sup>	Level of Concern <sup>(4)</sup>	Protection Status in China <sup>(5)</sup>	China Red Data Book <sup>(6)</sup>	IUCN Red List <sup>(7)</sup>	RS1	RS2	RS3	RS4	RS5
Large-billed Crow	<i>Corvus macrorhynchos</i>	Common	R	-	-	-	-					1
Crested Myna	<i>Acridotheres cristatellus</i>	Common	R	-	-	-	-	7	2	5	3	2
Black-collared Starling	<i>Gracupica nigricollis</i>	Common	R	-	-	-	-	2	3	2		2

Note:

- (1) All wild birds are protected under Wild Animal Protection Ordinance (Cap. 170).
- (2) AFCD (2013). Hong Kong Biodiversity Database.
- (3) R=resident.
- (4) Fellowes *et al.* (2002).
- (5) List of Wild Animals Under State Protection (promulgated by State Forestry Administration and Ministry of Agriculture on 14 January, 1989).
- (6) Zheng and Wang (1998).
- (7) IUCN (2013). IUCN Red List of Threatened Species. Version 2013.1.

**Table 3.9 Number of Dragonfly Recorded at Luk Tei Tong Bypass Channel (LBC)**

Common Name	Scientific Name	Distribution in Hong Kong <sup>(1)</sup>	Level of Concern <sup>(2)</sup>	Protection Status in China <sup>(3)</sup>	China Red Data Book <sup>(4)</sup>	IUCN Red List <sup>(5)</sup>	LBC1	LBC2	LBC3	LBC4	LBC5
Orange-tailed Midget	<i>Agriocnemis femina oryzae</i>	Abundant	-	-	-	-		1			
Wandering Glider	<i>Pantala flavescens</i>	Abundant	-	-	-	-	10			1	
Indigo Dropwing	<i>Trithemis festiva</i>	Abundant	-	-	-	-	1				

Note:

- (1) AFCD (2013). Hong Kong Biodiversity Database.
- (2) Fellowes *et al.* (2002).
- (3) List of Wild Animals Under State Protection (promulgated by State Forestry Administration and Ministry of Agriculture on 14 January, 1989).
- (4) Zheng and Wang (1998).
- (5) IUCN (2013). IUCN Red List of Threatened Species. Version 2013.1.

**Table 3.10 Relative Abundance of Aquatic Macroinvertebrate and Fish Recorded at Luk Tei Tong Bypass Channel (LBC)**

Fauna Groups	Scientific Name	Common Name	Distribution in Hong Kong <sup>(1)</sup>	Level of Concern <sup>(2)</sup>	Protection Status in China <sup>(3)</sup>	China Red Data Book <sup>(4)</sup>	IUCN Red List <sup>(5)</sup>	LBC1	LBC2	LBC3	LBC4	LBC5
Fish	<i>Goby sp.</i>	-	-	-	-	-	-	+				
Fish	<i>Terapon jarbua</i>	Jarbua Terapon	Common	-	-	-	-	+				
Snail	<i>Nerita sp.</i>	-	-	-	-	-	-	+++				
Snails	Lymnaeidae	-	-	-	-	-	-	+++				
Insects	<i>Ligia exotica</i>	Sea Slater	Common	-	-	-	-	+				

Note:

- (1) AFCD (2013). Hong Kong Biodiversity Database.
- (2) Fellowes *et al.* (2002).
- (3) List of Wild Animals Under State Protection (promulgated by State Forestry Administration and Ministry of Agriculture on 14 January, 1989).
- (4) Zheng and Wang (1998).
- (5) IUCN (2013). IUCN Red List of Threatened Species. Version 2013.1.  
 Relative abundance: + = occasional, less than 5 individuals were found; ++ = common, 5-20 individuals were found; +++ = abundant, more than 20 individuals were found.

### 3.2. Ecological Water Quality Monitoring (EWQM)

- 3.2.1. The post-construction phase EWQM was conducted on 25 October 2013. The monitoring results are presented in **Appendix 3** and summarised in **Table 3.11**, which includes reference to the key Water Quality Objectives (WQOs). Baseline surveys were conducted in 2007 prior to the start of the drainage improvement works. The baseline survey results are presented in **Table 3.12**.
- 3.2.2. The water quality monitoring results are discussed in **Section 5**.

**Table 3.11 Summarized Ecological Water Quality Monitoring Results (October 2013)**

Parameters	Key Water Quality Objectives <sup>(1)</sup>	WE1	WE2	WE3	WE4	WE5	WE6
Suspended Solids (mg/L)	<20	<2.0	<2.0	<2.0	6.0	3.0	3.0
Nitrogen (Ammonia) (mg/L)	-	0.12	0.11	0.05	0.14	1.64	0.05
Nitrogen (Nitrate) (mg/L)	-	0.10	0.10	0.09	0.21	0.06	0.10
Reactive Phosphorous (mg/L)	-	0.01	0.02	0.02	0.05	0.19	0.01
5-day Biochemical Oxygen Demand (BOD5) (mg/L)	<5	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
Dissolved Oxygen (mg/L)	>4	10.5	10.8	11.7	9.8	10.6	12.0
Temperature (°C)	-	21.3	21.45	22.1	25.55	24.9	22.4
pH	6.5 – 8.5	7.0	7.2	7.4	7.8	7.6	7.9
Salinity (ppt)	-	<0.1	<0.1	0.20	7.51	0.78	<0.1
Conductivity (µs/cm)	-	30.9	335.0	421.4	13089.0	1526.0	30.8
Water Flow (m/s)	-	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1
Water Depth (cm)	-	36	15	17	24	15	28

Note:

- (1) The available key Water Quality Objectives (WQOs) for River Monitoring Stations at Mui Wo River on Lantau Island (EPD, 2011).



**Table 3.12 Baseline Results of Ecological Water Quality Monitoring Results (September 2007)**

Parameters	WE1	WE2	WE3	WE4	WE5	WE6
Suspended Solids (mg/L)	1.0	2.0	3.0	3.0	<1.0	<1.0
Nitrogen (Ammonia) (mg/L)	0.07	0.12	0.11	0.23	0.03	0.02
Nitrogen (Nitrate) (mg/L)	0.12	0.13	0.13	0.31	0.04	0.05
Reactive Phosphorous (mg/L)	0.04	0.06	0.06	0.09	0.06	0.05
5-day Biochemical Oxygen Demand (BOD <sub>5</sub> ) (mg/L)	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0
Dissolved Oxygen (mg/L)	6.58	6.82	6.37	7.61	6.87	5.70
pH	6.4	7.1	7.0	6.8	6.6	6.1
Salinity (ppt)	<0.1	0.1	0.3	7.6	0.1	<0.1

#### 4. ECOLOGICAL MONITORING SCHEDULE

- 4.1. The next ecological surveys monitoring and ecological water quality is tentatively scheduled for mid-December 2013.

#### 5. DISCUSSION AND RECOMMENDATIONS

- 5.1. The aim of the monitoring programme is to provide data on the re-establishment of aquatic/riparian communities in the PNH and LTT, and allow an assessment of the relative success of the mitigation measures to be made. In addition, monitoring of the LBC will assess whether the proposed channel design has provided suitable compensation for the impacts to the Luk Tei Tong Marsh.
- 5.2. Key observations made during the October 2013 monitoring period in relation to the implemented mitigation measures are presented in **Table 5.1**. Where applicable, recommendations for improving the functionality of the mitigation measures have been made for DSD's consideration.
- 5.3. The fringe-of-marsh species, *Polygonum perfoliatum*, which was recorded in last monitoring period two months ago has no record this month. The vegetation composition at LBC3 to LBC5 continues to be dominated by the exotic species, *Wedelia trilobata*. Whilst a low abundance of marsh species (e.g. Taro, *Colocasia esculenta* and Ginger Lily, *Hedychium coronarium*) were also observed. Since *Polygonum perfoliatum* is a fringe-of-mangrove species that requires some moisture for establishment, its absence may imply a dry environment at LBC3 and LBC5.
- 5.4. Tree seedlings (such as *Acacia confusa* and *Sapium sebiferum*) are re-establishing at LBC2 and LBC3. These trees may hinder the re-establishment of marsh habitat.
- 5.5. Seedlings of the mangrove species (*Kandalia obovata*) were seen at other sections of LTT in addition to the existing mangrove stand at the junction at LTT2 and LTT3. Several mangrove seedlings were observed at LTT1, which is the confluence of PNH, TTT and LTT, as well as LTT5, which indicate natural re-colonization of mangrove.
- 5.6. Fish species of conservation importance, Predaceous Chub, that was last recorded in the December 2012 survey of post-construction monitoring, has reappeared at PNH2, LTT1 and LTT5. The status of the Predaceous Chub will continue to be monitored.
- 5.7. Whilst some differences between the original 2007 baseline surveys and the October 2013 monitoring surveys are evident, findings from water monitoring could be attributed to a range of factors including seasonal variations, and climatic conditions and/or the influence of tidal status at the time of survey. Taking this into account, the key Water Quality Objectives (WQOs) for River Monitoring Stations at Mui Wo River (EPD, 2011) have been included to provide a comparison with standard water quality goals applicable to the area (refer to **Table 3.12**).

- 5.8. The Environmental Protection Department (EPD) analyses and presents data from its annual water monitoring programme to express the level of compliance with the statutory WQOs including pH, Suspended Solids (SS), 5-day Biochemical Oxygen Demand (BOD<sub>5</sub>), and Dissolved Oxygen (DO). These WQOs specify the long-term water quality goals that the Government is to achieve and maintain for individual rivers in Hong Kong, including the Mui Wo River. As part of the programme five locations are sampled from the Mui Wo River, three of which are associated with the monitoring area for the drainage improvement works (MW1, MW2 and MW4). The objectives related to these sampling locations, have been used in this report. Water quality of the subject watercourses has met the WQOs during the survey.
- 5.9. Nitrogen (ammonia) has demonstrated a similar concentration at most monitoring stations compared to the last monitoring period, except at WE5. At WE5, concentration of nitrogen (ammonia) has increased double from 0.80 mg/L to 1.64 mg/L compared to the last monitoring period. No observable evidence of environmental changes such as odour, or discharge within the surveyed area, were recorded. Dissolved oxygen (DO) and conductivity were observed to increase at all monitoring stations compared to the last sampling record. Other monitoring parameters such as suspended solids, BOD<sub>5</sub> concentrations and pH value at all locations have demonstrated minimal change compared to the last sampling record.
- 5.10.

**Table 5.1** Key Observations/Comments and Recommendations Arising from the October 2013 Monitoring Period

Location	Mitigation Measure	Observations/Comments	Recommendations
PNH and LTT	Construction of a small fish ladder at the upstream end of the PNH	Vegetation has re-established in PNH4; however, the fish ladder is not currently overgrown or blocking water flow.	<p>The retention of native species, particularly at the edges of the river channel, during any future maintenance activities is recommended, to maintain existing habitat and minimize the re-colonization of exotic species.</p> <p>Some pits have been incorporated into the gabion banks, but do not appear to have been planted up. Planting of riparian vegetation, preferably with native species suggested in the EIA report Section 7.8.17 and Table 2.6 (e.g. <i>Albizia lebbek</i>, <i>Sterculia lanceolata</i>, <i>Cinnamomum camphora</i>, <i>Polyspora axillaris</i>, and <i>Rhaphiolepis indica</i>) is recommended.</p> <p>On-going weed management is recommended, as required, to maintain the open nature of the fish ladder.</p>
		The fish ladder does not meet the lip of the weir at the up-stream end of PNH4 due to a drop of approximately 30 cm. This could limit the overall function of the fish ladder for fish passage/movement up and downstream.	As per the current design of the fish ladder, the gap from the top of the fish ladder and the bottom of the weir is 30-40cm - presenting an obstacle to fish passage. Some improvement may be achieved by stacking

Location	Mitigation Measure	Observations/Comments	Recommendations
			additional boulders resembling that in PNH4 to form pools at the top of the fish ladder, which could facilitate fish movement.
	Re-establishment of aquatic / riparian communities	Predaceous Chub ( <i>Parazacco spilurus</i> ), which is of conservation importance, was recorded at PNH2, LTT1 and LTT5. Predaceous Chub was recorded at PNH4 and LTT5 in the December 2012 survey of post-construction monitoring. While the other fish species of conservation importance, Flagtail ( <i>Kuhlia marginata</i> ), was not recorded. Flagtail was recorded in the 2003-2004 EIA surveys.	The presence of species of conservation importance in both PNH and LTT including relative abundance will continue to be monitored.
LBC	Provision of suitable habitat compensation	Continued dominance by the exotic species, <i>Wedelia trilobata</i> , while limited marsh species were recorded.  Tree seedlings re-established at LBC2 and LBC3 that may hinder the re-establishment of a marsh habitat.	The regeneration of marsh species in the LBC is to be monitored.  Removal of tree seedling (species of <i>Acacia confusa</i> and <i>Sapium sebiferum</i> ) is suggested at LBC2 and LBC 3.
		The limited occurrence of typical marsh plant species (although this was also limiting in the RS) suggests that the water levels/availability within the channel may not be adequate to sustain a marsh habitat.	On-going monitoring of water levels and species composition within the channel are required. Further assessment should take into account the timing of the surveys (wet/dry season).

## 6. REFERENCES

AFCD (2013). Hong Kong Biodiversity Database. Available at <http://www.afcd.gov.hk/english/conservation/hkbiodiversity/database/resultlist.asp?lang=en> Accessed on 8 November 2013.

Carey, G. J., Chalmers, M. L., Diskin, D. A., Kennerley, P. R., Leader, P. J., Leven, M. R., Lewthwaite, R. W., Melville, D. S., Turnbull, M., and Young, L. (2001). *The Avifauna of Hong Kong*. Hong Kong Bird Watching Society, Hong Kong.

Chan, A., Cheung, J., Sze, P., Wong, A., Wong, E. and Yau, E. (2011). A Review of the Local Restrictedness of Hong Kong Butterflies. *Hong Kong Biodiversity Newsletter* 21: 1-6. Agriculture, Fisheries and Conservation Department, Hong Kong Special Administrative Region.

Chan, K.K. Benny and Caley, Kevin J. (2003). *Hong Kong Field Guides – Sandy Shores*. The Department of Ecology and Biodiversity, The University of Hong Kong.

Chan, S. K.-F., Cheung, K.-S., Ho, C.-Y., Lam, F.-N., Tang, W.-S., Lau, M. W.-N., Bogadek, A. (2005). *A Field Guide to the Amphibians of Hong Kong*. Agriculture, Fisheries and Conservation Department, Friends of the Country Parks and Cosmos Books Ltd.

Chan, S. K.-F., Cheung, K. S., Ho, C. Y., Lam, F. N., Tang, W. S., Tse, M. L. (2006). *A Field Guide to the Venomous Land Snakes of Hong Kong*. Agriculture, Fisheries and Conservation Department, Friends of the Country Parks and Cosmos Books Ltd.

Chan, S.K.F., Chan, A.S.W., Cheung, K.S., Ho, C.Y. Ng, C.K.Y. Tang, W.S. (2009). *The Skinks of Hong Kong*. *Hong Kong Biodiversity Newsletter: Issue 17*

Corlett, R., Xing, W. F., Ng, C. S., Chau, K. C. L., & Wong, M. Y. L. (2000). *Hong Kong Vascular Plants: Distribution and Status*. *Memoirs of the Hong Kong Natural History Society*, 23, 1-157.

Dudgeon (2003). *Hong Kong Field Guides – Hillstreams*. The Department of Ecology and Biodiversity, The University of Hong Kong.

Environmental Protection Department, 2005. *Wetland Restoration in Country Parks*. Advisory Council on the Environment Nature Conservation Subcommittee. Committee Paper NCSC 2/05

Environmental Protection Department (2011). *River Water Quality in Hong Kong in 2011*. The Government of the Hong Kong Special Administrative Region.

Fellowes, J. R., Lau, M. W., Dudgeon, D., Reels, G. T., Ades, G. W., & Carey, G. J. (2002). *Wild Animals to Watch: Terrestrial and Freshwater Fauna of Conservation Concern in Hong Kong*. *Memoirs of the Hong Kong Natural History Society*, 25, 123-159.

Hong Kong Herbarium and South China Botanical Garden (2007). *Flora of Hong Kong*. Volume 1. Agriculture, Fisheries and Conservation Department, Government of Hong Kong Special Administrative Region.

Hong Kong Herbarium and South China Botanical Garden (2008). Flora of Hong Kong. Volume 2. Agriculture, Fisheries and Conservation Department, Government of Hong Kong Special Administrative Region.

Hong Kong Herbarium and South China Botanical Garden (2009). Flora of Hong Kong. Volume 3. Agriculture, Fisheries and Conservation Department, Government of Hong Kong Special Administrative Region.

Hong Kong Herbarium and South China Botanical Garden (2011). Flora of Hong Kong. Volume 4. Agriculture, Fisheries and Conservation Department, Government of Hong Kong Special Administrative Region.

Hong Kong Herbarium (2012). Check List of Hong Kong Plants 2012. Agriculture, Fisheries and Conservation Department, HKSAR Government.

Hu, Q.M, Wu, T.L., Xia, N.H., Xing F.W., Patrick C.C.L., Yip, K.W. (2003). Rare and Precious Plants of Hong Kong. Agriculture, Fisheries and Conservation Department, Hong Kong Special Administrative Government.

IUCN (2013). IUCN Red List of Threatened Species. Version 2013.1. Available at [www.iucnredlist.org](http://www.iucnredlist.org). Accessed on 12 July 2013.

Lee, V. L. F., Lam, S. K. S., Ng, F. K. Y., Chan, T. K. T. and Young, M. L. C. (2004). Field Guide to the Freshwater Fish of Hong Kong. Agriculture, Fisheries and Conservation Department, Friends of the Country Parks and Cosmos Books Ltd. Hong Kong.

Lo, P.Y.F. (2005). Hong Kong Butterflies, 2nd edition. Agriculture, Fisheries and Conservation Department.

Karsen, S. J., Lau, M. W. N. and Bogadek, A. (1998). Hong Kong Amphibians and Reptiles. Urban Council, Hong Kong.

Shek, C.T. (2006). A Field Guide to the Terrestrial Mammals of Hong Kong. Agriculture, Fisheries and Conservation Department, Hong Kong.

Tam, T.W., Leung, K.K., Kwan, B.S.P., Wu, K.K.Y., Tang, S.S.H., So, I.W.Y., Cheng, J.C.Y., Yuen, E.F.M., Tsang, Y.M., and Hui, W.L. (2011). The Hong Kong Dragonflies. AFCD, Friends of Country Park and Cosmos Books Ltd. Hong Kong.

Viney, C., Phillips, K. and Ying, L.C. (2005). The birds of Hong Kong. Information Service Department.

William, G.A. (2003). Hong Kong Field Guides: Rocky Shore. The Department of Ecology and Biodiversity, The Hong Kong University of Hong Kong, Hong Kong.

Zheng, G. M. and Wang, Q. S. (1998). China Red Data Book of Endangered Animals: Aves. Science Press, Beijing.

## Appendix 1. Calibration certificate of the instruments (pH meter and multi-meter)



ALS Technichem (HK) Pty Ltd

### REPORT OF EQUIPMENT PERFORMANCE CHECK/CALIBRATION

**CONTACT:** MR MIKE SHEK  
**CLIENT:** AECOM ASIA COMPANY LIMITED  
**ADDRESS:** 11/F, TOWER 2, GRAND CENTRAL PLAZA,  
138 SHATIN RURAL COMMITTEE ROAD,  
SHATIN, N.T.,  
HONG KONG.

**WORK ORDER:** HK1321115  
**LABORATORY:** HONG KONG  
**DATE RECEIVED:** 06/08/2013  
**DATE OF ISSUE:** 09/08/2013

**PROJECT:** --

#### COMMENTS

It is certified that the item under calibration/checking has been calibrated/checked by corresponding calibrated equipment in the laboratory.  
Maximum Tolerance and calibration frequency stated in the report, unless otherwise stated, the internal acceptance criteria of ALS will be followed.

**Scope of Test:** Conductivity, Dissolved Oxygen, Salinity and Temperature  
**Equipment Type:** Multimeter  
**Brand Name:** YSI  
**Model No.:** Professional Plus  
**Serial No.:** 12M100515  
**Equipment No.:** W.040.01  
**Date of Calibration:** 08 August, 2013

#### NOTES

This is the Final Report and supersedes any preliminary report with this batch number.  
Results apply to sample(s) as submitted. All pages of this report have been checked and approved for release.

#### ISSUING LABORATORY: HONG KONG

##### **Address**

ALS Technichem (HK) Pty Ltd  
11/F Chung Shun Knitting Centre  
1-3 Wing Yip Street  
Kwai Chung  
HONG KONG

**Phone:** 852-2610 1044  
**Fax:** 852-2610 2021  
**Email:** [hongkong@alsglobal.com](mailto:hongkong@alsglobal.com)

  
Mr. Fung Lim Chee, Richard  
General Manager  
Greater China & Hong Kong

*This report may not be reproduced except with prior written approval from ALS Technichem (HK) Pty Ltd.*

Page 1 of 2

ADDRESS 11/F, Chung Shun Knitting Centre, 1-3 Wing Yip Street, Kwai Chung, N.T., Hong Kong | PHONE +852 2610 1044 | FAX +852 2610 2021  
ALS TECHNICHEM (HK) PTY LTD An ALS Limited Company

Life Sciences

[www.alsglobal.com](http://www.alsglobal.com)

RIGHT SOLUTIONS RIGHT PARTNER

## REPORT OF EQUIPMENT PERFORMANCE CHECK/CALIBRATION

**Work Order:** HK1321115  
**Date of Issue:** 09/08/2013  
**Client:** AECOM ASIA COMPANY LIMITED



**Description:** Multimeter  
**Brand Name:** YSI  
**Model No.:** Professional Plus  
**Serial No.:** 12M100515  
**Equipment No.:** W.040.01  
**Date of Calibration:** 08 August, 2013      **Date of next Calibration:** 08 November, 2013

**Parameters:**

**Conductivity**

**Method Ref: APHA (21st edition), 2510B**

Expected Reading (uS/cm)	Displayed Reading (uS/cm)	Tolerance (%)
146.9	143.4	-2.4
6667	6362	-4.6
12890	11960	-7.2
58670	56690	-3.4
Tolerance Limit (±%)		10.0

**Dissolved Oxygen**

**Method Ref: APHA (21st edition), 4500O: G**

Expected Reading (mg/L)	Displayed Reading (mg/L)	Tolerance (mg/L)
6.21	6.02	-0.19
7.26	7.28	0.02
8.40	8.26	-0.14
Tolerance Limit (±mg/L)		0.20

**Temperature**

**Method Ref: Section 6 of International Accreditation New Zealand Technical Guide No. 3 Second edition March 2008: Working Thermometer Calibration Procedure.**

Expected Reading (°C)	Displayed Reading (°C)	Tolerance (°C)
13.0	12.6	-0.4
21.0	21.1	0.1
38.0	38.7	0.7
Tolerance Limit (±°C)		2.0

**Salinity**

**Method Ref: APHA (21st edition), 2520B**

Expected Reading (ppt)	Displayed Reading (ppt)	Tolerance (%)
0	0.00	-
10	9.88	-1.2
20	21.09	5.5
30	30.67	2.2
Tolerance Limit (±%)		10.0

Remark: "Displayed Reading" presents the figures shown on item under calibration / checking regardless of equipment precision or significant figures.

  
 Mr. Fung Lim Chee, Richard  
 General Manager  
 Greater China & Hong Kong



## ALS Technichem (HK) Pty Ltd

### REPORT OF EQUIPMENT PERFORMANCE CHECK/CALIBRATION

**CONTACT:** MR MIKE SHEK  
**CLIENT:** AECOM ASIA COMPANY LIMITED  
**ADDRESS:** 11/F, TOWER 2, GRAND CENTRAL PLAZA,  
138 SHATIN RURAL COMMITTEE ROAD,  
SHATIN, N.T.,  
HONG KONG.

**WORK ORDER:** HK1328190  
**LABORATORY:** HONG KONG  
**DATE RECEIVED:** 11/10/2013  
**DATE OF ISSUE:** 21/10/2013

**PROJECT:** --

#### COMMENTS

It is certified that the item under calibration/checking has been calibrated/checked by corresponding calibrated equipment in the laboratory.  
Maximum Tolerance and calibration frequency stated in the report, unless otherwise stated, the internal acceptance criteria of ALS will be followed.

Scope of Test: pH and Temperature  
Equipment Type: pH Meter  
Brand Name: WTW  
Model No.: pH 3210  
Serial No.: 12340605  
Equipment No.: W.039.08  
Date of Calibration: 21 October, 2013

#### NOTES

This is the Final Report and supersedes any preliminary report with this batch number.  
Results apply to sample(s) as submitted. All pages of this report have been checked and approved for release.

#### ISSUING LABORATORY: HONG KONG

##### Address

ALS Technichem (HK) Pty Ltd  
11/F Chung Shun Knitting Centre  
1-3 Wing Yip Street  
Kwai Chung  
HONG KONG

**Phone:** 852-2610 1044  
**Fax:** 852-2610 2021  
**Email:** [hongkong@alsglobal.com](mailto:hongkong@alsglobal.com)

  
Mr. Fung Lim Chee, Richard  
General Manager -  
Greater China & Hong Kong

*This report may not be reproduced except with prior written approval from ALS Technichem (HK) Pty Ltd.*

Page 1 of 2

ADDRESS 11/F, Chung Shun Knitting Centre, 1-3 Wing Yip Street, Kwai Chung, N.T., Hong Kong | PHONE +852 2610 1044 | FAX +852 2610 2021  
ALS TECHNICHEM (HK) PTY LTD An ALS Limited Company

Life Sciences

[www.alsglobal.com](http://www.alsglobal.com)

RIGHT SOLUTIONS RIGHT PARTNER



## REPORT OF EQUIPMENT PERFORMANCE CHECK/CALIBRATION

Work Order: HK1328190  
Date of Issue: 21/10/2013  
Client: AECOM ASIA COMPANY LIMITED



Equipment Type: pH Meter  
Brand Name: WTW  
Model No.: pH 3210  
Serial No.: 12340605  
Equipment No.: W.039.08  
Date of Calibration: 21 October, 2013      Date of next Calibration: 21 January, 2014

### Parameters:

#### pH Value

Method Ref: APHA 21st Ed. 4500H:B

Expected Reading (pH Unit)	Displayed Reading (pH Unit)	Tolerance (pH unit)
4.0	4.046	0.05
7.0	7.123	0.12
10.0	10.018	0.02
Tolerance Limit ( $\pm$ pH unit)		0.20

#### Temperature

Method Ref: Section 6 of International Accreditation New Zealand Technical Guide No. 3 Second edition March 2008: Working Thermometer Calibration Procedure.

Expected Reading ( $^{\circ}$ C)	Displayed Reading ( $^{\circ}$ C)	Tolerance ( $^{\circ}$ C)
10.0	9.8	-0.2
25.0	24.2	-0.8
38.5	37.8	-0.7
Tolerance Limit ( $\pm$ $^{\circ}$ C)		2.0

Remark: "Displayed Reading" presents the figures shown on item under calibration / checking regardless of equipment precision or significant figures.

Appendix 2a: Plant Species Recorded in Pak Ngan Heung River and Luk Tei Tong River

Scientific Name	Growth Form	Native / Exotic to Hong Kong	Distribution in Hong Kong	PNH1	PNH2	PNH3	PNH4	LTT1	LTT2	LTT3	LTT4	LTT5
<i>Acanthus ilicifolius</i>	shrub	native	common						++			
<i>Ageratum houstonianum</i>	herb	exotic	common			+						
<i>Bidens alba</i>	herb	exotic	very common			+	++			++	+	+
<i>Canavalia maritima</i>	climber	native	common						+	+	+	+
<i>Cleome burmannii</i>	herb	exotic	-			+						
<i>Colocasia esculenta</i>	herb	native	-			+	+					
<i>Commelina diffusa</i>	herb	native	common			+	+					
<i>Cyperus</i> sp.	herb	-	-			+						
<i>Ficus hispida</i>	tree	native	very common		+	+						
<i>Ficus variegata</i>	shrub	native	common				+					
<i>Indigofera hirsuta</i>	shrub	native	common									+
<i>Ipomoea cairica</i>	climber	exotic	very common								+	+
<i>Itea chinensis</i>	shrub or small tree	native	very common						+			
<i>Kandelia obovata</i>	shrub or small tree	native	common		+			+	+	+		+
<i>Ludwigia octovalvis</i>	perennial herb	native	common			+	+					
<i>Mikania micrantha</i>	climber	exotic	very common			+++	+++		+	+	+	+
<i>Mimosa diplotricha</i>	herb	exotic	rare							+		
<i>Miscanthus sinensis</i>	perennial herb	native	very common							+		
<i>Neyraudia reynaudiana</i>	herb	native	very common								+	
<i>Polygonum barbatum</i>	herb	native	common			+	++					
<i>Praxelis clematidea</i>	perennial herb	exotic	very common			+	+					
<i>Rhus succedanea</i>	tree	native	common				+					
<i>Urena lobata</i>	shrub	native	common			+						
<i>Wedelia trilobata</i>	perennial herb	exotic	common	+			++		+			+

Note:

Code for Abundance: +++=abundant; ++=frequent; +=occasional

**Appendix 2b: Plant Species Recorded in Luk Tei Tong Bypass Channel and Reference Site**

LLT Bypass Channel (LBC)

Scientific Name	Growth Form	Native / Exotic to Hong Kong	Distribution in Hong Kong	LBC1	LBC2	LBC3	LBC4	LBC5	Average
<b>Species recorded in the quadrats along the transects</b>				<b>Average Percentage Cover</b>					
<i>Colocasia esculenta</i>	herb	native	-	0.00	0.00	0.00	0.00	0.01	0.00
<i>Crotalaria pallida</i>	herb	exotic	common	0.00	0.00	0.01	0.05	0.01	0.01
<i>Cyclosorus interruptus</i>	herb	native	common	0.00	0.00	0.04	0.01	0.00	0.01
<i>Cyperus</i> sp.	-	-	-	0.00	0.01	0.00	0.01	0.07	0.02
<i>Cyperus alternifolius</i> subsp. <i>flabelliformis</i>	herb	exotic	n/a	0.00	0.00	0.00	0.00	0.00	0.00
<i>Fimbristylis sieboldii</i>	herb	native	common	0.60	0.00	0.00	0.00	0.00	0.12
<i>Hedychium coronarium</i>	shrub	exotic	-	0.00	0.00	0.00	0.00	0.05	0.01
<i>Ipomoea cairica</i>	climber	exotic	very common	0.17	0.00	0.00	0.00	0.00	0.03
<i>Microstegium ciliatum</i>	perennial procumbent herb	native	very common	0.00	0.14	0.21	0.12	0.17	0.13
<i>Mikania micrantha</i>	climber	exotic	very common	0.00	0.00	0.00	0.00	0.01	0.00
<i>Mimosa diplotricha</i>	herb	exotic	rare	0.00	0.27	0.00	0.18	0.09	0.11
<i>Panicum maximum</i>	herb	exotic	very common	0.00	0.00	0.00	0.00	0.10	0.02
<i>Paspalum conjugatum</i>	perennial herb	exotic	common	0.00	0.08	0.04	0.00	0.00	0.02
<i>Praxelis clematidea</i>	perennial herb	exotic	very common	0.00	0.00	0.04	0.00	0.08	0.02
<i>Ruellia coerulea</i>	herb	exotic	-	0.23	0.00	0.00	0.00	0.00	0.05
<i>Solanum torvum</i>	shrub	exotic	common	0.00	0.00	0.01	0.00	0.00	0.00
<i>Wedelia trilobata</i>	perennial herb	exotic	common	0.00	0.50	0.65	0.63	0.41	0.44
<b>Species recorded during the walk-through survey</b>				<b>Occurrence of the Species</b>					
<i>Acacia confusa</i>	tree	exotic	-		+				
<i>Adiantum capillus-veneris</i>	herb	native	common		+				
<i>Apluda mutica</i>	herb	native	very common		+				
<i>Bidens alba</i>	herb	exotic	very common		+	+	+		
<i>Celtis sinensis</i>	tree	native	common			+	+		
<i>Colocasia esculenta</i>	herb	native	-		+	+	+	+	
<i>Commelina diffusa</i>	herb	native	common		+				
<i>Crotalaria pallida</i>	herb	exotic	common		+	+	+	+	
<i>Cyclosorus interruptus</i>	herb	native	common		+	+	+	+	
<i>Cynodon dactylon</i>	perennial herb	native	very common				+		
<i>Cyperus</i> sp.	-	-	-		+		+	+	
<i>Cyperus alternifolius</i> subsp. <i>flabelliformis</i>	herb	exotic	-	+					
<i>Fimbristylis sieboldii</i>	herb	native	common	+	+				
<i>Hedychium coronarium</i>	shrub	exotic	-		+		+	+	
<i>Ipomoea cairica</i>	climber	exotic	very common	+	+	+	+	+	
<i>Ipomoea triloba</i>	climber: twining herb	native	common			+		+	
<i>Kandelia obovata</i>	shrub or small tree	native	common	+					
<i>Lindernia</i> sp.	herb	native	-		+				
<i>Lophatherum gracile</i>	herb	native	common			+			

Note:

Code: +=occurrence of the species

**Appendix 2b: Plant Species Recorded in Luk Tei Tong Bypass Channel and Reference Site**

LLT Bypass Channel (LBC)

Scientific Name	Growth Form	Native / Exotic to Hong Kong	Distribution in Hong Kong	LBC1	LBC2	LBC3	LBC4	LBC5
<b>Other species recorded during the walk-through survey (Continue)</b>				<b>Occurrence of the Species</b>				
<i>Ludwigia octovalvis</i>	perennial herb	native	common		+		+	+
<i>Macaranga tanarius</i>	tree	native	common			+		
<i>Microstegium ciliatum</i>	perennial procumbent herb	native	very common		+	+	+	+
<i>Mikania micrantha</i>	climber	exotic	very common		+	+	+	+
<i>Mimosa diplotricha</i>	herb	exotic	rare			+	+	
<i>Panicum maximum</i>	herb	exotic	very common		+	+		+
<i>Paspalum conjugatum</i>	perennial herb	exotic	common		+	+	+	+
<i>Paspalum orbiculare</i>	herb	native	-	+	+			
<i>Polygonum barbatum</i>	herb	native	common			+	+	+
<i>Polygonum chinense</i>	herb	native	very common					+
<i>Polygonum</i> sp.	herb	n/a	n/a		+			
<i>Praxelis clematidea</i>	perennial herb	exotic	very common		+	+		+
<i>Pycneus polystachyus</i>	herb	native	common	+	+	+		+
<i>Ruellia coerulea</i>	herb	exotic	-	+				
<i>Sapium sebiferum</i>	tree	native	common		+	+		
<i>Solanum torvum</i>	shrub	exotic	common			+		
<i>Urena lobata</i>	shrub	native	common		+	+	+	
<i>Wedelia trilobata</i>	perennial herb	exotic	common		+	+		+

Note:

Code: +=occurrence of the species

**Appendix 2b: Plant Species Recorded in Luk Tei Tong Bypass Channel and Reference Site**

Reference Site (RS)

Scientific Name	Growth Form	Native / Exotic to Hong Kong	Distribution in Hong Kong	RS1	RS2	RS3	RS4	RS5	Average
<b>Species recorded in the quadrats along the transects</b>				<b>Average Percentage Cover</b>					
<i>Bidens alba</i>	herb	exotic	very common	0.00	0.05	0.02	0.00	0.22	0.06
<i>Colocasia esculenta</i>	herb	native	-	0.00	0.00	0.00	0.01	0.00	0.00
<i>Crotalaria pallida</i>	herb	exotic	common	0.02	0.00	0.01	0.00	0.00	0.01
<i>Hedychium coronarium</i>	shrub	exotic	-	0.00	0.00	0.01	0.00	0.00	0.00
<i>Imperata koenigii</i>	perennial herb	native	very common	0.02	0.00	0.00	0.01	0.00	0.01
<i>Ipomoea cairica</i>	climber	exotic	very common	0.00	0.00	0.00	0.01	0.00	0.00
<i>Lophatherum gracile</i>	herb	native	common	0.05	0.01	0.07	0.07	0.00	0.04
<i>Mikania micrantha</i>	climber	exotic	very common	0.00	0.00	0.00	0.03	0.00	0.01
<i>Mimosa diplotricha</i>	herb	exotic	rare	0.00	0.01	0.03	0.00	0.00	0.01
<i>Mimosa pudica</i>	herb	exotic	very common	0.09	0.00	0.00	0.05	0.16	0.06
<i>Paspalum conjugatum</i>	perennial herb	exotic	common	0.01	0.00	0.00	0.00	0.02	0.01
<i>Pueraria phaseoloides</i>	climber	native	very common	0.00	0.01	0.00	0.00	0.00	0.00
<i>Urena lobata</i>	shrub	native	common	0.03	0.05	0.01	0.04	0.02	0.03
<i>Wedelia trilobata</i>	perennial herb	exotic	common	0.67	0.53	0.83	0.78	0.48	0.66
<b>Other species recorded during the walk-through survey</b>				<b>Occurrence of the Species</b>					
<i>Acacia confusa</i>	tree	exotic	-	+					
<i>Adiantum capillus-veneris</i>	herb	native	common				+		
<i>Allamanda cathartica</i>	climbing shrub	exotic	-			+			
<i>Aster subulatus</i>	herb	exotic	-		+				
<i>Bauhinia blakeana</i>	tree	native	common	+	+				
<i>Bidens alba</i>	herb	exotic	very common	+	+	+		+	
<i>Canna indica</i>	herb	exotic	-				+		
<i>Celosia argentea</i>	herb	native	very common	+				+	
<i>Celtis sinensis</i>	tree	native	common	+			+		
<i>Chamaecrista mimosoides</i>	subshrubby Herb	native	common	+	+		+		
<i>Colocasia esculenta</i>	herb	native	-		+		+		
<i>Conyza canadensis</i>	herb	exotic	very common	+				+	
<i>Crotalaria pallida</i>	herb	exotic	common	+		+			
<i>Ficus hispida</i>	tree	native	very common	+	+	+		+	
<i>Ficus variegata</i>	shrub	native	common		+				
<i>Hedychium coronarium</i>	shrub	exotic	-		+	+			
<i>Imperata koenigii</i>	perennial herb	native	very common	+			+	+	

Note:

Code: +=occurrence of the species

**Appendix 2b: Plant Species Recorded in Luk Tei Tong Bypass Channel and Reference Site**

Reference Site (RS)

Scientific Name	Growth Form	Native / Exotic to Hong Kong	Distribution in Hong Kong	RS1	RS2	RS3	RS4	RS5
<b>Other species recorded during the walk-through survey (Continue)</b>				<b>Occurrence of the Species</b>				
<i>Ipomoea cairica</i>	climber	exotic	very common	+	+	+		+
<i>Ipomoea pes-caprae</i>	perennial herb	native	common		+		+	+
<i>Leucaena leucocephala</i>	tree	exotic	common				+	
<i>Lophatherum gracile</i>	herb	native	common	+	+	+	+	
<i>Mallotus paniculatus</i>	tree	native	very common					+
<i>Mikania micrantha</i>	climber	exotic	very common		+	+	+	+
<i>Mimosa diplotricha</i>	herb	exotic	rare	+	+	+		+
<i>Mimosa pudica</i>	herb	exotic	very common	+			+	+
<i>Paspalum conjugatum</i>	perennial herb	exotic	common	+		+		+
<i>Praxelis clematidea</i>	perennial herb	exotic	very common	+	+			+
<i>Pueraria phaseoloides</i>	climber	native	very common	+	+	+	+	
<i>Pycneus polystachyus</i>	herb	native	common		+			
<i>Sageretia thea</i>	shrub	native	very common	+	+			
<i>Sapium sebiferum</i>	tree	native	common		+	+	+	
<i>Sesbania cannabina</i>	herb	exotic	common			+		
<i>Sida rhombifolia</i>	herb	native	common			+		
<i>Solanum torvum</i>	shrub	exotic	common			+		
<i>Urena lobata</i>	shrub	native	common	+	+	+	+	+
<i>Wedelia trilobata</i>	perennial herb	exotic	common	+	+	+	+	+

Note:

Code: +=occurrence of the species

**Appendix 3: Ecological Water Quality Monitoring Raw Data  
(October 2013)**

Date of Monitoring: 25 October 2013

Weather : Sunny

Monitoring Location	Suspended Solids (mg/L)	Nitrogen (Ammonia) (mg/L)	Nitrogen (Nitrate) (mg/L)	Reactive Phosphorous (mg/L)	5-day Biochemical Oxygen Demand (BOD <sup>5</sup> ) (mg/L)	Dissolved Oxygen (mg/L)	
						M1	M2
WE1	<2.0	0.12	0.10	0.01	<2.0	10.31	10.61
WE2	<2.0	0.11	0.10	0.02	<2.0	10.50	11.04
WE3	<2.0	0.05	0.09	0.02	<2.0	11.22	12.10
WE4	6.0	0.14	0.21	0.05	<2.0	9.12	10.46
WE5	3.0	1.64	0.06	0.19	<2.0	10.21	10.97
WE6	3.0	0.05	0.10	0.01	<2.0	12.87	11.20
WE7	No water - Not sampled						
WE8	No water - Not sampled						
WE9	No water - Not sampled						
WE10	No water - Not sampled						

Monitoring Location	Temperature (°C)		pH	Salinity (ppt)		Conductivity (µs/cm)		Water Flow (m/s)		Water Depth (cm)
	M1	M2		M1	M2	M1	M2	M1	M2	
WE1	21.3	21.3	7.0	0.01	0.01	31.7	30.1	0.03	0.04	36
WE2	21.4	21.5	7.2	0.03	0.02	335.4	334.6	0.02	0.01	15
WE3	22.1	22.1	7.4	0.20	0.20	423.0	419.7	0.05	0.04	17
WE4	25.6	25.5	7.8	7.46	7.55	13025.0	13152.0	0.07	0.07	24
WE5	24.9	24.9	7.6	0.73	0.82	1398.0	1654	0.03	0.03	15
WE6	22.4	22.4	7.9	0.01	0.01	30.9	30.6	0.02	0.02	28
WE7	No water - Not sampled									
WE8	No water - Not sampled									
WE9	No water - Not sampled									
WE10	No water - Not sampled									

Note:

Where more than one measurement was taken, the data is represented by Measurement M1 and M2.



PNH1 and PNH2



PNH3 and PNH4



LTT1



LTT2 and LTT3



LTT4 and LTT5



LBC1



LBC2 - LBC5



Post-Construction Ecological Monitoring  
of Drainage Improvement Works in Southern Lantau

SCALE

N.T.S.

DATE

Nov-13

**Representative Photographs taken during  
the Monitoring**

CHECK

McmillanSE

DRAWN

CHIKYY

JOB NO.

60278381

DRAWING No.

Appendix 4





RS1



RS2



RS3 and RS4



RS5



Post-Construction Ecological Monitoring  
of Drainage Improvement Works in Southern Lantau

SCALE

N.T.S.

DATE

Nov-13

**Representative Photographs taken during  
the Monitoring**

CHECK

McmillanSE

DRAWN

CHIKYY

JOB NO.

60278381

DRAWING No.

Appendix 4