


A map of Hong Kong showing the locations where horseshoe crabs, including *Tachypleus tridentatus* (○: living adults, ●: dead adults, ⊗: living juveniles) and *Carcinoscorpius rotundicauda* (□: living adults, ■: dead adults, ⊠: living juveniles), have been found during the study period from March 1995 to June 1998.



Locations:

- | | | |
|-----------------------|-----------------------|-----------------------|
| 1 Tsim Bei Tsui | 2 Lau Fau Shan | 3 Sheung Pak Nai |
| 4 Ha Pak Nai | 5 Nim Wan | 6 Lung Kwu Sheung Tan |
| 7 Tuen Mun | 8 Sha Chau | 9 The Brothers |
| 10 Tung Chung Bay | 11 San Tau | 12 Sha Lo Wan |
| 13 Sham Wat Wan | 14 Tai O | 15 Yi O |
| 16 Shui Hau Wan | 17 Pui O Beach | 18 Shek Kwu Chau |
| 19 Peng Chau | 20 West Lamma Channel | 21 East Lamma Channel |
| 22 Tung O Wan | 23 Kau Pei Chau | 24 Waglan Island |
| 25 Silverstrand Beach | 26 Pak Sha Wan | 27 Sharp Island |

EXTRACTED FROM CHIU AND MORTON (1999)