

| Tree Group# | Relevant LRs | District | Botanical Name | Chinese Name | Estimated Total Number of Trees within this Tree Group | Estimated Number of Trees for this Species | Range of Trunk Diameter at DBH (mm) | Range of Height (m) | Range of Spread (m) | Overall Amenity Value (High/ Medium/ Low) | Overall Form (Good/ Fair/ Poor) | Overall Health (Good/ Fair/ Poor) | Overall Structural Condition (Good/ Fair/ Poor) | Overall Suitability for Transplanting (High/ Medium/ Low) | Overall Remark of Suitability for Transplanting | Overall Conservation Status (Normal/ Rare/ Endanger/ OVT/ Tree Register) | Overall Survival Rate after Transplant (High/ Medium/ Low) | Additional Remarks |
|-----------------------------------|---------------------|----------|---------------------------|--------------|--|--|-------------------------------------|---------------------|---------------------|---|---------------------------------|-----------------------------------|---|---|---|--|--|--------------------|
| J/O CWBR | | | | | | | | | | | | | | | | | | |
| TG1 | LR3.1, LR9.1, LR9.2 | J/O CWBR | Acacia confusa | 台灣相思 | 531 | 75 | 100-620 | 5-16 | 4-14 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Acacia dealbata | 銀荊 | | 13 | 100-250 | 4-5 | 3-4 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | on slope |
| | | | Aporosa dioica | 銀柴 | | 28 | 100-250 | 6-8 | 4-5 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Aquilaria sinensis | 土沉香 | | 1 | 130 | 8 | 5 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Protected under Cap 586; Rare and Precious Plants of Hong Kong (Near Threatened); China Plant Red Data Book (Vulnerable) | Low | on slope |
| | | | Bridelia tomentosa | 土蜜樹 | | 2 | 120 | 5-6 | 3-4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Casuarina equisetifolia | 木麻黃 | | 3 | 200-300 | 12-16 | 8-9 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | on slope |
| | | | Celtis sinensis | 朴樹 | | 35 | 100-350 | 4-16 | 4-11 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Cinnamomum burmannii | 陰香 | | 3 | 100-200 | 8 | 5-6 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | on slope |
| | | | Cinnamomum camphora | 樟 | | 13 | 330-400 | 12-16 | 7-9 | Low | Poor | Fair | Poor | Low | a, b, c, d, f, h | Normal | Low | on slope |
| | | | Cleistocalyx nervosum | 水翁 | | 1 | 200 | 10 | 6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Cratoxylum cochinchinense | 黃牛木 | | 15 | 120-170 | 4-8 | 3-5 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Dimocarpus longan | 龍眼 | | 3 | 100-170 | 7-9 | 4-5 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Duranta erecta | 假連翹 | | 2 | 120-150 | 2-5 | 4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Eriobotrya japonica | 枇杷 | | 1 | 105 | 4 | 4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Ficus hispida | 對葉榕 | | 37 | 100-400 | 4-11 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | on slope |
| | | | Ficus variegata | 青果榕 | | 10 | 220 | 6 | 5 | Low | Poor | Poor | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Ficus virens | 大葉榕 | | 1 | 235 | 9 | 6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Garcinia oblongifolia | 嶺南山竹子 | | 1 | 115 | 8 | 4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Leucaena leucocephala | 銀合歡 | | 17 | 140-290 | 9-12 | 6 | Low | Poor | Fair | Poor | Low | a, b, c, f, g, h | Normal | Low | on slope |
| | | | Ligustrum sinense | 山指甲 | | 7 | 120-230 | 3-8 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | on slope |
| Litsea glutinosa | 潺槁樹 | 45 | 100-250 | 4-8 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope | | | | |
| Macaranga tanarius var. tomentosa | 血桐 | 15 | 100-260 | 4-9 | 4-8 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope | | | | |
| Machilus breviflora | 短序潤楠 | 15 | 100-320 | 6-16 | 4-7 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope | | | | |
| Machilus chekiangensis | 浙江潤楠 | 18 | 110-300 | 7-12 | 4-8 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope | | | | |
| Machilus pauhoi | 刨花潤楠 | 35 | 100-350 | 6-16 | 4-9 | Low | Poor | Fair | Poor | Low | a, b, c, d, f, h | Normal | Low | on slope | | | | |

| Tree Group# | Relevant LRs | District | Botanical Name | Chinese Name | Estimated Total Number of Trees within this Tree Group | Estimated Number of Trees for this Species | Range of Trunk Diameter at DBH (mm) | Range of Height (m) | Range of Spread (m) | Overall Amenity Value (High/ Medium/ Low) | Overall Form (Good/ Fair/ Poor) | Overall Health (Good/ Fair/ Poor) | Overall Structural Condition (Good/ Fair/ Poor) | Overall Suitability for Transplanting (High/ Medium/ Low) | Overall Remark of Suitability for Transplanting | Overall Conservation Status (Normal/ Rare/ Endanger/ OVT/ Tree Register) | Overall Survival Rate after Transplant (High/ Medium/ Low) | Additional Remarks |
|-------------|--------------|----------|------------------------|--------------|--|--|-------------------------------------|---------------------|---------------------|---|---------------------------------|-----------------------------------|---|---|---|--|--|------------------------|
| | | | Mallotus paniculatus | 白楸 | | 25 | 100-210 | 5-11 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Mangifera indica | 芒果 | | 1 | 185 | 9 | 6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Microcos nervosa | 破布葉 | | 4 | 110-190 | 5-10 | 3-5 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Phyllanthus emblica | 餘甘子 | | 3 | 115-175 | 6-9 | 4-5 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Prunus phaeosticta | 腺葉桂櫻 | | 1 | 160 | 7 | 4 | Low | Poor | Poor | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Rhus succedanea | 野漆樹 | | 13 | 100-150 | 5-7 | 3-4 | Low | Poor | Poor | Poor | Low | a, c, f, h | Normal | Low | on slope |
| | | | Sapium sebiferum | 烏桕 | | 5 | 150-450 | 6-10 | 5-8 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Schefflera heptaphylla | 鵝掌柴 | | 25 | 100-220 | 4-10 | 4-6 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | on slope |
| | | | Schima superba | 木荷 | | 12 | 120-140 | 5-10 | 4-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Sterculia lanceolata | 假蒺藜 | | 35 | 110-300 | 5-12 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Syzygium jambos | 蒲桃 | | 1 | 590 | 12 | 12 | Low | Poor | Fair | Poor | Low | a, b, c, d, f, h | Normal | Low | on slope |
| | | | Syzygium levinei | 山蒲桃 | | 2 | 100-135 | 6 | 4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Tetradium glabrifolium | 棟葉吳茱萸 | | 3 | 100-130 | 6-8 | 3-4 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | on slope |
| | | | Viburnum odoratissimum | 珊瑚樹 | | 4 | 150-300 | 6-8 | 4-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Zanthoxylum avicennae | 筍櫨花椒 | | 1 | 170 | 8 | 4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Acronychia pedunculata | 山油柑 | | 10 | 100-150 | 2-5 | 2-3 | Low | Poor | Fair | Fair | Low | a, c, f, h | Normal | Low | on slope, shallow soil |
| | | | Celtis sinensis | 朴樹 | | 12 | 100-230 | 4-7 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h, i | Normal | Low | on slope, shallow soil |
| | | | Cinnamomum camphora | 樟 | | 10 | 100-230 | 4-7 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h, i | Normal | Low | on slope, shallow soil |
| | | | Gordonia axillaris | 大頭茶 | | 15 | 100-120 | 3-6 | 1-3 | Low | Poor | Fair | Fair | Low | a, c, f, h | Normal | Low | on slope, shallow soil |
| | | | Litsea glutinosa | 潺槁樹 | | 15 | 120-150 | 2-5 | 2-4 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | on slope, shallow soil |
| | | | Mallotus paniculatus | 白楸 | 117 | 20 | 100-120 | 3-7 | 2-4 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | on slope, shallow soil |
| | | | Pinus massoniana | 馬尾松 | | 5 | 100-120 | 3-9 | 2-4 | Low | Poor | Fair | Fair | Low | a, b, c, f, h | Normal | Low | on slope, shallow soil |
| | | | Reevesia thyrsoidea | 梭羅樹 | | 5 | 100-130 | 3-6 | 3-4 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | on slope, shallow soil |
| | | | Schefflera heptaphylla | 鵝掌柴(鴨腳木) | | 15 | 100-120 | 2-7 | 3-4 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | on slope, shallow soil |
| | | | Schima superba | 木荷 | | 10 | 100-130 | 3-6 | 3-4 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | on slope, shallow soil |
| | | | Acacia confusa | 台灣相思 | | 60 | 100-500 | 5-16 | 4-14 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Acacia dealbata | 銀荊 | | 5 | 100-250 | 4-5 | 3-4 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | on slope |

| Tree Group# | Relevant LRs | District | Botanical Name | Chinese Name | Estimated Total Number of Trees within this Tree Group | Estimated Number of Trees for this Species | Range of Trunk Diameter at DBH (mm) | Range of Height (m) | Range of Spread (m) | Overall Amenity Value (High/ Medium/ Low) | Overall Form (Good/ Fair/ Poor) | Overall Health (Good/ Fair/ Poor) | Overall Structural Condition (Good/ Fair/ Poor) | Overall Suitability for Transplanting (High/ Medium/ Low) | Overall Remark of Suitability for Transplanting | Overall Conservation Status (Normal/ Rare/ Endanger/ OVT/ Tree Register) | Overall Survival Rate after Transplant (High/ Medium/ Low) | Additional Remarks |
|------------------------|--------------|----------|-----------------------------------|--------------|--|--|-------------------------------------|---------------------|---------------------|---|---------------------------------|-----------------------------------|---|---|---|--|--|--------------------|
| T03 | LR3.2, LR9.1 | J/O CWBR | Aporosa dioica | 銀柴 | 320 | 12 | 100-250 | 6-8 | 4-5 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Bridelia tomentosa | 土蜜樹 | | 4 | 100-120 | 5-6 | 3-4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Casuarina equisetifolia | 木麻黃 | | 4 | 200-400 | 12-16 | 8-9 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | on slope |
| | | | Cinnamomum camphora | 樟 | | 3 | 300-400 | 12-16 | 7-9 | Low | Poor | Fair | Poor | Low | a, b, c, d, f, h | Normal | Low | on slope |
| | | | Cratoxylum cochinchinense | 黃牛木 | | 15 | 100-150 | 4-8 | 3-5 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Dimocarpus longan | 龍眼 | | 7 | 100-150 | 7-9 | 4-5 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Duranta erecta | 假連翹 | | 2 | 100-120 | 2-5 | 3-4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Ficus hispida | 對菜榕 | | 17 | 100-500 | 4-11 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | on slope |
| | | | Ficus variegata | 青果榕 | | 5 | 150-250 | 6-8 | 3-5 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Leucaena leucocephala | 銀合歡 | | 10 | 150-300 | 9-12 | 4-7 | Low | Poor | Fair | Poor | Low | a, b, c, f, g, h | Normal | Low | on slope |
| | | | Ligustrum sinense | 山指甲 | | 8 | 100-150 | 3-8 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | on slope |
| | | | Litsea glutinosa | 潺槁樹 | | 22 | 100-250 | 4-8 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Macaranga tanarius var. tomentosa | 血桐 | | 18 | 100-250 | 4-9 | 4-8 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Machilus spp. | 潤楠屬 | | 25 | 100-320 | 6-16 | 4-7 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Mallotus paniculatus | 白楸 | | 10 | 100-210 | 5-11 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Microcos nervosa | 破布葉 | | 14 | 100-200 | 5-10 | 3-5 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Phyllanthus emblica | 餘甘子 | | 11 | 100-150 | 4-6 | 3-5 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Rhus succedanea | 野漆樹 | | 13 | 100-150 | 5-7 | 3-4 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | on slope |
| | | | Sapium sebiferum | 烏桕 | | 15 | 150-450 | 6-10 | 5-8 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Schefflera heptaphylla | 鵝掌柴 | | 13 | 100-250 | 4-10 | 4-6 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | on slope |
| Schima superba | 木荷 | 12 | 120-140 | 5-10 | 4-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope | | | | |
| Viburnum odoratissimum | 珊瑚樹 | 15 | 150-300 | 6-8 | 4-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope | | | | |
| | | | Casuarina equisetifolia | 木麻黃 | | 2 | 150-300 | 8-13 | 8-10 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | at slope, leaning |
| | | | Acacia confusa | 台灣相思 | | 14 | 100-550 | 6-16 | 4-10 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, leaning |
| | | | Acacia dealbata | 銀荊 | | 3 | 100-250 | 4-5 | 3-4 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | on slope |
| | | | Bridelia tomentosa | 土蜜樹 | | 1 | 120 | 5 | 4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Celtis sinensis | 朴樹 | | 14 | 120-350 | 5-16 | 3-11 | Low | Poor | Fair | Poor | Low | b, c, h | Normal | Low | on slope |

| Tree Group# | Relevant LRs | District | Botanical Name | Chinese Name | Estimated Total Number of Trees within this Tree Group | Estimated Number of Trees for this Species | Range of Trunk Diameter at DBH (mm) | Range of Height (m) | Range of Spread (m) | Overall Amenity Value (High/ Medium/ Low) | Overall Form (Good/ Fair/ Poor) | Overall Health (Good/ Fair/ Poor) | Overall Structural Condition (Good/ Fair/ Poor) | Overall Suitability for Transplanting (High/ Medium/ Low) | Overall Remark of Suitability for Transplanting | Overall Conservation Status (Normal/ Rare/ Endanger/ OVT/ Tree Register) | Overall Survival Rate after Transplant (High/ Medium/ Low) | Additional Remarks |
|-----------------|--------------------------|----------|-----------------------------------|--------------|--|--|-------------------------------------|---------------------|---------------------|---|---------------------------------|-----------------------------------|---|---|---|--|--|------------------------|
| TG4 | LR3.3, LR4, LR8.1, LR9.1 | J/O CWBR | Duranta erecta | 假連翹 | 67 | 2 | 120-150 | 2-5 | 4 | Low | Poor | Fair | Poor | | | Normal | Low | on slope |
| | | | Eriobotrya japonica | 枇杷 | | 1 | 105 | 4 | 4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Ficus hispida | 對葉榕 | | 7 | 110-250 | 4-6 | 3-5 | Low | Poor | Fair | Poor | | | Normal | Low | on slope |
| | | | Ligustrum sinense | 山指甲 | | 3 | 120-230 | 3-5 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Low | on slope |
| | | | Litsea glutinosa | 潺槁樹 | | 8 | 100-210 | 4-8 | 2-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Macaranga tanarius var. tomentosa | 血桐 | | 4 | 100-260 | 4-7 | 3-8 | Low | Poor | Poor | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Microcos nervosa | 破布葉 | | 1 | 110 | 5 | 3 | Low | Poor | Poor | Poor | Low | a, c, f, h | Normal | Low | on slope |
| | | | Sapium sebiferum | 烏桕 | | 4 | 150-450 | 6-10 | 3-8 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Sterculia lanceolata | 假蘋婆 | | 2 | 130-290 | 6-8 | 5-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Syzygium jambos | 蒲桃 | | 1 | 590 | 12 | 12 | Low | Poor | Fair | Poor | Low | a, b, c, d, f, h | Normal | Low | on slope |
| TG5 | LR2, LR3.3, LR6, LR8.1 | J/O CWBR | Acronychia pedunculata | 山油柑 | 325 | 5 | 100-150 | 2-5 | 2-3 | Low | Poor | Fair | Fair | Low | a, c, f, h | Normal | Low | on slope, shallow soil |
| | | | Archontophoenix alexandrae | 假檳榔 | | 2 | 155-160 | 7 | 3 | Med | Fair | Fair | Fair | Med | b | Normal | Med | - |
| | | | Acacia confusa | 台灣相思 | | 100 | 100-150 | 5-7 | 3-5 | Low | Poor | Fair | Poor | Low | a, b, c, f, h, i | Normal | Low | on slope, shallow soil |
| | | | Cratogeomys cochinchinense | 黃牛木 | | 4 | 100 | 8 | 3 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | at slope, leaning |
| | | | Casuarina equisetifolia | 木麻黃 | | 5 | 150-200 | 8-13 | 6-8 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | leaning |
| | | | Celtis sinensis | 朴樹 | | 14 | 100-230 | 4-7 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h, i | Normal | Low | on slope, shallow soil |
| | | | Cinnamomum camphora | 樟 | | 10 | 100-230 | 4-7 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h, i | Normal | Low | on slope, shallow soil |
| | | | Cinnamomum parthenoxylon | 黃樟 | | 10 | 100-230 | 4-7 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h, i | Normal | Low | on slope, shallow soil |
| | | | Delonix regia | 鳳凰木 | | 2 | 200-500 | 9-16 | 6-9 | Med | Fair | Fair | Fair | Low | b, d, h | Normal | Med | at slope |
| | | | Ficus microcarpa | 細葉榕 | | 20 | 100-740 | 3-16 | 2-8 | Low | Poor | Fair | Poor | Low | a, b, d, f, h | Normal | Med | at slope |
| | | | Ficus virens | 大葉榕 | | 5 | 210 | 7 | 5 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, multi-trunks |
| | | | Glochidion hirsutum | 厚葉算盤子 | | 9 | 120-155 | 6-7 | 3-5 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, multi-trunks |
| | | | Gordonia axillaris | 大頭茶 | | 30 | 100-120 | 3-6 | 1-3 | Low | Poor | Fair | Fair | Low | a, c, f, h | Normal | Low | on slope, shallow soil |
| | | | Litsea glutinosa | 潺槁樹 | | 5 | 120-150 | 2-5 | 2-4 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | on slope, shallow soil |
| | | | Lophostemon confertus | 紅膠木 | | 15 | 200-350 | 5-9 | 4-5 | Low | Poor | Fair | Poor | Low | a, b, c, d, f, h | Normal | Low | on slope, shallow soil |
| | | | Machilus breviflora | 短序潤楠 | | 15 | 100-150 | 3-7 | 3-6 | Low | Poor | Fair | Fair | Low | a, c, f, h | Normal | Low | on slope, shallow soil |
| Machilus pauhoi | 刨花潤楠 | 10 | 100-180 | 3-8 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope, shallow soil | | | | |

| Tree Group# | Relevant LRs | District | Botanical Name | Chinese Name | Estimated Total Number of Trees within this Tree Group | Estimated Number of Trees for this Species | Range of Trunk Diameter at DBH (mm) | Range of Height (m) | Range of Spread (m) | Overall Amenity Value (High/ Medium/ Low) | Overall Form (Good/ Fair/ Poor) | Overall Health (Good/ Fair/ Poor) | Overall Structural Condition (Good/ Fair/ Poor) | Overall Suitability for Transplanting (High/ Medium/ Low) | Overall Remark of Suitability for Transplanting | Overall Conservation Status (Normal/ Rare/ Endanger/ OVT/ Tree Register) | Overall Survival Rate after Transplant (High/ Medium/ Low) | Additional Remarks |
|----------------------|--------------|-----------|--------------------------------------|--------------|--|--|-------------------------------------|---------------------|---------------------|---|---------------------------------|-----------------------------------|---|---|---|--|--|--------------------------------------|
| | | | Machilus chekiangensis | 浙江潤楠 | | 5 | 265 | 9 | 6 | Med | Fair | Fair | Fair | Low | b, c, h | Normal | Low | at slope |
| | | | Mallotus paniculatus | 白楸 | | 10 | 100-120 | 3-7 | 2-4 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | on slope, shallow soil |
| | | | Ormosia emarginata | 凹葉紅豆 | | 5 | 100-130 | 3-6 | 3-4 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | on slope, shallow soil |
| | | | Pinus massoniana | 馬尾松 | | 10 | 100-120 | 3-9 | 2-4 | Low | Poor | Fair | Fair | Low | a, b, c, f, h | Normal | Low | on slope, shallow soil |
| | | | Polyspora axillaris | 大頭茶 | | 5 | 130 | 7 | 5 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, leaning |
| | | | Reevesia thyrsoidea | 梭羅樹 | | 5 | 100-130 | 3-6 | 3-4 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | on slope, shallow soil |
| | | | Schefflera heptaphylla | 鵝掌柴 | | 15 | 100-120 | 2-7 | 3-4 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | on slope, shallow soil |
| | | | Viburnum odoratissimum | 珊瑚樹 | | 4 | 115-220 | 7-8 | 3-5 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, leaning |
| | | | Schima superba | 木荷 | | 5 | 100-130 | 3-6 | 3-4 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | on slope, shallow soil |
| TC6 | LRL1 | J/O CWBR | Melaleuca cajuputi | 白千層 | 20 | 20 | 250-450 | 9-14 | 4-6 | Medium | Fair | Fair | Fair | Low | b, h | Normal | Medium | shallow soil, root growth restricted |
| J/O NCWBR | | | | | | | | | | | | | | | | | | |
| TC7 | LRL34 | J/O NCWBR | Alstonia scholaris | 糖膠樹 | 69 | 3 | 410-470 | 17-19 | 6-7 | Med | Fair | Fair | Fair | Low | b, c, d | Normal | Low | Too large in size |
| | | | Araucaria heterophylla | 異葉南洋杉 | | 4 | 240-290 | 14-16 | 4-5 | Med | Fair | Fair | Fair | Med | b | Normal | Med | - |
| | | | Archontophoenix alexandrae | 假檳榔 | | 1 | 155 | 5 | 4 | Med | Fair | Fair | Fair | Med | - | Normal | Med | - |
| | | | Artocarpus communis | 麵包樹 | | 3 | 150-190 | 5-6 | 4-5 | Low | Poor | Fair | Poor | Low | a, b, f | Normal | Med | low branching |
| | | | Bischofia polycarpa | 重陽木 | | 4 | 150-250 | 7-8 | 5-7 | Med | Fair | Fair | Fair | Med | b | Normal | Med | - |
| | | | Callistemon rigidus | 紅千層 | | 2 | 180-240 | 5-13 | 4-5 | Med | Fair | Fair | Fair | Med | b | Normal | Med | - |
| | | | Chukrasia tabularis | 麻楝 | | 1 | 215 | 6 | 4 | Low | Poor | Fair | Poor | Low | a, b, c, f | Normal | Low | leaning |
| | | | Elaeis guineensis | 油棕 | | 2 | 450-600 | 11-12 | 5-7 | Med | Fair | Fair | Fair | High | - | Normal | Med | - |
| | | | Elaeocarpus hainanensis | 水石榕 | | 6 | 120-260 | 5-6 | 5-7 | Low | Poor | Fair | Poor | Low | a, b, c, f | Normal | Low | leaning, low branching |
| | | | Erythrina corallodendron | 龍牙花 | | 1 | 265 | 5 | 5 | Low | Poor | Fair | Poor | Low | a, b, c, f | Normal | Low | leaning, low branching |
| | | | Hypophorbe laginicaulis | 酒瓶椰子 | | 11 | 100-250 | 2-5 | 2-4 | Med | Fair | Fair | Fair | Med | - | Normal | Med | - |
| | | | Livistona chinensis | 蒲葵 | | 4 | 230-330 | 3-5 | 2 | Med | Fair | Fair | Fair | High | - | Normal | Med | - |
| | | | Melaleuca cajuputi subsp. cumingiana | 白千層 | | 8 | 150-350 | 12-16 | 4-7 | Med | Fair | Fair | Fair | Med | b | Normal | Med | - |
| | | | Phoenix roebelenii | 日本葵 | | 1 | 110 | 4 | 3 | Med | Fair | Fair | Fair | High | - | Normal | Med | Multi-trunks |
| | | | Roystonea regia | 大王椰子 | | 17 | 150-300 | 8-15 | 4-6 | Med | Fair | Fair | Fair | Med | b | Normal | Med | - |
| Washingtonia robusta | 華盛頓葵 | 1 | 330 | 9 | 4 | Med | Fair | Fair | Fair | Med | b | Normal | Med | - | | | | |

| Tree Group# | Relevant LRs | District | Botanical Name | Chinese Name | Estimated Total Number of Trees within this Tree Group | Estimated Number of Trees for this Species | Range of Trunk Diameter at DBH (mm) | Range of Height (m) | Range of Spread (m) | Overall Amenity Value (High/ Medium/ Low) | Overall Form (Good/ Fair/ Poor) | Overall Health (Good/ Fair/ Poor) | Overall Structural Condition (Good/ Fair/ Poor) | Overall Suitability for Transplanting (High/ Medium/ Low) | Overall Remark of Suitability for Transplanting | Overall Conservation Status (Normal/ Rare/ Endanger/ OVT/ Tree Register) | Overall Survival Rate after Transplant (High/ Medium/ Low) | Additional Remarks |
|--------------------|--------------|-----------|----------------------------|--------------|--|--|-------------------------------------|---------------------|---------------------|---|---------------------------------|-----------------------------------|---|---|---|--|--|--------------------------------------|
| TG8 | LRI.2 | J/O NCWBR | Melaleuca cajuputi | 白千層 | 37 | 32 | 100-200 | 4-8 | 3-5 | Medium | Fair | Fair | Fair | Low | b, h | Normal | Medium | shallow soil, root growth restricted |
| | | | Peltophorum tonkinense | 銀珠 | | 3 | 150-350 | 8-11 | 4-7 | Medium | Fair | Fair | Fair | Low | b, h | Normal | Medium | shallow soil, root growth restricted |
| | | | Ficus virens | 大葉榕 | | 2 | 300-400 | 6-9 | 6-8 | Medium | Fair | Fair | Fair | Low | b, h | Normal | Medium | shallow soil, root growth restricted |
| TG9 | LR8.3 | O NCWBR | Acacia auriculiformis | 耳果相思 | 755 | 6 | 310 | 12 | 7 | Med | Fair | Fair | Fair | Low | b, c, d | Normal | Low | at slope |
| | | | Acacia confusa | 台灣相思 | | 350 | 100-670 | 2-16 | 3-12 | Low | Poor | Fair | Poor | Low | a, b, c, d, f, h | Normal | Low | at slope |
| | | | Acronychia pedunculata | 山油柑 | | 5 | 325 | 6 | 7 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope |
| | | | Ailanthus fordii | 福氏臭椿 | | 1 | 130 | 9 | 3 | Low | Poor | Fair | Poor | Low | a, f | Normal | Med | at slope |
| | | | Albizia lebbek | 大葉合歡 | | 17 | 95-300 | 5-11 | 2-6 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | at slope |
| | | | Aleurites moluccana | 石栗 | | 10 | 110-300 | 5-13 | 2-4 | Low | Poor | Fair | Poor | Low | a, f, h | Normal | Med | at slope |
| | | | Aporosa dioica | 銀柴 | | 3 | 110-305 | 9-11 | 4-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope |
| | | | Bombax ceiba | 木棉 | | 9 | 95-290 | 5-12 | 5 | Med | Fair | Fair | Fair | Low | b, h | Normal | Med | at slope |
| | | | Casuarina equisetifolia | 木麻黃 | | 10 | 110-340 | 5-13 | 4-6 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | at slope |
| | | | Celtis sinensis | 朴樹 | | 24 | 95-320 | 3-10 | 3-7 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | at slope |
| | | | Cinnamomum burmannii | 陰香 | | 3 | 95-140 | 3-4 | 3-4 | Low | Poor | Fair | Poor | Low | a, f, h | Normal | Med | at slope |
| | | | Cinnamomum camphora | 樟 | | 2 | 235-250 | 9-11 | 6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope |
| | | | Crateva unilocularis | 樹頭菜 | | 1 | 170 | 7 | 4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope |
| | | | Cratogeomys cochinchinense | 黃牛木 | | 6 | 100-150 | 4-8 | 3-5 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | at slope |
| | | | Dalbergia odorifera | 降香黃檀 | | 1 | 300 | 8 | 5 | Med | Fair | Fair | Fair | Low | b, c | Normal | Low | at slope |
| | | | Delonix regia | 鳳凰木 | | 2 | 410-500 | 14-16 | 7-8 | Med | Fair | Fair | Fair | Low | b, d, h | Normal | Med | at slope |
| | | | Eucalyptus citriodora | 檸檬桉 | | 8 | 200-500 | 14-18 | 6-8 | Med | Fair | Fair | Fair | Low | b, c, d, h | Normal | Low | at slope |
| | | | Eucalyptus robusta | 大葉桉 | | 1 | 685 | 10 | 8 | Med | Fair | Fair | Fair | Low | b, c, d | Normal | Low | at slope |
| | | | Eucalyptus torelliana | 毛葉桉 | | 2 | 295-395 | 11-13 | 5-6 | Med | Fair | Fair | Fair | Low | b, c, d, h | Normal | Low | at slope |
| | | | Ficus hispida | 對葉榕 | | 21 | 100-220 | 3-6 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | at slope |
| | | | Ficus microcarpa | 細葉榕 | | 20 | 100-740 | 3-16 | 2-8 | Low | Poor | Fair | Poor | Low | a, b, d, f, h | Normal | Med | at slope |
| Ficus subpisocarpa | 筆管榕 | 5 | 210-400 | 3-9 | 3-7 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | at slope | | | | |
| Grevillea robusta | 銀樺 | 4 | 100-330 | 6-14 | 3-5 | Med | Fair | Fair | Fair | Low | b, h | Normal | Med | at slope | | | | |
| Ilex rotunda | 鐵冬青 | 12 | 100-300 | 2-12 | 2-6 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | at slope | | | | |

| Tree Group# | Relevant LRs | District | Botanical Name | Chinese Name | Estimated Total Number of Trees within this Tree Group | Estimated Number of Trees for this Species | Range of Trunk Diameter at DBH (mm) | Range of Height (m) | Range of Spread (m) | Overall Amenity Value (High/ Medium/ Low) | Overall Form (Good/ Fair/ Poor) | Overall Health (Good/ Fair/ Poor) | Overall Structural Condition (Good/ Fair/ Poor) | Overall Suitability for Transplanting (High/ Medium/ Low) | Overall Remark of Suitability for Transplanting | Overall Conservation Status (Normal/ Rare/ Endanger/ OVT/ Tree Register) | Overall Survival Rate after Transplant (High/ Medium/ Low) | Additional Remarks |
|--------------------|--------------|-----------|-----------------------------------|--------------|--|--|-------------------------------------|---------------------|---------------------|---|---------------------------------|-----------------------------------|---|---|---|--|--|--------------------|
| | | h | Leucaena leucocephala | 銀合歡 | | 15 | 95-120 | 4-5 | 3-4 | Low | Poor | Fair | Poor | Low | a, c, f, g, h | Normal | Low | at slope |
| | | | Ligustrum sinense | 山指甲 | | 13 | 95-160 | 3-6 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | at slope |
| | | | Litsea glutinosa | 潺槁樹 | | 20 | 100-275 | 2-10 | 3-7 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | at slope |
| | | | Litsea monopetala | 假柿樹 | | 1 | 360 | 9 | 7 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope |
| | | | Lophostemon confertus | 紅膠木 | | 6 | 100-300 | 6-12 | 2-6 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | at slope |
| | | | Macaranga tanarius var. tomentosa | 血桐 | | 78 | 95-275 | 3-9 | 3-9 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope |
| | | | Machilus chekiangensis | 浙江潤楠 | | 15 | 155 | 6 | 3 | Med | Fair | Fair | Fair | Low | c, h | Normal | Low | at slope |
| | | | Mallotus paniculatus | 白楸 | | 27 | 100-220 | 4-9 | 3-8 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | at slope |
| | | | Mangifera indica | 芒果 | | 3 | 120-150 | 6-9 | 3-4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope |
| | | | Melia azedarach | 苦楝 | | 4 | 150-330 | 6-12 | 5-6 | Low | Poor | Fair | Poor | Low | a, b, c, d, f, h | Normal | Low | at slope |
| | | | Michelia x alba | 白蘭 | | 3 | 110-195 | 6-9 | 2-5 | Med | Fair | Fair | Fair | Low | b | Normal | Med | at slope |
| | | | Microcos nervosa | 破布葉 | | 1 | 140 | 5 | 4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope |
| | | | Morus alba | 桑 | | 1 | 130 | 5 | 4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope |
| | | | Rhus succedanea | 野漆樹 | | 2 | 120-135 | 8 | 4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope |
| | | | Sapium discolor | 山烏柏 | | 1 | 100 | 7 | 2 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | at slope |
| | | | Sapium sebiferum | 烏柏 | | 5 | 100-300 | 5-10 | 3-9 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope |
| | | | Schefflera heptaphylla | 鵝掌柴 | | 15 | 95-250 | 4-8 | 2-7 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope |
| | | | Schima superba | 木荷 | | 1 | 110 | 6 | 4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope |
| | | | Spathodea campanulata | 火焰樹 | | 12 | 115-145 | 3-8 | 2-4 | Low | Poor | Poor | Poor | Low | a, c, f, h | Normal | Low | at slope |
| | | | Sterculia lanceolata | 假蒺藜 | | 6 | 100-150 | 5-8 | 2-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope |
| Syzygium cumini | 海南蒲桃 | 1 | 105 | 5 | 3 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | at slope | | | | |
| Thespesia populnea | 恒春黃槿 | 1 | 150 | 4 | 4 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | at slope | | | | |
| Vernicia montana | 木油樹 | 1 | 100-150 | 5-9 | 3-4 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | at slope | | | | |
| TG10 | LR4, LR5 | J/O NCWBR | Aleurites moluccana | 石栗 | 42 | 25 | 250-450 | 6-13 | 4-8 | Med | Fair | Fair | Fair | Med | b | Normal | Med | - |
| | | | Archontophoenix alexandrae | 假檳榔 | | 5 | 100-150 | 5-8 | 2-6 | Med | Fair | Fair | Fair | Med | b | Normal | Med | - |
| | | | Bauhinia spp. | 羊蹄甲屬 | | 2 | 150-200 | 4-8 | 2-5 | Med | Fair | Fair | Fair | Med | - | Normal | Med | - |
| | | | Lagerstroemia speciosa | 大花紫薇 | | 5 | 150-200 | 4-8 | 2-5 | Med | Fair | Fair | Fair | Med | - | Normal | Med | - |

| Tree Group# | Relevant LRs | District | Botanical Name | Chinese Name | Estimated Total Number of Trees within this Tree Group | Estimated Number of Trees for this Species | Range of Trunk Diameter at DBH (mm) | Range of Height (m) | Range of Spread (m) | Overall Amenity Value (High/ Medium/ Low) | Overall Form (Good/ Fair/ Poor) | Overall Health (Good/ Fair/ Poor) | Overall Structural Condition (Good/ Fair/ Poor) | Overall Suitability for Transplanting (High/ Medium/ Low) | Overall Remark of Suitability for Transplanting | Overall Conservation Status (Normal/ Rare/ Endanger/ OVT/ Tree Register) | Overall Survival Rate after Transplant (High/ Medium/ Low) | Additional Remarks |
|----------------------|-------------------------|----------|-----------------------------------|--------------|--|--|-------------------------------------|---------------------|---------------------|---|---------------------------------|-----------------------------------|---|---|---|--|--|------------------------|
| | | | Ficus religiosa | 菩提樹 | | 2 | 150-200 | 4-8 | 2-5 | Med | Fair | Fair | Fair | Med | - | Normal | Med | - |
| | | | Ficus elastica | 印度橡樹 | | 3 | 150-750 | 6-11 | 6-12 | Med | Fair | Fair | Fair | Low | b | Normal | Med | - |
| TG11 | LR3.3, LR4, LR8.2, LG11 | J/ONCWB | Acacia confusa | 台灣相思 | 447 | 230 | 100-670 | 2-16 | 3-12 | Low | Poor | Fair | Poor | Low | a, b, c, d, f, h | Normal | Low | at slope |
| | | | Acronychia pedunculata | 山油柑 | | 4 | 100-200 | 3-4 | 3-4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope |
| | | | Albizia lebbek | 大葉合歡 | | 7 | 95-300 | 5-11 | 2-6 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | at slope |
| | | | Aporosa dioica | 銀柴 | | 3 | 110-305 | 9-11 | 4-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope |
| | | | Casuarina equisetifolia | 木麻黃 | | 5 | 100-350 | 5-12 | 4-6 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | at slope |
| | | | Celtis sinensis | 朴樹 | | 14 | 95-200 | 3-8 | 3-5 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | at slope |
| | | | Cratoxylum cochinchinense | 黃牛木 | | 6 | 100-150 | 4-8 | 3-5 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | at slope |
| | | | Ficus hispida | 對葉榕 | | 21 | 100-220 | 3-6 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | at slope |
| | | | Ficus microcarpa | 細葉榕 | | 15 | 100-300 | 3-12 | 2-6 | Low | Poor | Fair | Poor | Low | a, b, d, f, h | Normal | Med | at slope |
| | | | Ficus subpisocarpa | 筆管榕 | | 6 | 210-400 | 3-9 | 3-7 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | at slope |
| | | | Ilex rotunda | 鐵冬青 | | 5 | 100-200 | 2-10 | 2-6 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | at slope |
| | | | Leucaena leucocephala | 銀合歡 | | 5 | 95-200 | 4-6 | 3-4 | Low | Poor | Fair | Poor | Low | a, c, f, g, h | Normal | Low | at slope |
| | | | Ligustrum sinense | 山指甲 | | 5 | 95-160 | 3-6 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | at slope |
| | | | Litsea glutinosa | 潺槁樹 | | 15 | 100-250 | 2-8 | 3-7 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | at slope |
| | | | Macaranga tanarius var. tomentosa | 血桐 | | 58 | 95-250 | 3-9 | 3-9 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope |
| | | | Mallotus paniculatus | 白楸 | | 17 | 100-220 | 4-9 | 3-8 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | at slope |
| | | | Pinus massoniana | 馬尾松 | | 5 | 100-120 | 3-9 | 2-4 | Low | Poor | Fair | Fair | Low | a, b, c, f, h | Normal | Low | on slope, shallow soil |
| | | | Rhus succedanea | 野漆樹 | | 2 | 120-130 | 5-8 | 3-4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope |
| | | | Sapium sebiferum | 烏柏 | | 5 | 100-200 | 5-6 | 3-5 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope |
| | | | Schefflera heptaphylla | 鵝掌柴 | | 13 | 95-250 | 4-8 | 2-7 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope |
| Sterculia lanceolata | 假蘋婆 | 6 | 100-150 | 5-8 | 2-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope | | | | |
| TG12 | LR3.3 | J/OCWB | Acacia confusa | 台灣相思 | 67 | 15 | 100-620 | 5-16 | 4-14 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Ficus hispida | 對葉榕 | | 5 | 100-400 | 4-11 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | on slope |
| | | | Litsea glutinosa | 潺槁樹 | | 15 | 100-250 | 4-8 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Macaranga tanarius var. tomentosa | 血桐 | | 5 | 100-260 | 4-9 | 4-8 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |

| Tree Group# | Relevant LRs | District | Botanical Name | Chinese Name | Estimated Total Number of Trees within this Tree Group | Estimated Number of Trees for this Species | Range of Trunk Diameter at DBH (mm) | Range of Height (m) | Range of Spread (m) | Overall Amenity Value (High/ Medium/ Low) | Overall Form (Good/ Fair/ Poor) | Overall Health (Good/ Fair/ Poor) | Overall Structural Condition (Good/ Fair/ Poor) | Overall Suitability for Transplanting (High/ Medium/ Low) | Overall Remark of Suitability for Transplanting | Overall Conservation Status (Normal/ Rare/ Endanger/ OVT/ Tree Register) | Overall Survival Rate after Transplant (High/ Medium/ Low) | Additional Remarks |
|-------------|--------------|----------|------------------------|--------------|--|--|-------------------------------------|---------------------|---------------------|---|---------------------------------|-----------------------------------|---|---|---|--|--|--------------------|
| | | | Mallotus paniculatus | 白楸 | | 15 | 100-210 | 5-11 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Rhus succedanea | 野漆樹 | | 7 | 100-150 | 5-7 | 3-4 | Low | Poor | Poor | Poor | Low | a, c, f, h | Normal | Low | on slope |
| | | | Schefflera heptaphylla | 鵝掌柴 | | 5 | 100-220 | 4-10 | 4-6 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | on slope |

J/O SMPR

| | | | | | | | | | | | | | | | | | | |
|------|-------|----------|--------------------------------------|-------|-----|----|---------|------|------|-----|------|------|------|-----|---------------------|--------|-----|-------------------------|
| TG13 | LRI.3 | J/O SMPR | Celtis sinensis | 朴樹 | 133 | 10 | 100-230 | 6-9 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h, i | Normal | Low | root growth restricted |
| | | | Melaleuca cajuputi subsp. cumingiana | 白千層 | | 27 | 100-360 | 6-14 | 2-6 | Med | Fair | Fair | Fair | Med | b | Normal | Med | - |
| | | | Leucaena leucocephala | 銀合歡 | | 63 | 100-370 | 6-14 | 2-8 | Low | Poor | Fair | Poor | Low | a, b, c, f, g, h, i | Normal | Low | imbalanced crown |
| | | | Liquidambar formosana | 楓香 | | 9 | 130-250 | 6-9 | 4-8 | Low | Poor | Fair | Poor | Low | a, b, c, f, i | Normal | Low | imbalanced crown |
| | | | Macaranga tanarius | 血桐 | | 4 | 100-150 | 6-7 | 4-6 | Low | Poor | Fair | Poor | Low | a, c, f, i | Normal | Low | Leaning |
| | | | Acacia confusa | 台灣相思 | | 9 | 150-380 | 7-10 | 6-9 | Low | Poor | Fair | Poor | Low | a, b, c, f, i | Normal | Low | imbalanced crown |
| | | | Thevetia peruviana | 黃花夾竹桃 | | 4 | 140-190 | 5-7 | 3-6 | Low | Poor | Fair | Poor | Low | a, c, f, h, i | Normal | Low | - |
| | | | Bauhinia x blakeana | 洋紫荊 | | 4 | 100-150 | 4-8 | 4-6 | Low | Poor | Fair | Poor | Low | a, c, f, i | Normal | Low | - |
| | | | Ficus elastica | 印度橡樹 | | 3 | 150-750 | 6-11 | 6-12 | Med | Fair | Fair | Fair | Low | b, d | Normal | Med | suspressed by structure |

| | | | | | | | | | | | | | | | | | | |
|--|--|--|----------------------------|-------|--|-----|----------|-------|------|-----|------|------|------|-----|------------------|--------|-----|---|
| | | | Acacia confusa | 台灣相思 | | 180 | 100-550 | 6-16 | 2-10 | Low | Poor | Fair | Poor | Low | a, b, c, d, f, h | Normal | Low | at slope, multi-trunks |
| | | | Acronychia pedunculata | 山油柑 | | 10 | 200 | 7 | 5 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at rock, leaning, low branching |
| | | | Albizia lebbeck | 大葉合歡 | | 20 | 120-500 | 6-16 | 2-8 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | at slope, many pruning wounds |
| | | | Bauhinia variegata | 宮粉羊蹄甲 | | 55 | 100-260 | 8 | 5 | Low | Poor | Fair | Poor | Low | a, b, c, f | Normal | Low | at raised planter, leaning, low branching |
| | | | Bridelia tomentosa | 土蜜樹 | | 5 | 130 | 5 | 4 | Low | Poor | Poor | Poor | Low | a, b, c, f, h | Normal | Low | at slope, leaning, low branching |
| | | | Casuarina equisetifolia | 木麻黃 | | 99 | 95-350 | 6-16 | 2-8 | Low | Poor | Fair | Poor | Low | a, f, h | Normal | Med | at slope, leaning |
| | | | Celtis sinensis | 朴樹 | | 34 | 120-300 | 5-10 | 2-7 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, leaning |
| | | | Cinnamomum camphora | 樟 | | 9 | 220 | 9 | 4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, leaning |
| | | | Cratogeomys cochinchinense | 黃牛木 | | 5 | 130-170 | 7 | 4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, leaning |
| | | | Duranta erecta | 假連翹 | | 17 | 100-200 | 4-6 | 3-4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, multi-trunks |
| | | | Eucalyptus citriodora | 檸檬桉 | | 13 | 100-400 | 13-14 | 4-6 | Med | Fair | Fair | Fair | Low | b, c, h, h | Normal | Low | at slope |
| | | | Ficus hispida | 對葉榕 | | 15 | 100-250 | 5-9 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | at slope, forked, low branching |
| | | | Ficus microcarpa | 細葉榕 | | 36 | 120-1000 | 4-14 | 3-10 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | at slope, multi-trunks |
| | | | Ficus subpisocarpa | 筆管榕 | | 18 | 95-550 | 5-9 | 4-12 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at rocky slope, multi-trunks, low branching |

| Tree Group# | Relevant LRs | District | Botanical Name | Chinese Name | Estimated Total Number of Trees within this Tree Group | Estimated Number of Trees for this Species | Range of Trunk Diameter at DBH (mm) | Range of Height (m) | Range of Spread (m) | Overall Amenity Value (High/ Medium/ Low) | Overall Form (Good/ Fair/ Poor) | Overall Health (Good/ Fair/ Poor) | Overall Structural Condition (Good/ Fair/ Poor) | Overall Suitability for Transplanting (High/ Medium/ Low) | Overall Remark of Suitability for Transplanting | Overall Conservation Status (Normal/ Rare/ Endanger/ OVT/ Tree Register) | Overall Survival Rate after Transplant (High/ Medium/ Low) | Additional Remarks |
|------------------------|--------------|----------|-----------------------------------|--------------|--|--|-------------------------------------|---------------------|---------------------|---|---------------------------------|-----------------------------------|---|---|---|--|--|---|
| TG14 | LR8.4 | J/O SMPR | Ficus variegata | 青果榕 | 775 | 5 | 300 | 8 | 5 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, leaning, forked |
| | | | Ficus virens | 大葉榕 | | 9 | 210 | 7 | 5 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, multi-trunks |
| | | | Garcinia oblongifolia | 嶺南山竹子 | | 4 | 100 | 7 | 4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, leaning, climbers |
| | | | Leucaena leucocephala | 銀合歡 | | 40 | 100-300 | 6-13 | 4-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, g, h | Normal | Low | at slope, multi-trunks, weedy species |
| | | | Litsea glutinosa | 潺槁樹 | | 6 | 130-210 | 6-11 | 4-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, multi-trunks, climbers |
| | | | Lophostemon confertus | 紅膠木 | | 12 | 170-270 | 8-13 | 5-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, leaning, climbers |
| | | | Macaranga tanarius var. tomentosa | 血桐 | | 32 | 100-400 | 5-12 | 3-7 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | at slope, leaning, many wounds, unbalanced form |
| | | | Machilus chekiangensis | 浙江潤楠 | | 12 | 350-400 | 9-14 | 6-8 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, leaning |
| | | | Mallotus paniculatus | 白楸 | | 33 | 100-250 | 6-11 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, leaning |
| | | | Melia azedarach | 苦楝 | | 3 | 120-250 | 7-8 | 4-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, crooked trunk, trunk conflicted with wall |
| | | | Phyllanthus emblica | 餘甘子 | | 12 | 100-200 | 5-6 | 3-4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, leaning |
| | | | Rhus succedanea | 野漆樹 | | 5 | 130-250 | 6-9 | 3-6 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | at slope, leaning |
| | | | Schefflera heptaphylla | 鵝掌柴 | | 10 | 100-250 | 5-7 | 3-4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, multi-trunks, climbers |
| | | | Sterculia lanceolata | 假蒴婆 | | 45 | 100-300 | 4-12 | 3-7 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, leaning |
| | | | Thevetia peruviana | 黃花夾竹桃 | | 28 | 100-250 | 4-8 | 3-6 | Low | Poor | Fair | Poor | Low | a, f, h | Normal | Med | at slope, bending trunk |
| Viburnum odoratissimum | 珊瑚樹 | 2 | 100-120 | 6-8 | 3-4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, double trunks | | | | |
| Zanthoxylum avicennae | 筋櫛花椒 | 1 | 155 | 5 | 4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, leaning, low branching | | | | |
| TG15 | LR3.5 | J/O SMPR | Acacia confusa | 台灣相思 | 125 | 25 | 100-620 | 5-16 | 4-14 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Ficus hispida | 對葉榕 | | 10 | 100-400 | 4-11 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | on slope |
| | | | Litsea glutinosa | 潺槁樹 | | 30 | 100-250 | 4-8 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Macaranga tanarius var. tomentosa | 血桐 | | 10 | 100-260 | 4-9 | 4-8 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Mallotus paniculatus | 白楸 | | 25 | 100-210 | 5-11 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Rhus succedanea | 野漆樹 | | 10 | 100-150 | 5-7 | 3-4 | Low | Poor | Poor | Poor | Low | a, c, f, h | Normal | Low | on slope |
| | | | Schefflera heptaphylla | 鵝掌柴 | | 15 | 100-220 | 4-10 | 4-6 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | on slope |
| TG16 | LR4 | J/O SMPR | Aleurites moluccana | 石栗 | 57 | 35 | 250-450 | 6-13 | 4-8 | Med | Fair | Fair | Fair | Med | - | Normal | Med | - |
| | | | Bauhinia spp. | 羊蹄甲屬 | | 2 | 150-200 | 4-8 | 2-5 | Med | Fair | Fair | Fair | Med | - | Normal | Low | - |
| | | | Lagerstroemia speciosa | 大花紫薇 | | 5 | 150-200 | 4-8 | 2-5 | Med | Fair | Fair | Fair | Med | - | Normal | Med | - |

| Tree Group# | Relevant LRs | District | Botanical Name | Chinese Name | Estimated Total Number of Trees within this Tree Group | Estimated Number of Trees for this Species | Range of Trunk Diameter at DBH (mm) | Range of Height (m) | Range of Spread (m) | Overall Amenity Value (High/ Medium/ Low) | Overall Form (Good/ Fair/ Poor) | Overall Health (Good/ Fair/ Poor) | Overall Structural Condition (Good/ Fair/ Poor) | Overall Suitability for Transplanting (High/ Medium/ Low) | Overall Remark of Suitability for Transplanting | Overall Conservation Status (Normal/ Rare/ Endanger/ OVT/ Tree Register) | Overall Survival Rate after Transplant (High/ Medium/ Low) | Additional Remarks |
|----------------------|--------------|----------|--------------------------------------|--------------|--|--|-------------------------------------|---------------------|---------------------|---|---------------------------------|-----------------------------------|---|---|---|--|--|--|
| | | | Senna surattensis | 黃槐 | | 5 | 100-120 | 4-6 | 3-4 | Med | Fair | Fair | Fair | Med | - | Normal | Med | - |
| | | | Delonix regia | 鳳凰木 | | 10 | 410-500 | 14-16 | 7-8 | Med | Fair | Fair | Fair | Low | b, d, h | Normal | Med | - |
| TG17 | LR8.5, LR9.2 | J/O SMPR | Melaleuca cajuputi subsp. cumingiana | 白千層 | 865 | 203 | 95-650 | 3-12 | 3-9 | Low | Poor | Fair | Poor | Low | a, b, h, c, f, h | Normal | Low | at slope, leaning |
| | | | Acacia confusa | 台灣相思 | | 81 | 200-350 | 8-10 | 4-9 | Low | Poor | Fair | Poor | Low | a, b, c, d, f, h | Normal | Low | at slope, leaning |
| | | | Albizia lebbbeck | 大葉合歡 | | 2 | 140-200 | 7-8 | 2-3 | Low | Poor | Fair | Fair | Low | a, f, h | Normal | Med | unbalanced crown, growth restricted |
| | | | Bombax ceiba | 木棉 | | 54 | 100-300 | 7-16 | 3-8 | Low | Poor | Fair | Poor | Low | a, f, h | Normal | Med | at slope, unbalanced crown |
| | | | Casuarina equisetifolia | 木麻黃 | | 24 | 100-220 | 3-12 | 2-8 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, leaning |
| | | | Celtis sinensis | 朴樹 | | 1 | 170 | 5 | 4 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Medium | at slope, leaning |
| | | | Cinnamomum burmannii | 陰香 | | 2 | 100-180 | 4-5 | 2-4 | Low | Poor | Fair | Poor | Low | a, c, f, h | Normal | Low | at slope, forked |
| | | | Cratoxylum cochinchinense | 黃牛木 | | 15 | 200-300 | 10-15 | 4-9 | Low | Poor | Fair | Poor | Low | a, b, h, c, f, h | Normal | Low | at slope, leaning |
| | | | Eucalyptus citriodora | 檸檬桉 | | 26 | 100-250 | 3-8 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, h, f, h | Normal | Med | at slope, multi-trunks |
| | | | Ficus hispida | 對葉榕 | | 55 | 100-500 | 5-9 | 5-9 | Low | Poor | Fair | Poor | Low | a, b, h, f, h | Normal | Med | at slope, multi-trunks |
| | | | Ficus microcarpa | 細葉榕 | | 75 | 200-450 | 7-9 | 4-9 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, leaning, low branching |
| | | | Ficus subpisocarpa | 筆管榕 | | 5 | 100-200 | 6-7 | 3-5 | Low | Poor | Fair | Poor | Low | a, b, h, c, f, h | Normal | Low | at slope, leaning, roots restricted by drain |
| | | | Ficus variegata | 青果榕 | | 88 | 95-400 | 7-14 | 3-10 | Low | Poor | Poor | Poor | Low | a, b, c, f, g | Normal | Low | at raised planter, leaning, weedy species |
| | | | Leucaena leucocephala | 銀合歡 | | 69 | 100-300 | 4-11 | 3-7 | Low | Poor | Fair | Poor | Low | a, b, h, c, f, h | Normal | Low | at slope, leaning, low branching |
| | | | Macaranga tanarius var. tomentosa | 血桐 | | 60 | 100-200 | 4-9 | 2-7 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, leaning |
| | | | Mallotus paniculatus | 白楸 | | 5 | 150-170 | 4-7 | 2 | Low | Poor | Fair | Poor | Low | a, f | Normal | Med | leaning, unbalanced crown |
| | | | Melaleuca cajuputi subsp. cumingiana | 白千層 | | 5 | 150-370 | 7-14 | 5-8 | Low | Poor | Fair | Poor | Low | a, b, h, c, f, h | Normal | Low | at slope, forked |
| | | | Melia azedarach | 苦楝 | | 1 | 170 | 12 | 6 | Low | Poor | Fair | Poor | Low | a, b, h, c, f, h | Normal | Low | at slope, leaning |
| | | | Morus alba | 桑 | | 1 | 190 | 11 | 6 | Low | Poor | Poor | Poor | Low | a, b, f, h | Normal | Medium | at slope, broken main trunk, climbers |
| | | | Pinus elliotii | 濕地松 | | 4 | 120-400 | 7-8 | 6-7 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | at slope, leaning |
| Rhus succedanea | 木蠟樹 | 9 | 100-170 | 3-10 | 2-6 | Low | Poor | Fair | Poor | Low | a, b, h, c, f, h | Normal | Low | at slope, trunk conflicted by soil erosion mat | | | | |
| Sterculia lanceolata | 假蒺藜 | 50 | 100-230 | 4-10 | 3-7 | Low | Poor | Fair | Poor | Low | a, f, h | Normal | Med | at slope, leaning, forked | | | | |
| Thevetia peruviana | 黃花夾竹桃 | 30 | 250-450 | 6-13 | 4-8 | Med | Fair | Fair | Fair | Low | b, h | Normal | Med | at slope | | | | |
| | | | Acacia confusa | 台灣相思 | | 15 | 100-420 | 5-13 | 4-10 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Ficus hispida | 對葉榕 | | 10 | 100-200 | 4-11 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | on slope |

| Tree Group# | Relevant LRs | District | Botanical Name | Chinese Name | Estimated Total Number of Trees within this Tree Group | Estimated Number of Trees for this Species | Range of Trunk Diameter at DBH (mm) | Range of Height (m) | Range of Spread (m) | Overall Amenity Value (High/ Medium/ Low) | Overall Form (Good/ Fair/ Poor) | Overall Health (Good/ Fair/ Poor) | Overall Structural Condition (Good/ Fair/ Poor) | Overall Suitability for Transplanting (High/ Medium/ Low) | Overall Remark of Suitability for Transplanting | Overall Conservation Status (Normal/ Rare/ Endanger/ OVT/ Tree Register) | Overall Survival Rate after Transplant (High/ Medium/ Low) | Additional Remarks |
|-------------|--------------|----------|-----------------------------------|--------------|--|--|-------------------------------------|---------------------|---------------------|---|---------------------------------|-----------------------------------|---|---|---|--|--|--------------------|
| TG18 | LR34, LR4 | J/O SMPR | Litsea glutinosa | 潺槁樹 | 90 | 10 | 100-200 | 4-8 | 3-4 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Macaranga tanarius var. tomentosa | 血桐 | | 15 | 100-250 | 4-9 | 4-8 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Mallotus paniculatus | 白楸 | | 20 | 100-210 | 5-10 | 3-6 | Low | Poor | Fair | Poor | Low | a, b, c, f, h | Normal | Low | on slope |
| | | | Schefflera heptaphylla | 鵝掌柴 | | 20 | 100-220 | 4-9 | 4-6 | Low | Poor | Fair | Poor | Low | a, b, f, h | Normal | Med | on slope |
| TG19 | LR4 | J/O SMPR | Livistona chinensis | 蒲葵 | 30 | 10 | 230-330 | 3-5 | 2 | Med | Fair | Fair | Fair | High | - | Normal | Med | - |
| | | | Ficus microcarpa | 細葉榕 | | 10 | 200-450 | 7-9 | 4-9 | Med | Fair | Fair | Fair | High | - | Normal | Med | - |
| | | | Elaeocarpus spp. | 杜英 | | 10 | 150-200 | 4-8 | 2-5 | Med | Fair | Fair | Fair | Med | - | Normal | Med | - |

Remarks for Suitability for Transplanting

- (a) Low amenity value;
- (b) Irrecoverable form after transplanting (e.g. if substantial crown and root pruning are necessary to facilitate the transplanting);
- (c) Low survival rate after transplanting;
- (d) Very large size (unless the feasibility to transplant has been considered financially reasonable and technically feasible during the feasibility stage);
- (e) With evidence of over-maturity and onset of senescence;
- (f) With poor health, structure or form (e.g. imbalanced form, leaning, with major cavity/cracks/splits); or cavity/cracks/splits); or
- (g) Undesirable species (e.g. Leucaena leucocephala which is an invasive exotic tree).
- (h) On steep slope or area where formation of a root ball of reasonable size is not practicable