

Appendix 8.11 Intertidal Organisms Recorded in the Quantitative Survey within the 500m Assessment Area (Wet Season)

No.	Sampling point	I-A			I-B		
	Transect	H	M	L	H	M	L
1	<i>Alpheus lobidens</i>	0	1	0	0	0	0
2	<i>Amphibalanus reticulatus</i>	0	0	1	0	0	9
3	<i>Batillaria multiformis</i>	0	12	0	58	10	8
4	<i>Batillaria zonalis</i>	0	24	0	12	0	0
5	<i>Brachidontes variabilis</i>	0	3	1	0	0	2
6	<i>Circe scripta</i>	0	0	1	0	0	1
7	<i>Clibanarius</i> sp.	0	0	5	0	0	0
8	<i>Clypeomorus humilis</i>	0	0	0	0	11	6
9	<i>Corallina</i> sp.	0	1	0	0	0	0
10	<i>Diogenes spinifrons</i>	0	6	16	0	0	0
11	<i>Gaetice</i> sp.	0	1	0	0	0	0
12	<i>Grafrarium pectinatum</i>	0	1	0	1	0	2
13	<i>Haliplanella lineata</i>	0	1	0	0	0	0
14	<i>Holothuria spilonota</i>	0	1	0	0	0	0
15	<i>Hydroides</i> sp.	0	8	13	0	0	8
16	<i>Leptodius exaratus</i>	0	1	0	0	0	0
17	<i>Lunella coronata</i>	0	19	0	0	0	0
18	<i>Nanosesarma minutum</i>	0	4	0	0	0	0
19	<i>Nassarius festivus</i>	0	0	10	0	0	0
20	<i>Nerita albicilla</i>	0	1	0	0	0	0
21	<i>Planaxis sulcatus</i>	0	1	0	0	0	0
22	<i>Saccostrea cucullata</i>	0	12	8	1	0	6
23	Sipunculid worm	0	2	0	0	0	0
24	<i>Thais clavigera</i>	0	4	0	0	0	0
25	<i>Thalamita crenata</i>	0	1	0	0	0	0