

**Appendix 8.16**

Biomass (g) of Subtidal Benthos Species Recorded at each Sampling Location within the 500m Assessment Area during Dry Season (replicates of sub-stations are pooled together)

No.	Phylum	Class	Order	Family	Species	B1	B2	B3
1	Annelida	Polychaeta	Amphinomida	Amphinomidae	<i>Amphinome rostrata</i>	0.0137	0	0
2	Annelida	Polychaeta	Eunicida	Onuphidae	<i>Diopatra chiliensis</i>	1.5187	0.083	0
3	Annelida	Polychaeta	Capitellida	Maldanidae	<i>Euclymene</i> sp.	0.0875	0	0
4	Annelida	Polychaeta	Phyllodocida	Syllidae	<i>Eusyllis</i> sp.	0.0009	0	0
5	Annelida	Polychaeta	Phyllodocida	Glyceridae	<i>Glycera onomichiensis</i>	0.0767	0	0
6	Annelida	Polychaeta	Phyllodocida	Goniadidae	<i>Glycinde gurjanovae</i>	0.0048	0	0
7	Annelida	Polychaeta	Eunicida	Eunicidae	<i>Marphysa sanguinea</i>	0.0475	0	0
8	Annelida	Polychaeta	Phyllodocida	Nereidae	<i>Nereis</i> sp.	0.0753	0	0.0024
9	Annelida	Polychaeta	Phyllodocida	Pilargiidae	<i>Sigambra hanaokai</i>	0.0117	0	0
10	Annelida	Polychaeta	Terebellida	Trichobranchidae	<i>Terebellides stroemii</i>	0.1484	0	0
11	Arthropoda	Crustacea	Decapoda	Alpheidae	<i>Alpheus</i> sp.	0.0529	0	0
12	Arthropoda	Crustacea	Stomatopoda	Squillidae	<i>Oratosquilla oratoria</i>	0.0612	0	0
13	Mollusca	Bivalvia	Veneroida	Veneridae	<i>Anomalodiscus aquamosus</i>	0	0	3.1405
14	Mollusca	Bivalvia	Veneroida	Veneridae	<i>Circe tumefacta</i>	0	0	2.8078
15	Mollusca	Bivalvia	Veneroida	Cardiidae	<i>Fulvia aperta</i>	0	0	0.2212
16	Mollusca	Bivalvia	Veneroida	Tellinidae	<i>Moerella culter</i>	1.4493	0.6037	0
17	Mollusca	Bivalvia	Veneroida	Tellinidae	<i>Nitidotellina iridella</i>	4.8048	2.0991	0.6999
18	Mollusca	Bivalvia	Veneroida	Veneridae	<i>Timoclea imbricata</i>	0	0.0722	0