

Appendix 8.9A Intertidal Organisms Recorded in the Quantitative Survey within the 500m Assessment Area (Wet season)

No.	Sampling Location	I-A			I-B		
	Species / Tidal Level	H	M	L	H	M	L
1	<i>Acanthopleura japonica</i>	0	1	0	0	0	0
2	<i>Anthocidaris crassipina</i>	0	0	1	0	0	0
3	<i>Barbatia virescens</i>	0	1	0	0	0	0
4	<i>Captiulum mitella</i>	0	1	0	0	0	0
5	<i>Cellana grata</i>	0	1	1	0	0	1
6	<i>Cellana toreuma</i>	0	0	1	0	1	0
7	<i>Clypeomorus humilis</i>	0	26	0	0	0	1
8	<i>Cronia margariticola</i>	1	1	1	0	0	10
9	<i>Donax</i> sp.	0	0	0	0	7	0
10	<i>Holothuria spilonota</i>	0	0	1	0	0	0
11	<i>Ligia exotica</i>	0	0	0	1	0	0
12	<i>Littoraria articulata</i>	0	1	0	0	0	0
13	<i>Lunella coronata</i>	1	14	16	0	1	5
14	<i>Monodonta neritoides</i>	1	0	0	0	0	0
15	<i>Patelloida pygmaea</i>	0	0	1	0	0	0
16	<i>Saccostrea cucullata</i>	0	2	6	0	0	0
17	<i>Siphonaria japonica</i>	0	0	0	0	2	0
18	<i>Thais clavigera</i>	0	3	1	0	6	0
19	Green algae	0	0	0	0	0	50%
20	Red encrusting algae	0	0	0	0	0	30%
21	<i>Bathygobius fuscus</i>	2	0	0	0	0	0
22	<i>Hapalospongidion gelatinosum</i>	0	0	0	0	0	5%