

Appendix 9.6a Benthos Survey Results (Abundance) during Wet Season

| Phylum | Family | Species | B1 | | | B2 | | |
|----------|------------|-------------------------------|----|----|----|----|----|----|
| | | | R1 | R2 | R3 | R1 | R2 | R3 |
| Mollusca | Arcidae | <i>Anadara consociata</i> | 8 | 1 | 6 | 0 | 0 | 0 |
| Mollusca | Corbulidae | <i>Corbula scaphoides</i> | 0 | 1 | 0 | 0 | 0 | 0 |
| Mollusca | Tellinidae | <i>Nitidotellina lischkei</i> | 0 | 0 | 2 | 0 | 0 | 0 |
| Mollusca | / | Unidentified juvenile bivalve | 1 | 0 | 0 | 0 | 0 | 0 |
| Annelida | Glyceridae | <i>Glycera chirori</i> | 1 | 0 | 2 | 0 | 0 | 0 |
| Annelida | Eunicidae | <i>Marphysa stragulum</i> | 0 | 0 | 2 | 0 | 0 | 0 |
| Annelida | Orbiniidae | <i>Naineris</i> sp. | 0 | 0 | 1 | 0 | 0 | 0 |
| Annelida | Nereididae | <i>Nereididae</i> spp. | 0 | 0 | 1 | 0 | 0 | 0 |
| Annelida | Onuphidae | <i>Onuphis eremita</i> | 13 | 11 | 30 | 0 | 0 | 0 |
| Annelida | Pilargidae | <i>Sigambra hanaokai</i> | 1 | 0 | 1 | 0 | 0 | 0 |
| Annelida | Syllidae | <i>Syllis</i> sp. | 1 | 1 | 1 | 0 | 0 | 0 |

Appendix 9.6b Benthos Survey Results (Biomass) during Wet Season

| Phylum | Family | Species | B1 | | | B2 | | |
|----------|------------|-------------------------------|-------|-------|-------|----|----|----|
| | | | R1 | R2 | R3 | R1 | R2 | R3 |
| Mollusca | Arcidae | <i>Anadara consociata</i> | 5.86 | 0.94 | 11.28 | 0 | 0 | 0 |
| Mollusca | Corbulidae | <i>Corbula scaphoides</i> | 0 | 0.05 | 0 | 0 | 0 | 0 |
| Mollusca | Tellinidae | <i>Nitidotellina lischkei</i> | 0 | 0 | 0.25 | 0 | 0 | 0 |
| Mollusca | / | Unidentified juvenile bivalve | >0.01 | 0 | 0 | 0 | 0 | 0 |
| Annelida | Glyceridae | <i>Glycera chirori</i> | 0.03 | 0 | 0.38 | 0 | 0 | 0 |
| Annelida | Eunicidae | <i>Marphysa stragulum</i> | 0 | 0 | 0.08 | 0 | 0 | 0 |
| Annelida | Orbiniidae | <i>Naineris</i> sp. | 0 | 0 | 0.02 | 0 | 0 | 0 |
| Annelida | Nereididae | <i>Nereididae</i> spp. | 0 | 0 | >0.01 | 0 | 0 | 0 |
| Annelida | Onuphidae | <i>Onuphis eremita</i> | 1.42 | 1.48 | 2.49 | 0 | 0 | 0 |
| Annelida | Pilargidae | <i>Sigambra hanaokai</i> | >0.01 | 0 | >0.01 | 0 | 0 | 0 |
| Annelida | Syllidae | <i>Syllis</i> sp. | >0.01 | >0.01 | >0.01 | 0 | 0 | 0 |

Appendix 9.6c Benthos Survey Results (Abundance) during Dry Season

| Phylum | Family | Species | B1 | | | B2 | | |
|----------|------------|--------------------------|----|----|----|----|----|----|
| | | | R1 | R2 | R3 | R1 | R2 | R3 |
| Annelida | Eunicidae | <i>Eunice indica</i> | 1 | 2 | 0 | 0 | 0 | 0 |
| Annelida | Nereididae | <i>Nereididae</i> spp. | 0 | 1 | 0 | 0 | 0 | 0 |
| Annelida | Pilargidae | <i>Sigambra hanaokai</i> | 2 | 2 | 0 | 0 | 0 | 0 |

Appendix 9.6d Benthos Survey Results (Biomass) during Dry Season

| Phylum | Family | Species | B1 | | | B2 | | |
|----------|------------|--------------------------|-------|-------|----|----|----|----|
| | | | R1 | R2 | R3 | R1 | R2 | R3 |
| Annelida | Eunicidae | <i>Eunice indica</i> | >0.01 | 0.01 | 0 | 0 | 0 | 0 |
| Annelida | Nereididae | <i>Nereididae</i> spp. | 0 | >0.01 | 0 | 0 | 0 | 0 |
| Annelida | Pilargidae | <i>Sigambra hanaokai</i> | >0.01 | >0.01 | 0 | 0 | 0 | 0 |