



B (809800,826000)




500m Study Area

A (809800,824600)



Scale: As shown
Figure 7-C1 Cumulative 24-hour RSP Concentration ($\mu\text{g}/\text{m}^3$) at 8.5mPD emitted from Source Location A4 (Mitigated Scenario 2 Phase II Eco Park without CEDD Fill Bank)



B (809800,826000)




500m Study Area

A (809800,824600)



Scale: As shown

Figure 7-C2 Cumulative 24-hour RSP Concentration ($\mu\text{g}/\text{m}^3$) at 8.5mPD emitted from Source Location A6 (Mitigated Scenario 2 Phase II Eco Park without CEDD Fill Bank)



B (809800,826000)




500m Study Area

A (809800,824600)



Scale: As shown
 Figure 7-C3 Cumulative 24-hour RSP Concentration ($\mu\text{g}/\text{m}^3$) at 8.5mPD emitted from Source Location A8 (Mitigated Scenario 2 Phase II Eco Park without CEDD Fill Bank)



Scale: As shown

Figure B7-C4 Cumulative 24-hour RSP Concentration ($\mu\text{g}/\text{m}^3$) at 22.5mPD emitted from Source Location A3 (Mitigated Scenario 2 Phase II Eco Park without CEDD Fill Bank)



B (809800,826000)



500m Study Area

A (809800,824600)



Scale: As shown

Figure 7-C5 Cumulative 24-hour RSP Concentration ($\mu\text{g}/\text{m}^3$) at 61.5mPD emitted from Source Location A2 (Mitigated Scenario 2 Phase II Eco Park without CEDD Fill Bank)



B (809800,826000)



500m Study Area

A (809800,824600)



Scale: As shown
 Figure 7-C6 Cumulative 24-hour RSP Concentration ($\mu\text{g}/\text{m}^3$) at 61.5mPD emitted from Source Location A5 (Mitigated Scenario 2 Phase II Eco Park without CEDD Fill Bank)