

Pearl River Delta Regional Air Quality Monitoring Network

A Report of Monitoring Results for the Period between January and June 2007

Report Number : **PRDAIR-2007-1**

Report Prepared by : **Guangdong Provincial
Environmental Protection
Monitoring Centre**

**Environmental Protection
Department, HKSAR**

Approved by : **Pearl River Delta Air Quality
Management and Monitoring
Special Panel**

Security Classification : **Unrestricted**

Purpose of the Report

This report provides the monitoring results from the Pearl River Delta Regional Air Quality Monitoring Network measured between January and June 2007 and their statistical analysis.

Contents

	<u>Page</u>
1. Introduction to the Pearl River Delta Regional Air Quality Monitoring Network	1
2. Operation of the Network	2
2.1 Quality Control (QC) and Quality Assurance (QA) Activities	2
3. Statistical Analysis of Pollutant Concentrations	3
3.1 Sulphur Dioxide (SO₂)	3
3.2 Nitrogen Dioxide (NO₂)	5
3.3 Ozone (O₃)	7
3.4 Respirable Suspended Particulates (PM₁₀)	9
3.5 Monthly Variations of Pollutant Concentrations	11
4. Statistical Analysis of the Regional Air Quality Index (RAQI)	12
4.1 Statistics on RAQI Grades	13
4.2 Spatial Distribution of Average RAQI Grades	14
4.3 Monthly Variations of Average RAQI	15
Annex A : Site Information of Monitoring Stations	16
Annex B : Measurement Methods of Air Pollutant Concentration	17

List of Tables

	<u>Page</u>
Table 3.1 a : The monthly maxima and minima of hourly averages of Sulphur Dioxide	4
Table 3.1 b : The monthly maxima and minima of daily averages of Sulphur Dioxide	4
Table 3.1 c : The monthly averages and overall averages of Sulphur Dioxide	4
Table 3.2 a : The monthly maxima and minima of hourly averages of Nitrogen Dioxide	6
Table 3.2 b : The monthly maxima and minima of daily averages of Nitrogen Dioxide	6
Table 3.2 c : The monthly averages and overall averages of Nitrogen Dioxide	6
Table 3.3 a : The monthly maxima and minima of hourly averages of Ozone	8
Table 3.3 b : The monthly maxima and minima of daily averages of Ozone	8
Table 3.3 c : The monthly averages and overall averages of Ozone	8
Table 3.4 a : The monthly maxima and minima of hourly averages of Respirable Suspended Particulates	10
Table 3.4 b : The monthly maxima and minima of daily averages of Respirable Suspended Particulates	10
Table 3.4 c : The monthly averages and overall averages of Respirable Suspended Particulates	10
Table 4.1 a : Statistics on RAQI grades of individual monitoring stations	13

List of Figures

	<u>Page</u>
Figure 1 : Spatial distribution of the PRD Regional Air Quality Monitoring Stations	1
Figure 2 : Spatial distribution of average concentrations of Sulphur Dioxide (SO₂) in the Network	3
Figure 3 : Spatial distribution of average concentrations of Nitrogen Dioxide (NO₂) in the Network	5
Figure 4 : Spatial distribution of average concentrations of Ozone (O₃) in the Network	7
Figure 5 : Spatial distribution of average concentrations of Respirable Suspended Particulates (PM₁₀) in the Network	9
Figure 6 : Monthly variations of average pollutant concentrations measured by the Network	11
Figure 7 : Stacked column chart of RAQI grades of individual monitoring stations	13
Figure 8 : Distribution of RAQI grades in the Network	14
Figure 9 : Spatial distribution of average RAQI grades in the Network	14
Figure 10 : The average RAQI of individual monitoring stations	15
Figure 11 : Monthly variations in average RAQI	15

1. Introduction to the Pearl River Delta Regional Air Quality Monitoring Network

The Pearl River Delta (PRD) Regional Air Quality Monitoring Network (the Network) was jointly established by the Guangdong Provincial Environmental Protection Monitoring Centre (GDEMC) and the Environmental Protection Department of the HKSAR (HKEPD) from 2003 to 2005. It came into operation on 30 November 2005 and has been providing data for reporting of Regional Air Quality Index (RAQI) to the public since then.

The Network comprises 16 automatic air quality monitoring stations (see Figure 1) across the PRD region. Ten of these stations are operated by the Environmental Protection Monitoring Centres of the individual cities in Guangdong while the 3 stations located in Hong Kong are managed by the HKEPD. The remaining 3 regional stations in the Network are operated by the GDEMC. The objectives of the Network are to :

- provide accurate air quality data that can help the Guangdong Provincial and HKSAR governments to appraise the air quality situation and pollution problems in the PRD region for formulating appropriate control measures;
- evaluate the effectiveness of the air pollution control measures through long-term monitoring;
- provide the public with information on the air quality of various places in the region.

In order to ensure the air quality monitoring results attain a high degree of accuracy and reliability, the two governments had jointly developed a set of “Standard Operational Procedures on Quality Assurance and Quality Control of the PRD Air Quality Monitoring System for Hong Kong and Guangdong” (QA/QC Operating Procedures). The design and operation of the Network comply with the requirements set out in the QA/QC Operating Procedures.

All stations are installed with equipment to measure the ambient concentrations of respirable suspended particulate (PM₁₀ or RSP), sulphur dioxide (SO₂), nitrogen dioxide (NO₂) and ozone (O₃).

Annexes A and B set out, respectively, the site information of the monitoring stations in the Network and the methods used for measuring air pollutant concentrations.




Figure 1 : Spatial distribution of the PRD Regional Air Quality Monitoring Stations

2. Operation of the Network

The Network was generally in smooth operation during the period from January to June 2007. The data capture rates of all monitoring stations in the Network averaged 90%.

In order to provide the public in both Guangdong and HKSAR with daily air quality information in different parts of the PRD region, the GDEMC and HKEPD established a daily reporting system of the Regional Air Quality Index (RAQI). The two Governments have been issuing the RAQI to the public at 4pm every day through the Internet since 30 November 2005.

2.1 Quality Control (QC) and Quality Assurance (QA) Activities

The two governments have fully carried out the agreed QA/QC activities, which include zero/span checks, precision checks, dynamic calibration, etc., in accordance with the QA/QC Operating Procedures so as to ensure that the air quality data from the monitoring stations are highly accurate and reliable. To ensure the operation of the Network complies continuously with the QA/QC requirements, the GDEMC and HKEPD have jointly set up the Guangdong-Hong Kong Quality Management Committee for the PRD Regional Air Quality Monitoring Network (the Quality Management Committee, QMC) to review, on a quarterly basis, the set-up of the network, its performance in QA/QC and the operation status of its data transmission system. The QMC will also conduct system audit once a year to evaluate the effectiveness of the quality management system. The findings of the system audit will be reported. The deficiencies found and corrective measures suggested will be listed and followed up by the QMC.

The performance audit results of Network accuracy and precision for 2007 will be discussed in details in the annual report.

3. Statistical Analysis of Pollutant Concentrations

3.1 Sulphur Dioxide (SO₂)

SO₂ comes mainly from the combustion of sulphur-containing fossil fuel. Its major sources of emissions include power plants, fuel combustion plants, vehicles and vessels. Apart from its impact on human respiratory system, SO₂ contributes substantially to acid rain. It can also be oxidized in the air to form sulphate which has a significant impact on the levels of respirable suspended particulates (PM₁₀) and visibility in the region.

The overall averages of SO₂ at various monitoring stations in the Network ranged from 0.016 mg/m³ to 0.127 mg/m³ for the period from January to June 2007. Figure 2 shows that the average levels of SO₂ at the north-western part of PRD and the Pearl River Estuary region were in general higher than those of other areas. The overall averages of SO₂ at various monitoring stations are shown in Table 3.1c.

During the period, 9 monitoring stations in the Network had recorded exceedance of the national daily air quality standard[#] (0.15 mg/m³) of SO₂ while the corresponding national hourly standard (0.50 mg/m³) was exceeded at 5 monitoring stations. Details are shown in Figure 2 and Table 3.1a - 3.1b.




Figure 2 : Spatial distribution of average concentrations of Sulphur Dioxide (SO₂) in the Network

National Standards refer to Class 2 of the “National Ambient Air Quality Standards (GB 3095 – 1996 – revised version)” [NAAQS], which are applicable to residential, mixed commercial/residential, cultural, industrial and village areas.

Table 3.1 a : The monthly maxima and minima of hourly averages of Sulphur Dioxide
[Class 2 NAAQS (Hourly) : 0.50 mg/m³]

Month	Jan 2007		Feb		Mar		Apr		May		Jun		Exceed- ance Hours	Exceed- ance Rate
	Min	Max	Min	Max	Min	Max	Min	Max	Min	Max	Min	Max		
Luhu Park(Guangzhou)	0.008	0.368	0.003	0.449	0.002	0.510	0.004	0.206	0.003	0.306	0.003	0.356	1	0.02%
Wanqingsha(Guangzhou)	0.022	0.340	0.009	0.466	0.009	0.297	0.009	0.222	0.008	0.346	0.009	0.473	0	0.00%
Tianhu(Guangzhou)	0.006	0.083	0.006	0.258	0.006	0.190	0.007	0.374	0.008	0.370	0.008	0.260	0	0.00%
Liyuan(Shenzhen)	0.014	0.197	0.011	0.148	0.002	0.108	0.000	0.180	0.001	0.188	0.001	0.198	0	0.00%
Tangjia(Zhuhai)	0.020	0.331	0.012	0.319	0.011	0.349	0.000	0.278	0.000	0.568	0.000	0.199	1	0.03%
Shunde Dangxiao(Foshan)	0.023	0.514	0.001	0.604	0.014	0.370	0.004	0.338	0.003	0.581	0.006	0.232	6	0.16%
Huijingcheng(Foshan)	0.047	0.891	0.009	0.593	0.027	0.547	0.019	0.456	0.016	0.675	0.012	0.627	41	1.02%
Donghu(Jiangmen)	0.022	0.279	0.003	0.338	0.005	0.240	0.001	0.209	0.000	0.117	0.000	0.095	0	0.00%
Chengzhong(Zhaoqing)	0.013	0.380	0.007	0.324	0.006	0.405	0.005	0.245	0.008	0.297	0.005	0.260	0	0.00%
Xiapu(Huizhou)	0.000	0.142	0.000	0.075	0.000	0.078	0.000	0.181	0.000	0.083	0.000	0.069	0	0.00%
Jinguowan(Huizhou)	0.006	0.074	0.005	0.067	0.005	0.118	0.005	0.126	0.006	0.099	0.006	0.075	0	0.00%
Haogang(Dongguan)	0.017	0.495	0.005	0.326	0.004	0.414	0.003	0.439	0.004	0.362	0.007	0.231	0	0.00%
Zimaling Park(Zhongshan)	0.015	0.539	0.004	0.356	0.004	0.409	0.004	0.251	0.004	0.256	0.005	0.153	1	0.04%
Tsuen Wan(HKSAR)	0.005	0.167	0.005	0.126	0.005	0.225	0.005	0.200	0.006	0.259	0.006	0.149	0	0.00%
Tap Mun(HKSAR)	0.009	0.257	0.004	0.069	0.004	0.223	0.005	0.106	0.005	0.132	0.003	0.175	0	0.00%
Tung Chung(HKSAR)	0.008	0.177	0.000	0.115	0.000	0.186	0.002	0.198	0.001	0.227	0.010	0.114	0	0.00%

Table 3.1 b : The monthly maxima and minima of daily averages of Sulphur Dioxide
[Class 2 NAAQS (Daily) : 0.15 mg/m³]

Month	Jan 2007		Feb		Mar		Apr		May		Jun		Exceed- ance Days	Exceed- ance Rate
	Min	Max	Min	Max	Min	Max	Min	Max	Min	Max	Min	Max		
Luhu Park(Guangzhou)	0.017	0.150	0.006	0.147	0.006	0.154	0.010	0.112	0.008	0.110	0.040	0.151	3	1.70%
Wanqingsha(Guangzhou)	0.040	0.212	0.011	0.135	0.012	0.163	0.012	0.120	0.016	0.118	0.012	0.096	5	2.96%
Tianhu(Guangzhou)	0.006	0.039	0.006	0.098	0.007	0.108	0.007	0.204	0.008	0.176	0.011	0.139	2	1.15%
Liyuan(Shenzhen)	0.023	0.078	0.024	0.093	0.005	0.070	0.004	0.063	0.007	0.069	0.004	0.042	0	0.00%
Tangjia(Zhuhai)	0.098	0.201	0.019	0.140	0.014	0.184	0.003	0.155	0.003	0.137	0.002	0.060	10	6.58%
Shunde Dangxiao(Foshan)	0.043	0.285	0.007	0.232	0.036	0.187	0.027	0.177	0.028	0.255	0.036	0.109	17	10.63%
Huijingcheng(Foshan)	0.077	0.436	0.021	0.319	0.043	0.362	0.043	0.227	0.049	0.267	0.034	0.281	46	26.90%
Donghu(Jiangmen)	0.039	0.127	0.005	0.133	0.013	0.109	0.006	0.090	0.004	0.045	0.002	0.029	0	0.00%
Chengzhong(Zhaoqing)	0.051	0.212	0.014	0.175	0.016	0.230	0.021	0.139	0.018	0.165	0.012	0.105	14	7.95%
Xiapu(Huizhou)	0.002	0.052	0.005	0.048	0.005	0.048	0.009	0.077	0.001	0.039	0.005	0.030	0	0.00%
Jinguowan(Huizhou)	0.009	0.033	0.006	0.045	0.007	0.039	0.007	0.065	0.007	0.033	0.006	0.032	0	0.00%
Haogang(Dongguan)	0.038	0.173	0.008	0.132	0.017	0.115	0.010	0.175	0.011	0.135	0.021	0.068	2	1.22%
Zimaling Park(Zhongshan)	0.037	0.219	0.007	0.170	0.006	0.137	0.007	0.146	0.004	0.095	0.008	0.028	11	9.73%
Tsuen Wan(HKSAR)	0.012	0.065	0.007	0.054	0.008	0.069	0.008	0.073	0.010	0.080	0.011	0.064	0	0.00%
Tap Mun(HKSAR)	0.016	0.056	0.006	0.043	0.007	0.029	0.007	0.035	0.007	0.036	0.003	0.021	0	0.00%
Tung Chung(HKSAR)	0.020	0.092	0.003	0.070	0.000	0.067	0.003	0.073	0.003	0.065	0.011	0.036	0	0.00%

Table 3.1 c : The monthly averages and overall averages of Sulphur Dioxide

Month	Jan 2007	Feb	Mar	Apr	May	Jun	Overall Average
Luhu Park(Guangzhou)	0.049	0.075	0.059	0.049	0.050	0.076	0.059
Wanqingsha(Guangzhou)	0.107	0.039	0.055	0.065	0.055	0.031	0.060
Tianhu(Guangzhou)	0.024	0.039	0.027	0.040	0.039	0.049	0.036
Liyuan(Shenzhen)	0.049	0.054	0.041	0.021	0.021	0.013	0.033
Tangjia(Zhuhai)	0.154*	0.047	0.060	0.059	0.033	0.011	0.049
Shunde Dangxiao(Foshan)	0.147*	0.091	0.094	0.094	0.090	0.068	0.095
Huijingcheng(Foshan)	0.179	0.104	0.137	0.128	0.115	0.108	0.127
Donghu(Jiangmen)	0.074	0.037	0.044	0.036	0.021	0.012	0.036
Chengzhong(Zhaoqing)	0.113	0.070	0.089	0.060	0.065	0.036	0.072
Xiapu(Huizhou)	0.025	0.020	0.024	0.027	0.019	0.019*	0.022
Jinguowan(Huizhou)	0.020*	0.016	0.014	0.018	0.015	0.013	0.016
Haogang(Dongguan)	0.074	0.054	0.043	0.052	0.052	0.042	0.053
Zimaling Park(Zhongshan)	0.131	0.049*	0.048	0.063	0.028*	0.016*	0.071
Tsuen Wan(HKSAR)	0.031	0.023	0.022	0.023	0.033	0.026	0.026
Tap Mun(HKSAR)	0.030	0.014	0.013	0.017	0.015	0.010	0.017
Tung Chung(HKSAR)	0.045	0.019	0.014	0.020	0.022	0.018	0.023

Remark : 1. All concentration units are in milligrams per cubic metre.

2. "*" denotes that the data capture rate does not meet the minimum requirements for determining a representative value.

3.2 Nitrogen Dioxide (NO₂)

Nitrogen Dioxide (NO₂) is mainly formed from oxidization of nitrogen monoxide (NO) emitted in the process of combustion. Its major emission sources include power plants, vehicles, industrial combustion plants, etc. Apart from the impact on human respiratory system, it can also be oxidized in the air to form nitrate, which has significant impact on the levels of particulates, acid rain and visibility in the region.

The overall averages of NO₂ at various monitoring stations in the Network ranged from 0.016 mg/m³ to 0.074 mg/m³ for the period from January to June 2007. During the period, 10 monitoring stations in the Network had recorded exceedance of the national daily air quality standard (0.12 mg/m³) of NO₂ while the corresponding national hourly standard (0.24 mg/m³) was exceeded at 9 monitoring stations. Details are shown in Figure 3 and Table 3.2a - 3.2c.




Figure 3 : Spatial distribution of average concentrations of Nitrogen Dioxide (NO₂) in the Network

Table 3.2 a : The monthly maxima and minima of hourly averages of Nitrogen Dioxide
[Class 2 NAAQS (Hourly) : 0.24 mg/m³]

Month	Jan 2007		Feb		Mar		Apr		May		Jun		Exceed- ance Hours	Exceed- ance Rate
	Min	Max	Min	Max	Min	Max	Min	Max	Min	Max	Min	Max		
Luhu Park(Guangzhou)	0.016	0.271	0.015	0.267	0.024	0.245	0.000	0.335	0.014	0.276	0.011	0.167	23	0.55%
Wanqingsha(Guangzhou)	0.030	0.192	0.003	0.182	0.005	0.159	0.003	0.152	0.000	0.207	0.000	0.129	0	0.00%
Tianhu(Guangzhou)	0.004	0.061	0.003	0.150	0.002	0.114	0.002	0.125	0.002	0.116	0.001	0.118	0	0.00%
Liyuan(Shenzhen)	0.005	0.299	0.003	0.158	0.003	0.220	0.000	0.210	0.008	0.225	0.010	0.144	1	0.02%
Tangjia(Zhuhai)	0.005	0.144	0.000	0.190	0.000	0.184	0.001	0.122	0.004	0.225	0.002	0.157	0	0.00%
Shunde Dangxiao(Foshan)	0.017	0.228	0.012	0.187	0.016	0.283	0.009	0.150	0.014	0.242	0.006	0.111	4	0.11%
Huijingcheng(Foshan)	0.026	0.323	0.000	0.306	0.016	0.217	0.016	0.212	0.023	0.184	0.016	0.209	19	0.48%
Donghu(Jiangmen)	0.016	0.183	0.007	0.202	0.002	0.147	0.001	0.279	0.004	0.182	0.001	0.101	7	0.18%
Chengzhong(Zhaoqing)	0.016	0.230	0.005	0.174	0.012	0.157	0.008	0.133	0.020	0.098	0.009	0.094	0	0.00%
Xiapu(Huizhou)	0.002	0.168	0.000	0.197	0.001	0.152	0.000	0.171	0.000	0.287	0.000	0.072	4	0.10%
Jinguowan(Huizhou)	0.005	0.122	0.004	0.106	0.004	0.096	0.005	0.089	0.004	0.113	0.002	0.075	0	0.00%
Haogang(Dongguan)	0.009	0.148	0.008	0.166	0.011	0.199	0.009	0.228	0.015	0.178	0.014	0.165	0	0.00%
Zimaling Park(Zhongshan)	0.028	0.331	0.010	0.172	0.006	0.186	0.005	0.160	0.005	0.166	0.013	0.111	3	0.10%
Tsuen Wan(HKSAR)	0.016	0.241	0.020	0.185	0.017	0.186	0.015	0.211	0.015	0.214	0.015	0.189	1	0.02%
Tap Mun(HKSAR)	0.006	0.117	0.005	0.077	0.003	0.067	0.004	0.090	0.002	0.116	0.002	0.049	0	0.00%
Tung Chung(HKSAR)	0.013	0.271	0.006	0.180	0.004	0.205	0.003	0.157	0.006	0.174	0.001	0.154	2	0.05%

Table 3.2 b : The monthly maxima and minima of daily averages of Nitrogen Dioxide
[Class 2 NAAQS (Daily) : 0.12 mg/m³]

Month	Jan 2007		Feb		Mar		Apr		May		Jun		Exceed- ance Days	Exceed- ance Rate
	Min	Max	Min	Max	Min	Max	Min	Max	Min	Max	Min	Max		
Luhu Park(Guangzhou)	0.031	0.178	0.028	0.179	0.036	0.138	0.039	0.199	0.039	0.157	0.030	0.116	15	8.57%
Wanqingsha(Guangzhou)	0.049	0.125	0.016	0.094	0.014	0.098	0.012	0.098	0.011	0.084	0.003	0.057	2	1.27%
Tianhu(Guangzhou)	0.006	0.024	0.006	0.070	0.004	0.057	0.003	0.067	0.003	0.058	0.004	0.050	0	0.00%
Liyuan(Shenzhen)	0.022	0.109	0.006	0.070	0.007	0.053	0.031	0.124	0.029	0.090	0.027	0.093	1	0.59%
Tangjia(Zhuhai)	0.052	0.096	0.013	0.068	0.010	0.097	0.006	0.072	0.009	0.074	0.007	0.066	0	0.00%
Shunde Dangxiao(Foshan)	0.028	0.156	0.022	0.131	0.028	0.116	0.026	0.101	0.026	0.123	0.021	0.061	5	3.16%
Huijingcheng(Foshan)	0.048	0.234	0.019	0.170	0.031	0.104	0.027	0.120	0.048	0.109	0.029	0.095	10	5.95%
Donghu(Jiangmen)	0.029	0.108	0.013	0.130	0.009	0.063	0.008	0.171	0.008	0.091	0.007	0.049	4	2.33%
Chengzhong(Zhaoqing)	0.029	0.114	0.010	0.104	0.022	0.112	0.017	0.068	0.027	0.058	0.019	0.050	0	0.00%
Xiapu(Huizhou)	0.014	0.073	0.016	0.078	0.017	0.073	0.019	0.074	0.005	0.055	0.010	0.041	0	0.00%
Jinguowan(Huizhou)	0.007	0.032	0.007	0.042	0.010	0.038	0.009	0.053	0.009	0.031	0.006	0.032	0	0.00%
Haogang(Dongguan)	0.027	0.094	0.021	0.101	0.025	0.099	0.023	0.142	0.039	0.089	0.036	0.092	1	0.61%
Zimaling Park(Zhongshan)	0.042	0.140	0.022	0.114	0.014	0.086	0.011	0.082	0.011	0.072	0.025	0.040	4	3.28%
Tsuen Wan(HKSAR)	0.052	0.147	0.052	0.118	0.044	0.099	0.046	0.131	0.049	0.128	0.030	0.100	6	3.39%
Tap Mun(HKSAR)	0.011	0.043	0.006	0.035	0.007	0.027	0.006	0.043	0.004	0.042	0.006	0.028	0	0.00%
Tung Chung(HKSAR)	0.043	0.139	0.025	0.102	0.011	0.122	0.008	0.110	0.015	0.108	0.010	0.075	3	1.69%

Table 3.2 c : The monthly averages and overall averages of Nitrogen Dioxide

Month	Jan 2007	Feb	Mar	Apr	May	Jun	Overall Average
Luhu Park(Guangzhou)	0.071	0.087	0.066	0.091	0.072	0.060	0.074
Wanqingsha(Guangzhou)	0.079	0.044	0.046	0.050	0.044	0.021	0.049
Tianhu(Guangzhou)	0.011	0.026	0.019	0.021	0.019	0.025	0.020
Liyuan(Shenzhen)	0.042	0.023	0.021	0.059	0.055	0.043	0.040
Tangjia(Zhuhai)	0.070*	0.033	0.040	0.037	0.033	0.015	0.033
Shunde Dangxiao(Foshan)	0.080*	0.064	0.061	0.061	0.063	0.037	0.060
Huijingcheng(Foshan)	0.096	0.075	0.058	0.071	0.073	0.059	0.071
Donghu(Jiangmen)	0.057	0.036	0.026	0.051	0.041	0.024	0.039
Chengzhong(Zhaoqing)	0.060	0.051	0.047	0.036	0.038	0.030	0.043
Xiapu(Huizhou)	0.033	0.041	0.037	0.040	0.025	0.023*	0.033
Jinguowan(Huizhou)	0.016*	0.019	0.020	0.022*	0.018	0.019	0.019
Haogang(Dongguan)	0.055	0.056	0.055	0.065	0.060	0.059	0.059
Zimaling Park(Zhongshan)	0.085	0.051*	0.042	0.046	0.036	0.034*	0.054
Tsuen Wan(HKSAR)	0.086	0.078	0.069	0.077	0.073	0.050	0.072
Tap Mun(HKSAR)	0.023	0.014	0.015	0.018	0.015	0.013	0.016
Tung Chung(HKSAR)	0.079	0.055	0.047	0.056	0.045	0.023	0.051

Remark : 1. All concentration units are in milligrams per cubic metre.

2. “*” denotes that the data capture rate does not meet the minimum requirements for determining a representative value.

3.3 Ozone (O₃)

Ozone (O₃) is not directly emitted from emission sources. It is formed by the photochemical reaction of oxygen, nitrogen oxides (NO_x) and volatile organic compounds (VOCs) in the air under sunlight, and is the main component of photochemical smog. Ozone can cause irritation to the eye, nose and throat. At elevated levels, O₃ can increase a person's susceptibility to respiratory diseases and aggravate pre-existing respiratory diseases such as asthma.

The precursors (NO_x and VOCs) of O₃ mainly originate from pollution sources in urban areas. However, as it usually takes several hours for O₃ to be formed and rise to its peak level, and O₃ and its precursors can be transported to rural areas downwind of their sources during this period, the concentrations of O₃ in rural areas are therefore often higher than in the urban areas. The overall averages of O₃ recorded by the Network ranged from 0.026 mg/m³ to 0.075 mg/m³ for the period from January to June 2007, with higher average values measured in rural areas such as Tianhu of Guangzhou, Tap Mun of Hong Kong and Jinguowan of Huizhou. During the period, 15 monitoring stations in the Network had recorded exceedance of the national hourly standard (0.20 mg/m³) of ozone. Details are shown in Figure 4 and Table 3.3a - 3.3c.




Figure 4 : Spatial distribution of average concentrations of Ozone (O₃) in the Network

Table 3.3 a : The monthly maxima and minima of hourly averages of Ozone**[Class 2 NAAQS (Hourly) : 0.20 mg/m³]**

Month	Jan 2007		Feb		Mar		Apr		May		Jun		Exceedance Hours	Exceedance Rate
	Min	Max	Min	Max	Min	Max	Min	Max	Min	Max	Min	Max		
Luhu Park(Guangzhou)	0.000	0.126	0.000	0.120	0.000	0.123	0.000	0.239	0.002	0.312	0.004	0.239	54	1.35%
Wanqingsha(Guangzhou)	0.000	0.205	0.002	0.142	0.000	0.196	0.001	0.374	0.001	0.402	0.005	0.367	80	1.97%
Tianhu(Guangzhou)	0.017	0.165	0.002	0.232	0.001	0.202	0.001	0.364	0.000	0.303	0.000	0.276	86	2.14%
Liyuan(Shenzhen)	0.015	0.203	0.004	0.222	0.005	0.164	0.000	0.088	0.000	0.142	0.000	0.067	2	0.05%
Tangjia(Zhuhai)	0.008	0.177	0.000	0.187	0.000	0.143	0.000	0.204	0.001	0.382	0.000	0.183	28	0.81%
Shunde Dangxiao(Foshan)	0.000	0.199	0.000	0.161	0.000	0.111	0.000	0.289	0.000	0.385	0.000	0.238	63	1.57%
Huijingcheng(Foshan)	0.002	0.132	0.000	0.198	0.001	0.108	0.000	0.370	0.001	0.285	0.002	0.226	30	0.77%
Donghu(Jiangmen)	0.002	0.219	0.003	0.267	0.001	0.145	0.003	0.261	0.001	0.330	0.002	0.257	55	1.36%
Chengzhong(Zhaoqing)	0.003	0.164	0.002	0.164	0.002	0.092	0.002	0.213	0.003	0.293	0.002	0.179	26	0.64%
Xiapu(Huizhou)	0.001	0.183	0.000	0.169	0.000	0.119	0.003	0.303	0.010	0.253	0.009	0.264	24	0.64%
Jinguowan(Huizhou)	0.011	0.172	0.003	0.198	0.003	0.143	0.003	0.358	0.004	0.267	0.003	0.244	43	1.09%
Haogang(Dongguan)	0.002	0.175	0.001	0.218	0.003	0.226	0.001	0.352	0.002	0.342	0.003	0.350	90	2.27%
Zimaling Park(Zhongshan)	0.000	0.228	0.000	0.181	0.000	0.116	0.000	0.255	0.000	0.290	0.000	0.245	21	0.76%
Tsuen Wan(HKSAR)	0.002	0.103	0.003	0.112	0.004	0.093	0.005	0.134	0.004	0.175	0.004	0.074	0	0.00%
Tap Mun(HKSAR)	0.006	0.191	0.005	0.171	0.002	0.183	0.002	0.226	0.002	0.303	0.000	0.160	15	0.36%
Tung Chung(HKSAR)	0.000	0.208	0.001	0.132	0.000	0.093	0.000	0.152	0.001	0.289	0.007	0.100	13	0.31%

Table 3.3 b : The monthly maxima and minima of daily averages of Ozone

Month	Jan 2007		Feb		Mar		Apr		May		Jun	
	Min	Max	Min	Max	Min	Max	Min	Max	Min	Max	Min	Max
Luhu Park(Guangzhou)	0.004	0.047	0.003	0.056	0.001	0.051	0.001	0.070	0.007	0.094	0.014	0.081
Wanqingsha(Guangzhou)	0.008	0.064	0.006	0.094	0.002	0.055	0.004	0.137	0.029	0.138	0.021	0.124
Tianhu(Guangzhou)	0.033	0.132	0.019	0.134	0.010	0.115	0.017	0.154	0.053	0.152	0.025	0.099
Liyuan(Shenzhen)	0.039	0.105	0.023	0.101	0.020	0.084	0.005	0.051	0.002	0.053	0.000	0.017
Tangjia(Zhuhai)	0.010	0.078	0.014	0.116	0.006	0.100	0.004	0.111	0.034	0.154	0.024	0.085
Shunde Dangxiao(Foshan)	0.000	0.055	0.004	0.061	0.001	0.052	0.001	0.114	0.010	0.125	0.000	0.081
Huijingcheng(Foshan)	0.003	0.053	0.003	0.058	0.006	0.055	0.006	0.148	0.005	0.081	0.010	0.091
Donghu(Jiangmen)	0.003	0.058	0.011	0.100	0.004	0.060	0.005	0.114	0.013	0.178	0.011	0.081
Chengzhong(Zhaoqing)	0.011	0.057	0.005	0.057	0.004	0.063	0.012	0.121	0.007	0.128	0.017	0.094
Xiapu(Huizhou)	0.015	0.096	0.023	0.102	0.011	0.069	0.023	0.137	0.041	0.110	0.024	0.076
Jinguowan(Huizhou)	0.029	0.120	0.033	0.132	0.013	0.096	0.036	0.185	0.042	0.146	0.021	0.078
Haogang(Dongguan)	0.018	0.085	0.016	0.077	0.009	0.070	0.010	0.113	0.016	0.127	0.013	0.115
Zimaling Park(Zhongshan)	0.002	0.055	0.008	0.071	0.001	0.072	0.000	0.103	0.015	0.078	0.019	0.076
Tsuen Wan(HKSAR)	0.006	0.059	0.006	0.058	0.005	0.054	0.009	0.080	0.007	0.073	0.006	0.020
Tap Mun(HKSAR)	0.020	0.105	0.027	0.135	0.017	0.102	0.032	0.168	0.027	0.146	0.017	0.058
Tung Chung(HKSAR)	0.002	0.081	0.005	0.096	0.002	0.075	0.003	0.096	0.014	0.107	0.016	0.059

Table 3.3 c : The monthly averages and overall averages of Ozone

Month	Jan 2007	Feb	Mar	Apr	May	Jun	Overall Average
Luhu Park(Guangzhou)	0.023	0.017	0.013	0.027	0.051	0.045	0.030
Wanqingsha(Guangzhou)	0.027	0.025	0.019	0.061	0.080	0.047	0.042
Tianhu(Guangzhou)	0.082	0.081	0.059	0.078	0.091	0.058	0.075
Liyuan(Shenzhen)	0.066	0.056	0.045	0.026	0.027	0.004	0.037
Tangjia(Zhuhai)	0.036*	0.059	0.032	0.055	0.075	0.045	0.053
Shunde Dangxiao(Foshan)	0.021*	0.026	0.015	0.037	0.053	0.018	0.029
Huijingcheng(Foshan)	0.024*	0.028	0.017	0.045	0.035	0.033	0.031
Donghu(Jiangmen)	0.029	0.046	0.023	0.047	0.053	0.024	0.037
Chengzhong(Zhaoqing)	0.033	0.031	0.020	0.048	0.062	0.039	0.039
Xiapu(Huizhou)	0.051	0.050	0.034	0.060	0.070	0.048*	0.051
Jinguowan(Huizhou)	0.071*	0.080	0.054	0.081	0.082	0.044	0.068
Haogang(Dongguan)	0.056	0.043	0.030	0.051	0.065	0.042	0.048
Zimaling Park(Zhongshan)	0.025	0.039*	0.028	0.046*	0.036*	0.040*	0.034
Tsuen Wan(HKSAR)	0.027	0.030	0.025	0.036	0.031	0.011	0.026
Tap Mun(HKSAR)	0.072	0.084	0.065	0.075	0.078	0.031	0.068
Tung Chung(HKSAR)	0.030	0.042	0.032	0.039	0.051	0.030	0.037

Remark : 1. All concentration units are in milligrams per cubic metre.

2. "*" denotes that the data capture rate does not meet the minimum requirements for determining a representative value.

3.4 Respirable Suspended Particulates (PM₁₀)

The respirable suspended particulates (PM₁₀ or RSP) in the atmosphere come from a great variety of emission sources, such as power plants, vehicles, cement and pottery manufacturing, fugitive dust, etc, while some are products of oxidization of gaseous pollutants in the air (e.g., sulphate formed from oxidation of SO₂) or from photochemical reactions. PM₁₀ can penetrate deeply into human lungs and cause impact on human respiratory system. Furthermore, finer particles in PM₁₀ have significant effect on visibility.

The overall averages of PM₁₀ at various monitoring stations in the Network ranged from 0.037 mg/m³ to 0.139 mg/m³ for the period from January to June 2007. As shown in Figure 5, the average levels of PM₁₀ in the central and northern parts of PRD were generally higher than those in the coastal areas in the south. During the period, 11 monitoring stations had recorded exceedance of the national daily standard (0.15mg/m³) of PM₁₀. Details are shown in Tables 3.4a - 3.4c.

Distribution of average PM10
from Jan 2007 to Jun 2007




Figure 5 : Spatial distribution of average concentrations of Respirable Suspended Particulates (PM₁₀) in the Network

Table 3.4 a : The monthly maxima and minima of hourly averages of Respirable Suspended Particulates

Month	Jan 2007		Feb		Mar		Apr		May		Jun	
	Min	Max	Min	Max	Min	Max	Min	Max	Min	Max	Min	Max
Luhu Park(Guangzhou)	0.007	0.385	0.000	0.307	0.000	0.273	0.008	0.294	0.002	0.187	0.001	0.249
Wanqingsha(Guangzhou)	0.017	0.376	0.000	0.311	0.000	0.459	0.000	0.243	0.000	0.331	0.000	0.214
Tianhu(Guangzhou)	0.000	0.235	0.000	0.225	0.000	0.150	0.000	0.228	0.000	0.298	0.000	0.204
Liyuan(Shenzhen)	0.003	0.426	0.000	0.311	0.000	0.330	0.000	0.176	0.000	0.269	0.002	0.160
Tangjia(Zhuhai)	0.002	0.116	0.000	0.180	0.002	0.100	0.000	0.223	0.000	0.312	0.000	0.122
Shunde Dangxiao(Foshan)	0.021	0.536	0.005	0.480	0.006	0.546	0.015	0.291	0.000	0.559	0.008	0.248
Huijingcheng(Foshan)	0.016	0.862	0.011	0.626	0.007	0.916	0.024	0.516	0.007	0.607	0.007	0.872
Donghu(Jiangmen)	0.000	0.312	0.000	0.423	0.001	0.224	0.000	0.302	0.000	0.283	0.000	0.245
Chengzhong(Zhaoqing)	0.006	0.304	0.001	0.250	0.001	0.252	0.001	0.221	0.004	0.192	0.000	0.173
Xiapu(Huizhou)	0.051	0.382	0.041	0.379	0.004	0.423	0.000	0.381	0.007	0.479	--	--
Jinguowan(Huizhou)	0.000	0.192	0.002	0.163	0.000	0.239	0.000	0.223	0.000	0.216	0.000	0.120
Haogang(Dongguan)	0.018	0.533	0.006	0.412	0.003	0.347	0.010	0.321	0.000	0.274	0.000	0.240
Zimaling Park(Zhongshan)	0.011	0.234	0.004	0.181	0.003	0.202	0.002	0.161	0.003	0.175	0.002	0.201
Tsuen Wan(HKSAR)	0.025	0.218	0.016	0.179	0.008	0.330	0.013	0.173	0.010	0.217	0.009	0.098
Tap Mun(HKSAR)	0.021	0.167	0.015	0.176	0.006	0.143	0.010	0.122	0.007	0.168	0.004	0.075
Tung Chung(HKSAR)	0.025	0.257	0.012	0.195	0.006	0.343	0.006	0.172	0.006	0.213	0.003	0.082

Table 3.4 b : The monthly maxima and minima of daily averages of Respirable Suspended Particulates**[Class 2 NAAQS (Daily) : 0.15 mg/m³]**

Month	Jan 2007		Feb		Mar		Apr		May		Jun		Exceed- ance Days	Exceed- ance Rate
	Min	Max	Min	Max	Min	Max	Min	Max	Min	Max	Min	Max		
Luhu Park(Guangzhou)	0.035	0.269	0.027	0.228	0.011	0.179	0.021	0.255	0.021	0.121	0.021	0.129	12	7.02%
Wanqingsha(Guangzhou)	0.053	0.236	0.038	0.165	0.024	0.311	0.025	0.133	0.030	0.170	0.011	0.071	14	9.27%
Tianhu(Guangzhou)	0.006	0.133	0.018	0.134	0.009	0.087	0.015	0.173	0.011	0.186	0.019	0.103	3	1.72%
Liyuan(Shenzhen)	0.047	0.183	0.023	0.136	0.021	0.092	0.024	0.112	0.020	0.157	0.018	0.069	3	1.75%
Tangjia(Zhuhai)	0.027	0.058	0.024	0.056	0.022	0.049	0.021	0.105	0.019	0.105	0.010	0.045	0	0.00%
Shunde Dangxiao(Foshan)	0.055	0.355	0.053	0.265	0.039	0.267	0.047	0.204	0.034	0.256	0.022	0.124	29	18.01%
Huijingcheng(Foshan)	0.057	0.512	0.074	0.368	0.051	0.552	0.031	0.293	0.026	0.291	0.023	0.276	57	34.76%
Donghu(Jiangmen)	0.033	0.156	0.028	0.102	0.023	0.115	0.022	0.143	0.022	0.154	0.013	0.089	2	1.18%
Chengzhong(Zhaoqing)	0.016	0.186	0.034	0.170	0.014	0.174	0.026	0.135	0.025	0.105	0.016	0.071	7	3.98%
Xiapu(Huizhou)	0.079	0.242	0.065	0.220	0.043	0.213	0.023	0.204	0.017	0.157	--	--	35	29.66%
Jinguowan(Huizhou)	0.028	0.114	0.022	0.119	0.017	0.083	0.024	0.124	0.019	0.115	0.010	0.068	0	0.00%
Haogang(Dongguan)	0.058	0.266	0.039	0.207	0.019	0.152	0.034	0.183	0.024	0.131	0.028	0.106	10	6.06%
Zimaling Park(Zhongshan)	0.028	0.121	0.019	0.108	0.014	0.097	0.013	0.078	0.013	0.065	0.012	0.036	0	0.00%
Tsuen Wan(HKSAR)	0.039	0.141	0.028	0.130	0.028	0.121	0.038	0.102	0.021	0.119	0.024	0.060	0	0.00%
Tap Mun(HKSAR)	0.039	0.119	0.026	0.126	0.023	0.088	0.020	0.092	0.019	0.103	0.012	0.046	0	0.00%
Tung Chung(HKSAR)	0.038	0.155	0.027	0.144	0.024	0.192	0.023	0.096	0.015	0.116	0.013	0.042	3	1.67%

Table 3.4 c : The monthly averages and overall averages of Respirable Suspended Particulates

Month	Jan 2007	Feb	Mar	Apr	May	Jun	Overall Average
Luhu Park(Guangzhou)	0.101	0.093	0.063	0.091	0.059	0.065	0.078
Wanqingsha(Guangzhou)	0.131	0.083	0.078	0.077	0.073*	0.036*	0.084
Tianhu(Guangzhou)	0.071	0.073	0.047	0.068	0.066	0.057	0.064
Liyuan(Shenzhen)	0.087	0.068	0.047	0.056	0.066	0.032	0.059
Tangjia(Zhuhai)	0.042*	0.035	0.035	0.038	0.055*	0.024	0.037
Shunde Dangxiao(Foshan)	0.161*	0.120	0.100	0.106	0.098	0.062	0.105
Huijingcheng(Foshan)	0.207	0.147	0.159	0.143	0.074	0.116	0.139
Donghu(Jiangmen)	0.087	0.058	0.054	0.065	0.061	0.037	0.059
Chengzhong(Zhaoqing)	0.091	0.082	0.066	0.065	0.062	0.037	0.066
Xiapu(Huizhou)	0.153	0.145	0.110	0.083	0.063*	--	0.120
Jinguowan(Huizhou)	0.074*	0.059	0.048	0.063	0.059	0.033	0.056
Haogang(Dongguan)	0.125	0.097	0.066	0.079	0.074	0.054	0.082
Zimaling Park(Zhongshan)	0.068	0.041*	0.031	0.035	0.038	0.027*	0.044
Tsuen Wan(HKSAR)	0.088	0.066	0.055	0.061	0.061	0.034	0.061
Tap Mun(HKSAR)	0.077	0.060	0.046	0.054	0.056	0.022	0.052
Tung Chung(HKSAR)	0.098	0.063	0.051	0.057	0.052	0.022	0.057

Remark : 1. All concentration units are in milligrams per cubic metre.

2. "*" denotes that the data capture rate does not meet the minimum requirements for determining a representative value.

3. "--" denotes lack of data due to equipment maintenance works.

3.5 Monthly Variations of Pollutant Concentrations

Figure 6 shows the monthly variations of the major pollutants (SO_2 , NO_2 , O_3 , and PM_{10}) recorded by the Network during the period from January to June 2007. The overall concentrations of SO_2 , NO_2 and PM_{10} were generally higher in the winter month of January and lower in June as summer approached. This was due to the heavier rainfall and higher mixing layer height in the summer months. Apart from heavier rainfall and higher mixing layer which favoured the dispersion of pollutants, the relatively clean maritime air stream prevailed in the PRD region under the influence of southern monsoon also accounts for a lower level of pollution in summer time. As for ozone, the variation of monthly average during the first half of 2007 was relatively small as the highest monthly average usually occurred in autumn. That notwithstanding, the monthly variations in concentrations of pollutants may change from year to year. Long-term monitoring is thus required before a general pattern can be established.




Figure 6 : Monthly variations of average pollutant concentrations measured by the Network

4. Statistical Analysis of the Regional Air Quality Index (RAQI)

The two governments of Guangdong and HKSAR jointly started reporting the RAQI since 30 November 2005 to provide the public with information about the air quality in different parts of the PRD region.

The RAQI is a composite indicator of the aggregate level of the four major regional air pollutants, namely sulphur dioxide (SO₂), nitrogen dioxide (NO₂), ozone (O₃) and respirable suspended particulates (PM₁₀). The higher the index value, the higher the regional air pollution levels. The RAQI is divided into the following five grades :

Grade	Regional Air Quality Index (RAQI) value [#]	Air Quality Condition in the Monitored Area
I	0 – 1	Concentrations of all pollutants are well within Class 2 of the National Ambient Air Quality Standards (NAAQS)
II	1 – 2	Concentrations of all pollutants are generally within Class 2 NAAQS
III	2 – 3	Concentrations of individual pollutants may approach or exceed Class 2 NAAQS
IV	3 – 4	Class 2 NAAQS are generally exceeded
V	>4	Class 2 NAAQS are significantly exceeded

The formula for calculating the RAQI is as follows:

$$I_c = \sum_{i=1}^4 \frac{C_i}{R_i}$$

where I_c stands for the RAQI, an indicator of the aggregate pollution level of four pollutants, namely, SO₂, NO₂, PM₁₀ and O₃. For SO₂, NO₂ and PM₁₀, C_i is the daily average concentration while R_i represents the daily average concentration limits of the corresponding pollutants as specified in Class 2 NAAQS. For O₃, C_i is the highest hourly average of a day while R_i represents the hourly average concentration limit in Class 2 NAAQS (refer to “NAAQS (GB 3095 – 1996)” revised version).

[#] The upper limits of the value ranges for RAQI Grades I, II, III and IV are inclusive.

4.1 Statistics on RAQI Grades

Table 4.1a and Figure 7 summarise the statistics on the RAQI grades of all monitoring stations in the Network from January to June 2007. As shown in the table, the percentages of days with valid RAQI at various monitoring stations were quite high, averaging 91%. With continuous improvement in the operation of the Network, the overall percentage of valid RAQI is expected to further improve in the future.

Table 4.1 a : Statistics on RAQI grades of individual monitoring stations

Monitoring Stations	District	Days with valid RAQI	Distribution of RAQI grades (%) (January – June 2007)				
			Grade I	Grade II	Grade III	Grade IV	Grade V
Luhu Park	Guangzhou	169	13.61	36.69	39.05	8.28	2.37
Wanqingsha	Guangzhou	173	17.92	42.77	25.43	13.87	0.00
Tianhu	Guangzhou	170	27.06	51.76	17.06	2.94	1.18
Liyuan	Shenzhen	170	34.71	54.12	9.41	1.18	0.59
Tangjia	Zhuhai	148	34.46	50.00	14.19	0.68	0.68
Shunde Dangxiao	Foshan	161	4.97	47.20	31.68	9.32	6.83
Huijingcheng	Foshan	170	0.59	31.18	32.35	19.41	16.47
Donghu	Jiangmen	168	37.50	39.29	20.24	2.98	0.00
Chengzhong	Zhaoqing	172	19.19	50.58	18.60	11.05	0.58
Xiapu	Huizhou	162	22.22	58.02	18.52	1.23	0.00
Jinguowan	Huizhou	164	39.02	55.49	4.88	0.61	0.00
Haogang	Dongguan	166	6.63	46.99	33.13	12.05	1.20
Zimaling Park	Zhongshan	121	28.10	42.15	19.83	9.92	0.00
Tsuen Wan	HKSAR	171	19.88	67.25	12.28	0.58	0.00
Tap Mun	HKSAR	167	39.52	57.49	2.99	0.00	0.00
Tung Chung	HKSAR	170	40.00	43.53	15.29	1.18	0.00




Figure 7 : Stacked column chart of RAQI grades of individual monitoring stations

Figure 8 shows the overall distribution of different RAQI grades recorded by the Network during the period from January to June 2007. On the whole, 72.42% of the RAQI values are in Grade I or II, meaning the pollutant concentrations are within Class 2 NAAQS, followed by 19.72% in Grade III, 5.95% in Grade IV and 1.91% in Grade V.

**Distribution of RAQI Grades
(January - June 2007)**




Figure 8 : Distribution of RAQI grades in the Network

4.2 Spatial Distribution of Average RAQI Grades

Figure 9 shows the spatial distribution of RAQI overall average grades during the period from January to June 2007. It can be seen that average RAQI values measured in most parts of the PRD region were in Grade II while the average values in the middle part of the region were in Grade III, similar to the corresponding period last year. Average RAQI values measured at individual monitoring stations in the Network are shown in Figure 10.

**Distribution of average RAQI
from Jan 2007 to Jun 2007**




Figure 9 : Spatial distribution of average RAQI grades in the Network




Figure 10 : The average RAQI of individual monitoring stations

4.3 Monthly Variations of Average RAQI

Figure 11 shows the monthly variations in the average RAQI values of the Network from January to June 2007. Except for January where average RAQI value reached the Grade III level, the values in other months were within the Grade II category. The highest and lowest monthly averages of RAQI were recorded in January and June, respectively.




Figure 11 : Monthly variations in average RAQI

Annex A : Site Information of Monitoring Stations

Monitoring Stations	Address	Area Type	Sampling Height (Above P.D.)	Above Ground	Date Commenced Operation
Luhu Park (Guangzhou)	Inside Jufong Garden of Luhu Park (Big yard, No. 11 Luhu Park)	City	30m	9m	1993
Wanqingsha (Guangzhou)	Wanqingsha Secondary School, Nansha	Mixed educational/commercial and residential/industrial	13m	12m	Oct 2004
Tianhu (Guangzhou)	Tianhu Park, Conghua City	Background : rural	251m	13m	Oct 2004
Liyuan (Shenzhen)	Shennan Zhong Road, Shenzhen City	City	38m	12m	Sep 1997
Tangjia (Zhuhai)	Building No. 1, Rong Yuan, Zhongshan University, Tangjia, Zhuhai City	Mixed educational/commercial and residential/industrial	24m	19m	Jan 2003
Shunde Dangxiao (Foshan)	Roof-top of Educational Building, Foshan City Communist Party Shunde	Tourist and cultural/educational	27m	17m	Oct 1999
Huijingcheng (Foshan)	No. 127, Fenjiang Nan Road, Chancheng Area	Urban: mixed residential/commercial/industrial	24m	14m	Feb 2000
Donghu (Jiangmen)	Inside Donghu Park, Jiangmen City	City	17.5m	5m	Nov 2001
Chengzhong (Zhaoqing)	No. 17, Qintian Road, Zhaoqing City	Urban: mixed residential/commercial	21m	16m	Jun 2001
Xiapu (Huizhou)	No. 4 Xiabuhengjiang Road No. 3, Huicheng Area	Urban: commercial	49m	20m	Dec 1999
Jinguowan (Huizhou)	Jinguowan Ecological Farm, Huizhou City	Residential	77m	8m	Oct 2004
Haogang (Dongguan)	Haogang Primary School, NanchengQu, Dongguan City	Mixed residential/commercial/industrial	18m	14m	1998
Zimaling Park (Zhongshan)	Zimaling Park, Zhongshan City	Mixed residential/commercial	45 m	7m	Aug 2002
Tsuen Wan (HKSAR)	60 Tai Ho Road, Tsuen Wan	Urban: mixed residential/commercial/industrial	21m	17m	Aug 1988
Tap Mun (HKSAR)	Tap Mun Police Station	Background: rural	26m	11m	Apr 1998
Tung Chung (HKSAR)	6 Fu Tung Street, Tung Chung	New Town: residential	34.5m	27.5m	Apr 1999

Annex B : Measurement Methods of Air Pollutant Concentration

Pollutants	Measuring Principles
Sulphur Dioxide	UV fluorescence / Differential Optical Absorption Spectroscopy
Nitrogen Dioxide	Chemiluminescence / Differential Optical Absorption Spectroscopy
Ozone	UV absorption / Differential Optical Absorption Spectroscopy
Respirable Suspended Particulates	Oscillating microbalance (TEOM) Beta particulate monitor